1985
NEW MEXICO
SELECTED HEALTH STATISTICS


Rx
HEALTH PROMOTION
DISEASE PREVENTION

THE RIGHT PRESCRIPTION

State of New Mexico HEALTH and ENVIRONMENT DEPARTMENT Health Services Division

STATE OF NEW MEXICO

1912

GARREY CARRUTHERS
Governor

NEW MEXICO
HEALTH AND ENVIRONMENT
DEPARTMENT

LARRY GORDON
Secretary for Health & Environment

MARY LOU MARTINEZ
Director
Health Services Division

SAM CULBERTSON
State Registrar
Vital Statistics

PUBLISHED JULY 1987

SELECTED HEALTH STATISTICS
NEW MEXICO 1985

This report of New Mexico Health Statistics gives an overview of selected vital and health statistics for calendar year 1985. To portray the patterns of temporal change, data from earlier years have also been included. The primary sources of these summary data are the reportable certificates of birth and death and the federal and state reporting systems.

Unless otherwise indicated, the tabulations of vital events are by place of residence. Thus, births and deaths of New Mexico residents that occurred in other states are included in the tabulations, while the vital events that occurred in New Mexico to non-residents are not included. Tabulations are based on 1985 birth and death certificates received in the vital records office through May 15, 1986. Extending the cut-off date so late into the following year allows for the inclusion of more than 99% of the vital events occurring in 1985 into the tabulations.

The following map depicts the alignment of New Mexico counties into five Health Service Districts. The majority of county data presented in this report are grouped according to these districts.


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
1
2
MORA
HARDING
LOS ALAMOS
3
SANDOVAL
SANTA FE
McKINLEY
SAN MIGUEL
BERNALILLO
CIBOLA
VALENCIA
TORRANCE
GUADALUPE
QUAY
CURRY
DE BACA
ROOSEVELT
CATRON
SOCORRO
4
LINCOLN
CHAVES
5
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO

Comments on the content and format of this report, or requests for additional data or analysis should be directed to:

Health & Environment Department
Health Services Division
Vital Statistics Section
P.O. Box 968—Harold Runnels Bldg.
Santa Fe, New Mexico 87504-0968

REPORT PREPARED BY THE HEALTH STATISTICS SECTION
Tony Ortiz, Supervisor

In addition to the statistical tabulation and analysis, the staff is responsible for the review and query of the source documents; the coding and data preparation for submission to key entry; the maintenance of the statistical data file; and the graphical layouts of the report.

ACKNOWLEDGEMENTS

Grateful acknowledgement is made to the administration of the Health and Environment Department for its support, as well as to the various sections of the department for their support and help. Particular thanks to the Vital Records Section for its registration of the documents; to the Automated Data Processing Bureau for their programming support; and to the sections of the Adult Health and Nutrition Bureau, the Office of Epidemiology, and the Division of Health Planning and Development, for their cooperation and help with various aspects of the report.

Acknowledgement and thanks are also extended to the various State and Federal agencies whose data aided in the preparation of this report. Among these agencies are: Human Services Department, Cooperative Extension Service, New Mexico Tumor Registry, Bureau of Business and Economic Research, National Center for Health Statistics, Bureau of the Census and the Centers for Disease Control.

TABLE OF CONTENTS

HIGHLIGHTS

New Mexicans on the Average 1
New Mexico Health Statistics 2

DEFINITIONS 3

INTRODUCTION

POPULATION 4
INCOME 11
MARRIAGES/DIVORCES 13
ABORTIONS 14

NATALITY

FERTILITY 18
NATALITY BY AGE OF MOTHER 20
LIVE BIRTH ORDER 22
BIRTHS TO SINGLE MOTHERS 24
LOW BIRTH WEIGHT 27
GESTATIONAL AGE 29
CONGENITAL ANOMALIES 30
PARENT'S LEVEL OF EDUCATION 32
MULTIPLE BIRTHS 32
HOSPITAL/ATTENDANT 32
LEVEL OF PRENATAL CARE 33

MORTALITY

ALL CAUSES 37
LIFE EXPECTANCY 38
LEADING CAUSES OF DEATH 38
CAUSES BY AGE 42
CAUSES BY COUNTY 46
STANDARDIZED MORTALITY RATIOS 50
YEARS OF POTENTIAL LIFE LOST 52
INFANT MORTALITY 53
MATERNAL MORTALITY 58
FETAL MORTALITY 59

COMMUNICABLE DISEASES

VENEREAL DISEASE 62
PLAGUE 64
RABIES 65
DIPHTHERIA 65
TUBERCULOSIS 66
RUBELLA 67
AIDS 67
MEASLES 68
OTHER COMMUNICABLE DISEASES 68

SPECIAL FEATURES

NEW MEXICO COUNTY HEALTH PROFILES 70

APPENDIX 137

NEW MEXICANS

ON THE AVERAGE

12 PERSONS PER SQUARE MILE

MARRIAGE



40 MARRIAGES A DAY

INCOME

PER CAPITA
INCOME 10,741

1 out of 7 families below poverty
1 out of 6 persons below poverty

DIVORCE



25 DIVORCES A DAY

MEDIAN AGE 29.0

10 OUT OF 100 ARE 65 YEARS OF AGE OR OLDER
10 OUT OF 100 ARE LESS THAN 5 YEARS OF AGE
16 OUT OF 100 ARE 5-14 YEARS OF AGE
34 OUT OF 100 ARE 15-34 YEARS OF AGE
30 OUT OF 100 ARE 35-64 YEARS OF AGE

7 out of 100 are Indian
2 out of 100 are Black
4 out of 10 are Hispanic White
5 out of 10 are Non-Hispanic White

BIRTHS

75 BIRTHS A DAY

1 out of 6 to teen mothers
1 out of 3 to women 20-24
1 out of 4 to single women
12 out of 1000 reported congenital anomalies
11 maternal deaths for every 100,000 live births
6 fetal deaths for every 1000 live births
1 infant death before age one for every 100 live births
13 abortions a day - 1 out of 3 to women 20-24

40 out of 100 were first born
1 out of 14 were low birth weight
88 out of 100 born in a hospital

DEATHS

26 DEATHS A DAY

7 a day from heart disease
5 a day from cancer
2 a day from accidents
2 a day from stroke
1 a day from suicide or homicide

LIFE EXPECTANCY

79.0 YEARS FOR FEMALES

70.3 YEARS FOR MALES

HIGHLIGHTS NEW MEXICO HEALTH STATISTICS 1985

POPULATION	As the ninth fastest growing State between 1980-1985, New Mexico reached a 1985 population total of 1,450,000 persons ranking it 37th in total population and seventh lowest in population density. New Mexico's population remains relatively young; however, the percentage of population 65 and older is increasing at a rate over ⅔ faster than nationally - a rate of growth exceeded in only five states. Projections indicate a continued increase in the percentage of older population in the State.
INCOME	New Mexico's per capita personal income of $10,741 was 20% lower than the national figure and ranked New Mexico 41st among the states. 14% of New Mexico families are below poverty level, ranking New Mexico sixth highest among the States.
MARRIAGE/DIVORCE	The State's marriage rate and divorce rate both continued dropping. New Mexico's marriage rate was just 1% lower than nationally for 1985, while the divorce rate was 24% higher.
ABORTIONS	4651 New Mexico resident women obtained legal induced abortions in 1985, a 13% decrease since 1980. 26% were under 20 years of age, and 54% were 20-29 years of age. 77% were single, 60% had had no previous abortions, and 47% obtained the abortion prior to the ninth week of gestation.
BIRTHS	New Mexico experienced an increasing birth rate from 1976 to 1980 followed by a slight general decrease in rate thru 1985. Despite the decrease, New Mexico's rate ranked fourth among the states, and exceeded the national rate by 20%. The number of births in the State in 1985 and the previous three years represent the highest numbers of resident annual births since 1963.
• **AGE OF MOTHER**	Birth rates are higher in New Mexico than nationally for all age groups of women ages 15 and older. In 1985 the State showed general increases for younger women and decreases for women over 24. The highest rate is for women 20-24. The birth rate for teenage women in the State remained 20% higher than nationally in 1984.
• **LIVE BIRTH ORDER**	40% of 1985 births to New Mexico residents were first order births. The general long-term trend has been toward a greater percentage of first and second order births and a decreased percentage of higher order births in the State.
• **BIRTHS TO SINGLE MOTHERS**	26.4% of live births in New Mexico during 1985 were to single mothers. The percentage of births to single mothers is highest for teenagers, but it is for mothers age 20 and older that percentages in New Mexico exceed national percentages.
• **LOW BIRTH WEIGHT**	The incidence of low birth weight babies in New Mexico dropped eight percent from 1984. In 1984 the incidence in the State was 12% higher than nationally, and is higher for younger and older women and lowest for women 25-34.
• **MULTIPLE BIRTHS**	There were 418 live births in multiple deliveries. New Mexico's multiple birth ratio of 15 live births in multiple deliveries per 1,000 total live births in 1985 was lower than the national ratio of 20 for 1984.
• **CONGENITAL ANOMALIES**	Congenital anomalies of varying degrees of severity were reported for 1.35% of resident live births in New Mexico (1981-85).
MORTALITY	Because of a younger population structure, New Mexico's crude death rates are well below the national figures: age-adjustment of the rates reveals only slightly lower overall mortality rates for New Mexico than nationally for both males and females. Mortality in New Mexico is generally higher than nationally for ages less than 45 years of age and lower for older ages.
• **LIFE EXPECTANCY**	Life expectancy at birth for New Mexico males is the same as nationally, and is 1.1 years more for New Mexico females than for females nationally. In New Mexico, females have a life expectancy 8.7 years greater than do males.
• **LEADING CAUSES**	Heart disease, cancer and accidents are the leading causes of death in New Mexico. Mortality from heart disease and cancer is lower in New Mexico than nationally, but accidents account for two times the percentage of deaths in New Mexico as nationally. Mortality in the State decreased in 1985 for heart disease, cancer, accidents, stroke and influenza/pneumonia, while increasing for COPD, suicide, diabetes, cirrhosis, homicide and alcoholism. The mortality rates for accidents, diabetes, suicide, homicide, COPD, cirrhosis and alcoholism notably exceeded national rates for these causes of death. Nearly twice the percentage of deaths occurred to persons under 35 years of age in New Mexico than nationally. Accidents are the leading cause of death for New Mexicans age 1-44. Homicide and suicide rank next as causes of death for New Mexicans 15-24 years of age.
INFANT MORTALITY	In 1985 New Mexico's infant mortality rate was the same as the national rate at 10.6, resulting from a ten percent increase over the State's previous year all time low of 9.6. New Mexico's neonatal mortality rate was 9% lower than nationally, while the postneonatal component of infant mortality in the State was 19% higher.
MATERNAL MORTALITY	3 maternal deaths were reported in 1985 for a rate of 10.9 maternal deaths per 100,000 live births as compared to 9.3 nationally. This is the same number of maternal deaths for the State reported in 1980, 1982, and 1983.
FETAL MORTALITY	157 resident fetal deaths were reported in 1985, for a fetal death ratio of 5.7 fetal deaths per thousand live births; a 16% decrease in the number from the previous year. The State's fetal mortality ratio remains lower than nationally for all ages of mothers.
COMMUNICABLE DISEASES:	
• **GONORRHEA**	The number of reported cases in the State decreased for the ninth successive year; since 1978 New Mexico's case rate has been below the national rate.
• **SYPHILIS**	The number of cases of infectious syphilis increased in 1985 by 25% over the total cases for the prior year and the reported cases of non-infectious syphilis increased by 31%.
• **PLAGUE**	An outbreak of 14 cases were reported in 1985, the third highest number ever reported in the State.
• **TUBERCULOSIS**	The State's incidence rate for tuberculosis decreased slightly from 7.9 cases per 100,000 population in 1984 to 6.5 in 1985 and had a 16% lower than incidence nationally in 1984.
• **OTHER**	There was one [redacted] reported case of rubella in 1984 and two [redacted] reported in 1985. No cases of diphtheria have been reported in New Mexico the last eight years. After a minor outbreak of measles in 1979 (38 cases) the number of reported cases had dropped dramatically during the last four years, but in 1984 soared to 88 cases, the highest since 1977, but dropped back to only eight cases in 1985.

DEFINITIONS

RESIDENCE DATA—Data allocated by place of residence of the decedent (deaths) or by place of residence of the child's mother (births, fetal deaths).

OCCURRENCE DATA—Data allocated by place where event occurred, regardless of the person's place of residence.

LIVE BIRTH—A birth that shows any sign of life after delivery.

LOW BIRTH WEIGHT—A live birth weighing less than 2500 grams (5½ pounds).

FETAL DEATH—A fetal death is defined as a death in utero (in the womb prior to birth), sometimes referred to as "stillbirths". New Mexico statute requires the reporting of all fetal deaths weighing 500 or more grams. Prior to 1980, the criteria had been 20 or more weeks of gestation.

INFANT DEATH—Death of an infant under one year of age (neonatal + postneonatal).

NEONATAL DEATH—Death of a newborn under 28 days of age.

POSTNEONATAL DEATH—Death of an infant 28 days of age and older, but not yet having attained one year of age.

MATERNAL DEATH—Death attributed to complications of pregnancy, childbirth or the puerperium.

PUERPERIUM—Period after delivery of placenta, which extends until the organs of reproduction have returned to normal, usually 6 to 8 weeks.

NATURAL INCREASE—Excess of births minus deaths.

(CRUDE) BIRTH RATE—Number of live births per 1,000 population.

FERTILITY RATE—Number of live births per 1,000 women 15-44 years of age.

FETAL DEATH RATIO—Number of fetal deaths per 1,000 live births.

INFANT DEATH RATE—Number of infant deaths per 1,000 live births.

NEONATAL DEATH RATE—Number of neonatal deaths per 1,000 live births.

POSTNEONATAL DEATH RATE—Number of postneonatal deaths per 1,000 live births.

MATERNAL DEATH RATE—Number of maternal deaths per 100,000 live births.

(CRUDE) DEATH RATE—Number of deaths per 1,000 population.

AGE-ADJUSTED DEATH RATE (Direct Method)—Elimination of the effects of difference in age composition by adjustment of the rate to a standard.

INTRODUCTION

POPULATION

The total United States population for 1985 was estimated to be 238,740,000 persons. This total represents a 5.4% increase in the nation's population since 1980-an average yearly growth of 1.1%. This rate of growth continues into the Eighties the average yearly growth experienced between 1970 and 1980 which had resulted in a total population increase of 11.4% during those ten years. This rate is lower than the 13.4% change from 1960 to 1970, and considerably lower than 18.5% change from 1950 to 1960.

The internal population shifts and growths of the various regions in the country have been notable and varied. Between 1970 and 1980, with a 37.1% increase, the Mountain Division was the fastest growing in the nation by far. The Seventies was a period of tremendous growth for the Mountain Division, exceeding the 35.1% increase of the Fifties. So far into the Eighties, the Mountain Division continues to be the fastest growing division having increased 12.5% in 1980-1985. The second fastest for 1980-1985 was the West South Central Division with an 11.6% increase (Figure 1.1).

The Pacific Division, including Alaska and Hawaii, was third fastest growing in 1980-1985 with an overall increase of 10.2%.

The East North Central Division lost population as two of the five States of that region recorded population decreases in 1980-1985. The Middle Atlantic Division rebounded slightly after experiencing an overall population decrease in 1970-1980.

FIGURE 1.1
UNITED STATES
PERCENTAGE INCREASE IN POPULATION BY DIVISION
1980-1985



Alaska, Arizona, Nevada and Florida have been the nation's fastest growing states of the Eighties. Nevada, Arizona and Florida were the fastest growing states in 1970-1980. During the first half of the Eighties, New Mexico has been the 9th fastest growing State, as it had been during the Seventies. New Mexico's average yearly growth of 2.8% during the Seventies has dropped to 2.3% so far in the Eighties (Table 1.1).

New Mexico's population has continued to increase over the years (Figure 1.2), however, the rate of growth and the components of growth have shown notable variation. From 1920 to 1930, the State's population increased 17.5%; 1930 to 1940 - 25.8%; 1940 to 1950 - 28.0%; 1950 to 1960 - 39.6%; 1960 to 1970 - 6.9%; and from 1970 to 1980, the population increased 28.1%. The State's population reached a 1980 census count of 1,302,894 persons, increasing to an estimated 1,450,000 persons in 1985.

TABLE 1.1
TWELVE FASTEST GROWING STATES
1980-85 AND 1970-80

RANK	STATE	PERCENT INCREASE 1980-1985	STATE	PERCENT INCREASE 1970-1980
1	Alaska	29.7	Nevada	63.8
2	Arizona	17.2	Arizona	53.1
3	Nevada	16.9	Florida	43.5
4	Florida	16.6	Wyoming	41.3
5	Texas	15.0	Utah	37.9
6	Utah	12.6	Alaska	32.8
7	Colorado	11.8	Idaho	32.4
8	California	11.4	Colorado	30.8
9	New Mexico	11.3	New Mexico	28.1
10	Georgia	9.4	Texas	27.1
11	Hawaii	9.2	Oregon	25.9
12	Oklahoma	9.1	Hawaii	25.3

SOURCE: Bureau of Census

FIGURE 1.2
NEW MEXICO
POPULATION COUNTS, ESTIMATES, PROJECTIONS
1920-2000



CENSUS YEARS 1920-1980 TAKEN FROM BUREAU OF CENSUS COUNTS. 1985 ESTIMATE FROM CURRENT POPULATION REPORTS P-25 BUREAU OF CENSUS, 1990, 2000 PROJECTIONS, NEW MEXICO POPULATION BY COUNTY, 1980-2005, BBER, NOV. 1986

In the Fifties, New Mexico experienced its highest percentage growth, mainly due to a high rate of natural increase *(excess of births over deaths)* coupled with a high rate of in-migration (Table 1.2).

During the Sixties, the State's growth continued but at a much slower rate due in part to a lower rate of natural increase, reflecting the lower birth rates of the period, but also reflecting the impact of an out-migration experienced in New Mexico during the Sixties.

The Seventies was a period of renewed growth for the State. As birth rates in the State have been lower than those previous periods, the natural increase component of the State's growth for 1970-1980 has decreased from previous periods, less than half of what it was for 1950-1960. However, a markedly increased in-migration component of population growth resulted in a 28.1% population increase in New Mexico from 1970 to 1980. Net migration accounted for half of the growth of the Seventies.

TABLE 1.2
NEW MEXICO
COMPONENTS OF POPULATION CHANGE
1950-1985

	NET CHANGE		COMPONENTS OF CHANGE (%)	
TIME PERIOD	NUMBER (Persons)	PERCENT	NATURAL INCREASE	NET MIGRATION
1980-85	147,106	11.3	7.3	4.0
1984-85	24,000	1.7	1.3	0.4
1983-84	27,000	1.9	1.3	0.6
1982-83	32,000	2.3	1.4	1.0
1981-82	33,000	2.5	1.4	1.1
1980-81	31,106	2.4	1.7	0.7
1970-80	285,839	28.1	14.0	14.1
1979-80	19,894	1.6	1.0	0.6
1978-79	29,000	2.3	1.3	1.0
1977-78	27,000	2.2	1.2	1.0
1976-77	31,000	2.6	1.2	1.4
1975-76	32,000	2.7	1.1	1.6
1974-75	34,000	3.0	1.2	1.8
1973-74	26,000	2.4	1.2	1.2
1972-73	26,000	2.4	1.2	1.2
1971-72	25,000	2.4	1.3	1.1
1970-71	35,945	3.5	1.8	1.7
1960-70	66,032	6.9	19.4	-12.4
1969-70	6,055	0.6	1.1	-0.5
1968-69	17,000	1.7	1.4	0.3
1967-68	-6,000	-0.6	1.3	-1.9
1966-67	-7,000	-0.7	1.5	-2.2
1965-66	-5,000	-0.5	1.6	-2.1
1964-65	6,000	0.6	1.9	-1.3
1963-64	17,000	1.7	2.1	-0.4
1962-63	10,000	1.0	2.2	-1.2
1961-62	14,000	1.5	2.4	-1.0
1960-61	13,977	1.5	3.2	-1.7
1950-60	269,836	39.6	29.7	9.9

Between 1980 and 1985, natural increase has been the major component of the State's growth accounting for 65% of that growth.

The effect of the changing components of New Mexico's population growth has been to create changes in the structural composition of the State's population. Because of the general decrease in birth rates, the increased life span, and the increased rate of the in-migration component of population change, the age distribution of New Mexico's population has changed over the years.

The most notable change is the increased percentage of older population. From 1950 to 1980 (Table 1.3), the percentage of the State's population 55 to 64 increased 34%, the percentage 65 and older increased 82%, while the percentage of population under 5 decreased by 37%.

With the exception of the 55-64 age group, population projections indicate a continuing increase in the percentage of older age groups in the State, while younger age groups comprise the decreasing percentages of New Mexico's population.

TABLE 1.3
NEW MEXICO POPULATION
PERCENTAGE DISTRIBUTION BY AGE GROUP
1950, 1960, 1970, 1980
PROJECTED 1990 and 2000

AGE GROUP	CENSUS	CENSUS	CENSUS	CENSUS	PROJECTED*		% CHANGE
	1950	1960	1970	1980	1990	2000	1980 TO 2000
< 5	13.9%	14.3%	9.5%	8.8%	9.4%	8.1%	- 8.0%
5-14	20.9	23.6	23.8	17.2	16.6	16.4	- 4.7
15-24	16.9	15.1	18.5	19.8	14.9	14.9	-24.7
25-34	15.8	13.9	12.3	16.6	16.6	13.5	-18.7
35-44	12.7	12.5	11.4	11.1	14.5	15.0	+35.1
45-54	8.9	9.2	10.0	9.4	9.4	12.5	+33.0
55-64	6.1	5.9	7.5	8.2	7.8	8.0	- 2.4
65 +	4.9	5.4	6.9	8.9	10.8	11.7	+31.5

SOURCE: Data based on Bureau of Census figures for Census years.

SOURCE OF PROJECTIONS: *"New Mexico Population Projections by County 1980-2005," Bureau of Business and Economic Research, Institute for Applied Research Services, University of New Mexico, Nov 86.

Figure 1.3 graphically illustrates the growth rates of various age groups for New Mexico and the U.S. populations for 1970-1980 and 1980-1985. New Mexico's 65 and older population was the fastest growing of any age group in 1970-1980 with a rate increase over 2 times greater than the national rate for this age group. The 18-44 age group ranked as the State's second fastest growing at a rate 64% higher than nationally. The rate of growth for New Mexico's population age 45-64 was over 4½ times the national rate for this age group in 1970-1980.

For the period 1980-1985 the 18-44 age group showed the greatest percentage increase in population, followed by the under five and 65 and older age groups. The increase in the less than 5 age group most likely reflects the increased birth rates of the early Eighties.

FIGURE 1.3
PERCENTAGE POPULATION CHANGE BY AGE GROUP
NEW MEXICO AND UNITED STATES
1970-1980 & 1980-1985



Please note Scale Differences

In spite of the changing age structure, as seen in Figure 1.4, New Mexico's population remains comparatively young.

Whereas 26% of the U.S. population in 1985 was under 18 years of age, 31% of New Mexico's population falls into that age group. The 1980 Census showed the median age in New Mexico to be 27.4 years as compared to 30.0 nationally. Only three States, (Utah, Alaska, Wyoming) showed a younger median age.

The percentage of population in 1985 that was 65 years of age and older is 18% lower in New Mexico than nationally (9.8% as compared to 12.0%). Only five states have proportionately fewer people 65 years of age and older than does New Mexico.

FIGURE 1.4
1985 POPULATION AGE DISTRIBUTION NEW MEXICO AND UNITED STATES



SOURCE: Bureau of the Census

The State's relatively young age structure is evident at the county level. In only six of the State's counties is the percentage of population under 20 years of age less than that for the nation as a whole according to 1985 estimates.

TABLE 1.4
NEW MEXICO POPULATION PERCENTAGE DISTRIBUTION BY AGE GROUP BY COUNTY
1985 ESTIMATES

HEALTH DISTRICTS	COUNTY	PERCENTAGE DISTRIBUTION BY AGE				
		< 5	5-19	20-44	45-64	65 +
	UNITED STATES	7.5%	22.0%	39.7%	18.8%	12.0%
	NEW MEXICO	9.7%	24.1%	39.1%	17.3%	9.8%
	CIBOLA	11.1	29.8	32.3	19.6	7.2
1	MCKINLEY	13.5	30.4	37.7	13.4	5.1
	SAN JUAN	13.2	27.6	38.0	14.7	6.5
	COLFAX	9.0	24.8	35.9	17.2	13.1
	HARDING	7.6	21.8	29.3	23.5	17.8
	LOS ALAMOS	5.9	23.2	37.3	27.0	6.5
	MORA	8.8	23.8	33.3	18.9	15.2
2	RIO ARRIBA	11.0	27.4	37.8	15.9	7.9
	SAN MIGUEL	10.2	25.1	38.4	14.9	11.4
	SANTA FE	8.4	22.6	41.7	17.3	10.0
	TAOS	10.2	25.7	37.6	16.8	9.7
	UNION	8.5	23.5	33.9	18.3	15.8
	BERNALILLO	8.5	21.6	43.2	17.5	9.2
	SANDOVAL	9.7	25.6	35.3	17.6	11.8
3	TORRANCE	8.7	26.4	32.4	18.9	13.6
	VALENCIA	9.8	25.6	35.9	18.1	10.6
	CHAVES	9.6	23.2	32.8	19.2	15.2
	CURRY	10.6	24.9	37.4	15.6	11.5
	DE BACA	8.3	19.4	29.7	20.5	22.1
	EDDY	10.0	24.9	34.0	17.7	13.4
4	GUADALUPE	9.8	26.3	34.0	19.1	10.9
	LEA	12.4	26.5	36.5	16.2	8.5
	LINCOLN	8.2	21.1	37.4	21.1	12.2
	QUAY	6.9	22.4	31.9	22.4	16.4
	ROOSEVELT	9.1	20.7	40.9	15.9	13.4
	CATRON	7.0	24.5	35.1	19.1	14.2
	DONA ANA	9.1	25.7	40.5	16.3	8.3
	GRANT	9.9	26.1	35.3	17.6	11.0
5	HIDALGO	10.8	28.4	33.9	17.0	9.9
	LUNA	8.3	21.7	27.8	21.1	21.1
	OTERO	11.0	24.1	41.2	16.4	7.3
	SIERRA	5.2	14.3	19.7	25.7	35.1
	SOCORRO	9.4	26.6	38.1	16.5	9.4

SOURCE OF PROJECTION: Projections of the population of New Mexico by county, 1980-2005; Bureau of Business and Economic Research, Institute for Applied Research Services, University of New Mexico. November 1986.

In three of New Mexico's counties (McKinley, San Juan, and Cibola) over 40% of each county's population is under 20 years of age (Table 1.4).

Only fourteen of New Mexico's counties exceeded the national percentage of population in the 65 and older age group. Over one-third of Sierra County's population is 65 years of age or older. McKinley County showed the smallest percentage, with just about 5% of the county population being 65 years of age or older.

Sandoval County showed the largest percentage population increase for the period 1970-1980 (Table 1.5), nearly doubling its population. San Juan and Valencia were the next fastest growing counties. Seven counties registered decreases in population for the period.

The components affecting the county population changes varied from county to county. In the fastest growing county (Sandoval) in-migration accounted for 81% of population growth, while in San Juan, the second fastest growing, migration accounted for 59% of growth. Other counties in which in-migration accounted for a substantial portion of growth were: Santa Fe, Bernalillo, Torrance, Valencia, Lincoln, Catron, Dona Ana, Luna, Sierra, and Socorro counties.

According to the 1980 Census New Mexico's population increased by 28.1% during 1970-80, an increase of 285,839 persons, resulting in a total population of 1,302,894. The state's overall in-migration accounted for 50.2% of the increase.

TABLE 1.5
NEW MEXICO POPULATION BY COUNTY
1980

HEALTH DISTRICTS	COUNTY	1980 CENSUS	1970 CENSUS	NET CHANGE 1970-1980		COMPONENTS OF CHANGE 1970-1980	
				NUMBER	PERCENT	NATURAL INCREASE* %	NET MIGRATION %
	NEW MEXICO	1,302,894	1,017,055	285,839	28.1	14.0	14.1
	CIBOLA	Cibola County included in Valencia County.					
1	MCKINLEY	56,449	43,208	13,241	30.6	25.9	4.7
	SAN JUAN	81,433	52,517	28,916	55.1	22.8	32.3
	COLFAX	13,667	12,170	1,497	12.3	7.7	4.6
	HARDING	1,090	1,348	- 258	-19.1	0.8	-19.9
	LOS ALAMOS	17,599	15,198	2,401	15.8	9.0	6.8
	MORA	4,205	4,673	- 468	-10.0	6.8	-16.8
2	RIO ARRIBA	29,282	25,170	4,112	16.3	16.7	- 0.4
	SAN MIGUEL	22,751	21,951	800	3.6	11.5	- 7.9
	SANTA FE	75,360	54,774	20,586	37.6	12.7	24.9
	TAOS	19,456	17,516	1,940	11.1	13.0	- 1.9
	UNION	4,725	4,925	- 200	- 4.1	3.6	- 7.7
	BERNALILLO	419,700	315,774	103,926	32.9	12.8	20.1
	SANDOVAL	34,799	17,492	17,307	98.9	18.9	80.0
3	TORRANCE	7,491	5,290	2,201	41.6	8.7	32.9
	VALENCIA	61,115	40,576	20,539	50.6	17.4	33.2
	CHAVES	51,103	43,335	7,768	17.9	8.2	9.7
	CURRY	42,019	39,517	2,502	6.3	17.8	-11.5
	DE BACA	2,454	2,547	- 93	- 3.7	0.1	- 3.8
	EDDY	47,855	41,119	6,736	16.4	9.7	6.7
4	GUADALUPE	4,496	4,969	- 473	- 9.5	8.1	-17.6
	LEA	55,993	49,554	6,439	13.0	13.9	- 0.9
	LINCOLN	10,997	7,560	3,437	45.5	9.8	35.7
	QUAY	10,577	10,903	- 326	- 3.0	4.6	- 7.6
	ROOSEVELT	15,695	16,479	- 784	- 4.8	8.4	-13.2
	CATRON	2,720	2,198	522	23.7	8.3	15.4
	DONA ANA	96,340	69,773	26,567	38.1	16.6	21.5
	GRANT	26,204	22,030	4,174	18.9	14.5	4.4
5	HIDALGO	6,049	4,734	1,315	27.8	17.2	10.6
	LUNA	15,585	11,706	3,879	33.1	11.2	21.9
	OTERO	44,665	41,097	3,568	8.7	17.9	- 9.2
	SIERRA	8,454	7,189	1,265	17.6	- 7.6	25.2
	SOCORRO	12,566	9,763	2,803	28.7	13.6	15.1

*/Excess of births minus deaths; net migration is the difference between net change and natural increase.

SOURCE: BUREAU OF CENSUS, CURRENT POPULATION REPORT

Forty percent of the State's counties experienced a net out-migration during 1970-80. In counties such as Harding (with the largest overall percentage population loss), the natural increase component was not sufficiently large to negate the loss due to out-migration.

Eleven of New Mexico's counties exceeded the overall State growth rate during this period. In Sandoval County the rate was more than triple the State average for 1970-1980.

Though not the fastest growing Bernalillo County remained by far the State's most populous county followed by Dona Ana, San Juan and Santa Fe. In all, seven counties exceeded 50,000 in population accounting for 64% of the state population in 1980. Harding County was the least populated of New Mexico's counties (Table 1.6).

Between 1980 and 1985, an additional increase of 147,106 persons was added to the State's population bringing the 1985 total population estimate to 1,450,000. Sandoval County followed closely by Lincoln County is the fastest growing of the State's counties in the Eighties.

In-migration accounted for 35% of the overall New Mexico population increase during this period. For the State's fastest growing counties - Sandoval, Lincoln, Dona Ana, Valencia, and Torrance - in-migration played a major role; however in Lea County natural increase was the major component of growth.

Four New Mexico counties registered overall declines in population from 1980 to 1985 (Table 1.6).

TABLE 1.6
NEW MEXICO
ESTIMATES OF POPULATION BY COUNTY
1985

HEALTH DISTRICTS

COUNTY	1985 ESTIMATES	1980 CENSUS	NET CHANGE 1980-1985		COMPONENTS OF CHANGE 1980-1985	
			NUMBER	PERCENT	NATURAL INCREASE * %	NET MIGRATION %
NEW MEXICO	1,450,000	1,302,894	147,106	11.3	7.3	4.0
CIBOLA	25,300	30,346	-5,046	-16.6	7.7	-24.3
1 MCKINLEY	63,300	56,449	6,851	12.1	12.4	-0.3
SAN JUAN	91,100	81,433	9,667	11.9	12.3	-0.4
COLFAX	14,300	13,667	633	4.6	5.0	-0.4
HARDING	1,000	1,090	-90	-8.3	1.0	-9.3
LOS ALAMOS	18,400	17,599	801	4.6	4.1	0.5
MORA	4,600	4,205	395	9.4	5.3	4.1
2 RIO ARRIBA	32,400	29,282	3,118	10.6	8.1	2.5
SAN MIGUEL	24,800	22,751	2,049	9.0	6.8	2.2
SANTA FE	84,600	75,360	9,240	12.3	5.8	6.5
TAOS	21,900	19,456	2,444	12.6	8.2	4.4
UNION	5,100	4,725	375	7.9	3.0	4.9
BERNALILLO	464,300	419,700	44,600	10.6	6.1	4.5
3 SANDOVAL	44,100	34,799	9,301	26.7	8.1	18.6
TORRANCE	8,600	7,491	1,109	14.8	5.1	9.7
VALENCIA	35,900	30,768	5,132	16.7	7.4	9.3
CHAVES	56,500	51,103	5,397	10.6	5.7	4.9
CURRY	42,600	42,019	581	1.4	7.7	-6.3
DE BACA	2,400	2,454	-54	-2.2	0.9	-3.1
4 EDDY	52,500	47,855	4,645	9.7	6.3	3.4
GUADALUPE	4,400	4,496	-96	-2.1	5.6	-7.7
LEA	65,500	55,993	9,507	17.0	9.9	7.1
LINCOLN	13,900	10,997	2,903	26.4	6.2	20.2
QUAY	11,700	10,577	1,123	10.6	2.7	7.9
ROOSEVELT	16,500	15,695	805	5.1	5.4	-0.3
CATRON	2,800	2,720	80	2.9	2.5	0.4
DONA ANA	118,300	96,340	21,960	22.8	8.3	14.5
GRANT	27,300	26,204	1,096	4.2	6.7	-2.5
5 HIDALGO	6,200	6,049	151	2.5	7.7	-5.2
LUNA	17,500	15,585	1,915	12.3	3.8	8.5
OTERO	49,400	44,665	4,735	10.6	9.3	1.3
SIERRA	9,500	8,454	1,046	12.4	-3.1	15.5
SOCORRO	13,600	12,566	1,034	8.2	6.9	1.3

* Excess of births minus deaths; net migration is the difference between net change and natural increase.

State estimate is to the nearest thousand; county estimates to the nearest hundred.

The overall percentage increases in population by county for 1980-1985 are presented graphically in Figure 1.5.

Although the percentage of population increase as shown in Figure 1.5, does indicate the rate of change in a county's population in relation to its previous population, it does not provide an indication of the actual population involved.

Sandoval County, the fastest growing county in the State for the period 1980-1985 increased by 26.7% during that period, an actual increase of 9,301 persons. Dona Ana County, the third fastest growing increased in population by 22.8% for an actual population increase of 21,960 persons, over two times as many people as for Sandoval County.

The percentage change was 10.6%, but in actual number of people Bernalillo County registered the largest gain in the State during the period 1980-1985 (44,600 persons).

FIGURE 1.5
NEW MEXICO
PERCENTAGE CHANGE IN POPULATION BY COUNTY
1980-1985



Of the State's population increase of 147,106 persons during the period 1980-1985, Bernalillo County accounted for 30% of that growth. Dona Ana County received 15% of the State's total increase while San Juan County accounted for another 7%.

Figure 1.6, on the following page, graphically depicts the contribution by each of the counties toward the overall State increase. As noted Bernalillo, Dona Ana and San Juan counties were the largest contributors, together accounting for 52% of the State's population increase. These three counties plus Lea, Sandoval, Santa Fe, McKinley, Chaves, Valencia, and Otero, in that ranking, contributed 86% of the total population increase in New Mexico for the period 1980-1985.

As noted earlier, population changes are composed of two factors: natural increase and net migration. During 1980-

FIGURE 1.6
NEW MEXICO
PERCENTAGE DISTRIBUTION OF POPULATION INCREASE
1980-1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
LOS ALAMOS
MORA
HARDING
McKINLEY
SANDOVAL
SANTA FE
SAN MIGUEL
BERNALILLO
QUAY
CIBOLA
VALENCIA
TORRANCE
GUADALUPE
CURRY
DE BACA
CATRON
SOCORRO
ROOSEVELT
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
TOTAL STATE INCREASE 147,106 PERSONS
30.3
1.3-3.7
4.7-14.9
0.0-0.8
LOSS IN POPULATION

FIGURE 1.7
NEW MEXICO
PERCENTAGE DISTRIBUTION
OF NET MIGRATION
1980-1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
LOS ALAMOS
MORA
HARDING
McKINLEY
SANDOVAL
SANTA FE
SAN MIGUEL
BERNALILLO
QUAY
CIBOLA
VALENCIA
TORRANCE
GUADALUPE
CURRY
DE BACA
CATRON
SOCORRO
ROOSEVELT
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
TOTAL STATE
NET MIGRATION 52,625 PERSONS
26.2-36.0
2.5-4.7
5.4-12.3
0.0-1.6
NEGATIVE NET MIGRATION

1985, 52,625 persons were added to the State's population as a result of migration. Net migration accounted for 35% of the State's total population increase, as New Mexico's fastest growing counties relied heavily on in-migration for growth. In Sandoval County, the fastest growing county, migration accounted for 70% of the growth.

Bernalillo County, which accounted for 45% of New Mexico's in-migration for 1970-1980, accounted for 36% in 1980-1985. Dona Ana accounted for 27 % of migration in 1980-85. Sandoval County accounted for 12% and Santa Fe County accounted for 9% of the State's net migration.

Eleven of the State's counties experienced a net out-migration during the period 1980-1985. The largest loss of population due to out-migration during this period occurred in Cibola County.

Though eleven of the State's counties experienced a net out-migration during 1980-1985, only four counties experienced overall losses in population (Cibola, Harding, De Baca and Guadalupe) as the natural increase components in the other seven counties compensated for the losses due to out-migration.

All but seven of New Mexico's counties exceeded the national birth to death ratio for natural increase during this period—1.8 births for every death. Only in one county did the number of births not exceed the number of deaths (Figure 1.8), as natural increase ranged from .7 births for every death in Sierra County, to five births for every death in McKinley and San Juan counties. Overall, the State's ratio of births to deaths is 67% higher than nationally.

FIGURE 1.8
NEW MEXICO NATURAL INCREASE
BIRTH TO DEATH RATIO
(Number of Births for Every Death)
AVERAGE BY COUNTY
1980-1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
LOS ALAMOS
MORA
HARDING
McKINLEY
SANTA FE
SANDOVAL
SAN MIGUEL
BERNALILLO
QUAY
CIBOLA
GUADALUPE
VALENCIA
TORRANCE
CURRY
CATRON
DE BACA
ROOSEVELT
SOCORRO
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
STATE 3.0 BIRTHS FOR EVERY DEATH
4.0-5.2
1.1-1.9
3.0-3.9
≤1.0
2.0-2.9

Thus from 1980 to 1985, eleven of the State's counties have experienced a net out-migration, however natural increase has offset these losses in all but four counties.

In-migration, which accounted for half of the State's population growth for 1970-1980, has accounted for only 35% for 1980-1985, but has been a major factor in the State's fastest growing counties. The net result being a total State increase of 285,839 persons from 1970 to 1980 with an additional population increase of 147,106 persons during the period from 1980 to 1985.

County population projections to the year 2000 indicate that Sandoval, Dona Ana and McKinley counties will experience the fastest growth rates of any of the counties, 56.2%, 52.1% and 40.9% respectively. Bernalillo County will remain New Mexico's most populous county - surpassing the half million mark by 1990 (Table 1.7).

By 1990, San Juan County should approach the hundred thousand mark, with Santa Fe County exceeding 100,000 by the year 2000 at which time Dona Ana should exceed 170,000. Four counties (Curry, De Baca, Harding, and Union) are projected to experience overall population decreases between 1985 and the year 2000. All other counties are expected to show an increase, with Quay showing the smallest increase.

The State then is projected to be well past the 1½ million mark by 1990, and to be over 1.8 million people by the year 2000.

TABLE 1.7
NEW MEXICO
PROJECTED COUNTY POPULATIONS
1985-2000

HEALTH DISTRICTS	COUNTY	1985 ESTIMATE	1990 PROJECTION	2000 PROJECTION	% CHANGE 1985 TO 1990	TO 2000
	NEW MEXICO	1,450,000	1,583,500	1,853,300	9.2	27.8
	CIBOLA	25,300	26,600	31,100	5.1	22.9
1	MCKINLEY	63,300	70,900	89,200	12.0	40.9
	SAN JUAN	91,100	99,500	128,000	9.2	40.5
	COLFAX	14,300	15,000	16,600	4.9	16.1
	HARDING	1,000	1,000	900	0.0	-10.0
	LOS ALAMOS	18,400	18,800	20,800	2.2	13.0
	MORA	4,600	4,900	5,400	6.5	17.4
2	RIO ARRIBA	32,400	35,300	42,300	9.0	30.6
	SAN MIGUEL	24,800	27,000	31,300	8.9	26.2
	SANTA FE	84,600	90,300	102,600	6.7	21.3
	TAOS	21,900	23,600	28,000	7.8	27.9
	UNION	5,100	5,000	4,900	-2.0	-3.9
	BERNALILLO	464,300	510,200	588,500	9.9	26.7
	SANDOVAL	44,100	51,900	68,900	17.7	56.2
3	TORRANCE	8,600	9,400	11,400	9.3	32.6
	VALENCIA	35,900	39,400	48,600	9.7	35.4
	CHAVES	56,500	57,500	63,200	1.8	11.9
	CURRY	42,600	41,900	41,700	-1.6	-2.1
	DE BACA	2,400	2,400	2,300	0.0	-4.2
	EDDY	52,500	55,000	61,500	4.8	17.1
4	GUADALUPE	4,400	4,600	4,800	4.5	9.1
	LEA	65,500	66,300	77,100	1.2	17.7
	LINCOLN	13,900	15,600	18,000	12.2	29.5
	QUAY	11,700	11,800	11,800	0.9	0.9
	ROOSEVELT	16,500	17,200	18,200	4.2	10.3
	CATRON	2,800	3,000	3,300	7.1	17.9
	DONA ANA	118,300	144,000	179,900	21.7	52.1
	GRANT	27,300	29,500	32,200	8.1	17.9
5	HIDALGO	6,200	6,700	7,600	8.1	22.6
	LUNA	17,500	19,800	22,500	13.1	28.6
	OTERO	49,400	54,300	62,500	9.9	26.5
	SIERRA	9,500	10,200	10,500	7.4	10.5
	SOCORRO	13,600	15,100	18,100	11.0	33.1

SOURCE OF PROJECTIONS: Projections of the population of New Mexico by county, 1980-2005; Bureau of Business and Economic Research, Institute for Applied Research Services, University of New Mexico. November 1986.

The nation's most populous states in 1985 were California (26.4 million). New York (17.7 million), Texas (16.4 million), Pennsylvania (11.9 million), and Illinois (11.5). In general, the states of the eastern U.S. are more populous than those of the West, and though the Mountain Division was the fastest growing in the nation for 1970-1980 and for 1980-1985, only 5.4% of the total United States population resides in the Mountain Division of the country (Figure 1.9).

FIGURE 1.9
UNITED STATES POPULATION
1985 PERCENTAGE DISTRIBUTION
BY DIVISION



1985 RESIDENT U.S. POPULATION
238,740,000

Such a relatively small percentage of the nation's population residing in the vast expanse of the Mountain Division results in that division being the least densely populated in the nation, however Alaska remains the least densely populated state. New Mexico, in spite of its population growth, is one of the less densely populated Mountain States with a population density of about 12 persons per square mile (Figure 1.10). The Middle Atlantic Division is the most densely populated in the nation.

FIGURE 1.10
UNITED STATES
PERSONS PER SQUARE MILE*
BY DIVISION
1985



* BASED ON LAND AREA IN SQUARE MILES

New Mexico's county population densities ranged from a high of 397 persons per square mile in Bernalillo County to less than half a person per square mile in Catron County (Figure 1.11).

Only 9 of the State's counties exceeded the State overall average population density of 11.9 persons per square mile.

Los Alamos was the second most densely populated county, followed by Santa Fe County. Nearly half (16) of the counties have population densities less than half of the overall State figure.

FIGURE 1.11
NEW MEXICO
POPULATION DENSITY BY COUNTY
PERSONS PER SQUARE MILE
1985



Much of the State's population is classified as rural. According to the 1980 Census, in six of the State's counties 100% of the population is classified as rural. In only seven counties is 25% or less of the population classified as rural.

According to the 1980 Census there are in New Mexico only six cities with populations in excess of 30,000 people. Albuquerque by far and away the largest city in the State, had a 1980 population of 331,767 persons, increasing to over 350,000 for 1984; Santa Fe, Las Cruces, Roswell, Farmington and Clovis, make up the remainder of the cities exceeding 30,000, in 1980 with Hobbs joining the group for 1984. Of the cities listed in Table A-1 in the Appendix three had populations in the twenty thousands, one in the ten thousands, and eleven cities over five thousand but under ten thousand in population in 1980 (Table A-1 in the Appendix).

Ethnically, New Mexico's population is composed of four major components: non-Hispanic White comprise 52.6% of the State's population, the Hispanic White 36.3%, the Indian 8.1% and the Black 1.8% (Table 1.8).

As noted from the data of Table 1.8, some shift in the percentage composition of the State's ethnic makeup has occurred since 1970.

Examination of the data in Tables 1.8 and 1.9 indicate that: the Indian have been the fastest growing of the State's ethnic groups during 1970-80 due to a high rate of natural increase and now compose 8.1% of the State's population; the non-Hispanic White were second fastest growing comprising over 50% of New Mexico's population, though a comparatively low rate of natural increase component of growth is overshadowed by a large in-migration; the Black continue to comprise about 2% of the State's population maintaining this level through natural increase growth; the Hispanic White comprise 36% of the State's population, a percentage which has decreased through the Seventies, for though experiencing a high rate of natural increase, the Hispanic experienced only a 1.9% net in-migration increase (Table 1.9).

TABLE 1.8
NEW MEXICO
ETHNIC/RACIAL COMPOSITION
1970 CENSUS AND 1980 ESTIMATE

ETHNIC/RACE	PERCENTAGE	
	1970 CENSUS	1980 CENSUS
WHITE	90.1	88.9
Non-Hispanic	*50.0*	*52.6*
Hispanic	*40.1*	*36.3*
BLACK	1.9	1.8
INDIAN	7.2	8.1
OTHER	.8	1.1

* Ethnic breakdown extracted and adjusted from the Census by the New Mexico Tumor Registry.

TABLE 1.9
NEW MEXICO POPULATION GROWTH CHARACTERISTICS BY ETHNICITY 1970-1980

ETHNIC/RACE	NET POPULATION INCREASE 1970-80 %	NATURAL INCREASE		NET MIGRATION
		BIRTH/ 1 DEATH	%	
ALL RACES	28.1	2.8	14.0	14.2
WHITE	25.6	2.6	12.4	13.2
Non-Hispanic	28.9	1.8	7.5	21.4
Hispanic	21.0	4.2	19.1	1.9
BLACK	24.4	3.7	19.8	4.6
INDIAN	48.0	4.4	30.3	17.7

The Hispanic comprise a portion of all the counties' populations (11%-86%). Five counties have notably high percentages of Indian population, while the counties showing the highest percentages of Black population are mainly those of the southeastern corner of the State (Table 1.10).

The Hispanic, Black and Indian are young populations. While only 38% of New Mexico's non-Hispanic White population is under 25 years of age, this figure is 53% for the Hispanic, 54% for the Black, and 60% of the Indian population is under 25 years of age (Table 1.11).

TABLE 1.10
NEW MEXICO
RACIAL DISTRIBUTION BY COUNTY
1980

HEALTH DISTRICTS	COUNTY	WHITE Non-Hispanic	WHITE Hispanic	BLACK	INDIAN	OTHER
	NEW MEXICO	52.6%	36.3%	1.8%	8.1%	1.1%
	CIBOLA	Cibola County included in Valencia County.				
1	MCKINLEY	20.3	12.5	.6	65.7	.9
	SAN JUAN	54.6	11.4	.4	33.0	.5
	COLFAX	51.3	47.3	.4	.5	.5
	HARDING	55.1	44.3	--	.1	.5
	LOS ALAMOS	86.1	11.4	.4	.5	1.5
	MORA	12.9	86.4	0.0	.2	.4
2	RIO ARRIBA	14.5	73.3	0.1	11.4	.7
	SAN MIGUEL	17.3	81.3	.4	.4	.6
	SANTA FE	40.8	55.0	.5	2.8	.8
	TAOS	23.9	68.8	.2	6.4	.7
	UNION	68.3	31.2	--	.3	.3
	BERNALILLO	56.9	36.4	2.3	2.7	1.7
	SANDOVAL	43.9	27.3	.9	27.3	.7
3	TORRANCE	57.3	41.0	.2	.5	.9
	VALENCIA	41.2	43.5	.6	14.1	.6
	CHAVES	66.1	30.4	2.2	.6	.7
	CURRY	71.6	19.2	6.8	.4	1.9
	DE BACA	68.5	30.9	0.0	.1	.4
	EDDY	66.9	30.6	1.8	.3	.4
4	GUADALUPE	16.5	82.4	.1	.2	.7
	LEA	73.1	21.2	4.8	.5	.4
	LINCOLN	72.0	26.5	.5	.6	.5
	QUAY	62.2	35.3	1.3	.4	.8
	ROOSEVELT	75.3	21.4	1.3	.7	1.4
	CATRON	69.4	28.3	.4	1.5	.4
	DONA ANA	44.5	51.8	1.7	.8	1.3
	GRANT	47.4	51.2	.5	.5	.4
5	HIDALGO	52.0	47.0	.2	.1	.7
	LUNA	58.3	39.2	1.6	.5	.4
	OTERO	67.0	21.3	4.8	4.8	2.0
	SIERRA	73.9	25.2	.1	.6	.2
	SOCORRO	41.9	46.4	.4	9.9	1.3

Percentage distribution based upon 1980 census counts.

TABLE 1.11
NEW MEXICO
1980 RACE/ETHNIC POPULATIONS
Percentage Distribution by Age Group

AGE	ALL RACES	WHITE Non-Hispanic	WHITE Hispanic	Black	Indian	Other
TOTAL	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
< 5	8.8	6.8	10.7	10.4	13.1	10.2
5-14	17.2	13.7	20.6	19.0	23.7	16.6
15-24	19.8	17.5	22.1	24.6	23.2	20.8
25-34	16.6	17.2	15.9	16.7	15.1	23.6
35-44	11.1	12.1	10.2	9.1	9.4	12.0
45-54	9.4	10.7	8.2	8.4	6.4	8.4
55-64	8.2	10.5	6.0	5.8	4.3	4.4
65 +	8.9	11.5	6.3	6.1	4.8	4.0

Percentage distribution by age group for various race/ethnic groups based upon 1980 Census counts with adjustments for Hispanic White breakdown.

In the population pyramids, Figure 1.12, the shift toward a younger age among New Mexico Hispanic, Black and Indian is graphically evident.

At its 1985 level of 1,450,000 people, New Mexico, the fifth largest state in land area, ranks 37th among the states in total population and is the seventh lowest in population density.

FIGURE 1.12
POPULATION DISTRIBUTION BY AGE
NEW MEXICO 1980 CENSUS
UNITED STATES 1983 ESTIMATE



INCOME

Although monetary income is not the sole determinant of the quality of health, it is evident from various health statistics presented in reports for previous years, and in other studies, that low levels of income are associated with poor showings in various statistical health indicators. According to the DHEW report, *Health—United States 1978,* "Low income people in general have worse health than people of higher income."

Though New Mexico has in recent years shown improvement in such indices as personal income and number of families below poverty level, the State continues to compare poorly with other states.

New Mexico's personal income has increased over the years (Table 1.12); however, its ranking, compared to other states has varied, and based on latest revised figures, New Mexico now ranks forty-first among the states. New Mexico's 1985 per capita personal income figure of $10,741 was **20%** lower than the national figure of $13,451.

TABLE 1.12
PER CAPITA PERSONAL INCOME
NEW MEXICO AND UNITED STATES
1975-1985

YEAR	UNITED STATES	NEW MEXICO	
		INCOME	RANK
1985	$ 13451	$ 10741	41
1984	$ 12707	$ 10330	42
1983	$ 11687	$ 9656	41
1982	$ 11113	$ 9301	39
1981	$ 10544	$ 8780	40
1980	$ 9194	$ 7947	39
1979	$ 8651	$ 7207	40
1978	$ 7772	$ 6445	42
1977	$ 6984	$ 5745	44
1976	$ 6367	$ 5280	45
1975	$ 5842	$ 4785	47

Source: U.S. Dept. of Commerce

Figure 1.13 gives a general idea of the variation of per capita personal income throughout the nation by divisions based on 1985 data.

Alaska ranked highest among the states with a per capita income of $17,756, followed by Connecticut, New Jersey, Massachusetts and rounding off the top five - Maryland. The East South Central was lowest of any region; Mississippi was the lowest among the states, with a per capita personal income of $9,035, a level which New Mexico exceeded by 19%.

FIGURE 1.13
PER CAPITA INCOME 1985
BY DIVISIONS



U.S. DEPT. OF COMMERCE, BUREAU OF ECONOMIC ANALYSIS

1984 data shows only one of New Mexico's counties exceeded the national figure for per capita personal income (Figure 1.14). Over half of the counties registered a per capita personal income figure which was less than 75% of the national figure, and for four counties the figure was less than half that of the nation as a whole.

FIGURE 1.14
PER CAPITA PERSONAL INCOME
NEW MEXICO 1984
RANKING BY COUNTY



Los Alamos County had the highest per capita income in the state, which was 58% higher than nationally, nearly double the State figure, and 63% higher than Bernalillo County (the State's most populous county) which ranked second.

A greater percentage of persons in New Mexico are at or below poverty level compared to the total U.S. population.

According to the 1980 Census, 17.4% of the people in New Mexico are at or below poverty level compared to the na-

TABLE 1.13
PERCENTAGE OF PERSONS AT OR BELOW POVERTY LEVEL
BY ETHNIC/RACE
NEW MEXICO AND UNITED STATES
1980 CENSUS

ETHNIC/RACE	UNITED STATES	NEW MEXICO
TOTAL POPULATION	12.5%	17.4%
White	9.4	13.3
Spanish Origin	23.8	23.2
All Other	NA	37.5
Black	30.2	29.3
Indian	NA	40.2
Asian & Pacific Islander	13.9	23.7

tional figure of 12.5%. The percentage of persons in New Mexico at or below the poverty level is 39% greater than nationally (Table 1.13).

Table 1.13 further shows the high percentage of persons in New Mexico's various race/ethnic populations who are at or below poverty level. Over 40% of the Indian and nearly ¼ of the Spanish origin populations in New Mexico are at or below the poverty level.

Data by state for 1980 indicates that New Mexico ranks sixth highest among the states in percentage of families below the poverty level. The highest is Mississippi (19.5%) and the lowest Nevada and Wyoming, (5.9%). This is a decline from 1975, when data from the *"Survey of Income and Education"* ranked New Mexico second highest among the states. Despite the decrease, the State's percentage remains notably higher than the national rate.

FIGURE 1.15
NEW MEXICO FAMILIES BELOW POVERTY
PERCENTAGE BY COUNTY
RANKING BY COUNTY
1980 CENSUS



All but two of the State's counties exceed the national percent and twenty one exceed the overall state average for families below poverty level. In two of the counties in New Mexico (McKinley and Guadalupe) the percentage of families below poverty level triples the national average and in Mora County, the percentage is over four times the national average.

The five counties which in 1980 displayed the highest percentage of families below the poverty level in New Mexico are Mora, McKinley, Guadalupe, San Miguel and Rio Arriba (Figure 1.15).

Figure 1.15 presents a percentage ranking of New Mexico families below the poverty level by county. In 1980, 14.0% of the State's families were below poverty level, compared with 9.1% for the United States overall.

Los Alamos County showed the state's lowest percentage of families below the poverty level in 1980, with the percentage about a third of the national rate. The only other county in the State with a percentage below the national rate was Lea County at 9.0%. Bernalillo County, the State's most populous county showed 10.1% of the families below poverty level.

MARRIAGES AND DIVORCES

Nationally the marriage rate fell 3% between 1984 and 1985, from 10.5 to 10.2 per 1,000 population. The number of marriages fell between 1984 and 1985, from 2,487,000 to 2,425,000, a drop of 2%. The 1985 rate is the lowest since 1977, but the actual number of marriages was the fourth greatest annual total for the nation.

The Mountain Division reported the highest marriage rate nationally in 1985 (Figure 1.16) due mainly to Nevada's extremely high rate of 114.2 marriages per 1,000 population. All other states had marriage rates between 15.8 (South Carolina) and 7.5 (Pennsylvania).

National data indicate that remarriages comprise an increasing portion of all marriages. June is still the preferred month for tying the matrimonial knot nationally.

FIGURE 1.16
UNITED STATES
MARRIAGE RATE
BY DIVISION
1985



PROVISIONAL RATE PER 1,000 POPULATION

Nationally both the rate and number of divorces rose between 1984 and 1985: the rate increased by 2% to 5.0 per 1,000 population and the number by 3% to 1,187,000 which is the third highest total ever reported.

Nevada reported the nation's highest divorce rate (14.4). National data revealed the median duration of marriage prior to divorce to be 6.9 years; the median age at divorce was 31.7 for women and 34.3 for men; 46.5% of divorcing couples had no children under 18 years of age.

There has been a general decline in the rate of both marriage and dissolution of marriage in New Mexico in recent years (Table 1.14). In 1985, the State's marriage rate was 1% lower than nationally, while the divorce rate was 24% higher.

Regretably there is no centralized reporting of marriage or divorce data in New Mexico therefore data is limited and restricted to state totals.

TABLE 1.14
MARRIAGES AND DISSOLUTIONS OF MARRIAGE
Rates Per 1,000 Population
NEW MEXICO AND UNITED STATES
SELECTED YEARS 1940-85

YEAR	MARRIAGES		DISSOLUTIONS OF MARRIAGE	
	NEW MEXICO	U.S.	NEW MEXICO	U.S.
1985	10.1	10.2	6.2	5.0
1984	10.6	10.5	6.5	4.9
1983	12.0	10.5	6.5	5.0
1982	12.8	10.6	7.0	5.1
1981	12.9	10.6	7.7	5.3
1980	13.2	10.6	8.0	5.2
1979	13.6	10.4	8.0	5.3
1978	13.7	10.3	7.9	5.1
1977	13.8	9.9	7.7	5.0
1976	13.3	9.9	7.8	5.0
1975	13.8	10.0	7.4	4.8
1974	14.2	10.5	7.1	4.6
-				
-				
-				
1970	NA	10.6	NA	3.5
1965	NA	9.3	NA	2.5
1960	11.6	8.5	3.0	2.2
1955	28.4	9.3	2.7	2.3
1950	33.3	11.1	3.9	2.6
1945	22.6	12.2	6.3	3.5
1940	22.9	12.1	2.3	2.0

New Mexico data by occurrence
Rates per 1,000 population

SOURCE: NCHS-Monthly Vital Statistics Report

ABORTIONS

In New Mexico, reporting of legal induced abortion became law in 1977, and 1978 was the first full year of reporting. In 1978, 4,493 abortions were reported increasing yearly to 5,474 in 1981 and then decreasing to 5,010 in 1983 followed by a slight increase to 5,034 in 1984, and dropping by 8% to 4,651 abortions reported for 1985.

Consistently the abortion ratio has been highest for New Mexico's non-Hispanic White population and lowest for the State's Indian population (Table 1.15).

In 1985, about a third of New Mexico resident women obtaining abortions are age 20-24. The next largest percentage is to women 15-19, followed by women 25-29 years of age (Table 1.16). The percentage of abortions to women 30-39 has increased reflecting to some extent a general shift in the age distribution of the female population.

65% of New Mexico resident women obtaining abortions in 1985 reported a 12th grade education or higher (Table 1.17). The percentage with less than a 12th grade education was highest for Hispanic women.

The latest national data (1983) shows 1.27 million abortions reported nationally; the third consecutive year that the national abortion ratio has declined since reporting began and the first year in which the number of abortions actually declined.

TABLE 1.15
NEW MEXICO RESIDENTS
REPORTED LEGAL INDUCED ABORTIONS
1980-1985
Ratio Per 1,000 Live Births

RACE/ETHNIC	1980	1981	1982	1983	1984	1985
STATE TOTAL						
Abortions	5,373	5,474	5,213	5,010	5,034	4,651
Ratio	202.1	206.1	188.7	182.1	184.1	169.4
WHITE:						
Abortions	4,718	4,936	4,797	4,576	4,625	4,298
Ratio	210.9	222.0	208.6	200.3	203.6	190.0
Non-Hispanic						
Abortions	2,917	3,089	3,027	2,774	2,765	2,608
Ratio	273.9	292.3	273.5	263.6	267.3	254.4
Hispanic						
Abortions	1,801	1,847	1,770	1,802	1,860	1,690
Ratio	153.6	158.3	148.6	146.3	150.4	136.6
BLACK:						
Abortions	133	126	119	122	104	102
Ratio	233.7	197.5	165.3	172.6	162.5	151.1
INDIAN:						
Abortions	393	341	267	262	249	215
Ratio	116.2	101.3	74.8	72.5	68.0	56.5
OTHER:						
Abortions	61	17	26	42	56	36
Ratio	231.1	51.8	71.6	122.1	166.2	103.2
UNKNOWN:						
Abortions	68	54	4	8	0	0
Ratio*	*	*	*	*	*	*

*Ratio not computed

TABLE 1.16
NEW MEXICO RESIDENTS
REPORTED LEGAL INDUCED ABORTIONS
1980-1985
Percentage Distribution by Age Group by Race/Ethnic

		WHITE				
AGE GROUP	YEAR	NON HISPANIC	HISPANIC	BLACK	INDIAN	TOTAL
<15	1980	0.5	0.8	0.8	0.8	0.6
	1981	0.9	0.9	1.6	--	0.8
	1982	0.7	1.0	--	--	0.7
	1983	0.6	1.0	1.6	0.4	0.8
	1984	0.5	1.2	1.9	0.8	0.8
	1985	0.6	0.9	1.0	1.4	0.8
15-19	1980	26.2	31.6	35.3	20.4	28.2
	1981	25.7	31.4	30.3	20.2	27.4
	1982	24.3	31.0	30.3	26.6	26.8
	1983	23.2	28.9	30.3	22.1	25.3
	1984	23.1	29.7	32.7	22.1	25.5
	1985	22.8	28.3	28.4	23.7	24.9
20-24	1980	33.0	36.4	33.8	36.1	34.3
	1981	32.7	36.0	45.9	42.7	34.7
	1982	32.5	38.6	34.5	33.7	34.6
	1983	32.2	38.1	37.7	34.4	34.5
	1984	29.7	36.7	36.5	40.2	33.0
	1985	29.4	36.2	42.2	29.3	32.1
25-29	1980	22.4	17.4	21.1	25.2	20.5
	1981	22.4	18.5	13.9	23.5	20.9
	1982	22.0	17.2	21.8	25.1	20.7
	1983	21.1	18.6	16.4	24.4	20.2
	1984	22.4	17.6	19.2	18.5	20.5
	1985	23.2	19.5	15.7	22.8	21.7
30-34	1980	11.9	8.5	6.0	10.4	10.6
	1981	12.0	7.9	6.6	9.1	10.4
	1982	13.4	8.0	5.9	7.9	11.1
	1983	15.0	8.2	9.0	10.7	12.1
	1984	15.1	9.7	6.7	12.4	12.9
	1985	13.4	10.2	9.8	13.0	12.2
35-39	1980	4.0	3.2	3.0	3.8	3.7
	1981	4.6	3.7	0.8	3.3	4.1
	1982	5.3	2.8	3.4	5.2	4.4
	1983	6.1	3.8	3.3	7.3	5.4
	1984	7.1	3.8	2.9	4.0	5.6
	1985	8.0	3.8	2.9	7.4	6.3
>39	1980	1.4	1.7	--	2.3	1.5
	1981	1.3	1.4	0.8	1.3	1.3
	1982	1.4	1.2	2.5	1.1	1.3
	1983	1.4	1.1	1.6	0.4	1.3
	1984	1.7	0.8	--	1.6	1.4
	1985	1.9	0.7	--	2.3	1.4

TABLE 1.17
NEW MEXICO RESIDENTS
REPORTED LEGAL INDUCED ABORTIONS
1982-1985
Percent Distribution by Level of Education

			WHITE					
EDUCATION LEVEL	YEAR	STATE TOTAL	NON-HISPANIC	HISPANIC	BLACK	INDIAN	OTHER	UNKNOWN
< 8	1982	2.3	1.6	3.7	0.8	0.4	11.5	0.0
	1983	2.7	1.8	4.3	1.6	1.9	9.5	0.0
	1984	2.4	1.4	3.8	1.9	2.0	7.1	-
	1985	1.5	0.5	2.5	1.0	3.3	11.1	-
9-11	1982	13.4	9.9	18.9	12.6	16.9	3.8	-
	1983	12.7	10.7	15.6	11.5	14.9	11.9	-
	1984	14.7	11.8	19.0	14.4	18.5	3.6	-
	1985	12.6	10.4	16.2	10.8	13.0	8.3	-
12	1982	37.7	32.9	45.8	37.0	39.0	38.5	-
	1983	37.7	32.1	45.1	41.0	46.2	38.1	-
	1984	38.3	33.2	45.1	35.6	46.2	32.1	-
	1985	37.5	33.1	44.1	38.2	40.5	27.8	-
> 12	1982	31.3	35.8	22.9	30.3	39.0	11.5	0.0
	1983	32.7	38.8	24.2	23.8	34.7	21.4	-
	1984	31.0	36.9	22.0	32.7	30.5	37.5	-
	1985	27.9	32.1	20.1	30.4	38.1	27.8	-
UNKNOWN/NS	1982	15.4	19.8	8.7	19.3	4.9	34.6	100.0
	1983	14.1	16.6	10.9	22.1	2.3	19.0	100.0
	1984	13.6	16.8	10.1	15.4	2.8	19.6	-
	1985	20.5	24.0	17.1	19.6	5.1	25.0	-

Nationally women obtaining abortions were most likely to be young, white, single, and of low parity.

Over three-quarters of New Mexico resident women obtaining abortions during 1985 were single at the time of abortion; this percentage represents a slight decrease from the previous years. The percentage single was highest for Black women (Table 1.18), while the previous year it had been highest for the state's Hispanic women.

Nationally, 78.6% of women obtaining abortions were single, this percentage has been increasing yearly, up 12% since 1972.

TABLE 1.18
NEW MEXICO RESIDENTS
REPORTED LEGAL INDUCED ABORTIONS
1981-1985
Percentage Distribution by Marital Status

YEAR/ STATUS	ALL RACES	TOTAL WHITE	NON HISPANIC WHITE	HISPANIC WHITE	BLACK	INDIAN	OTHER	UNKNOWN
1981								
MARRIED	22.5	22.2	23.0	20.9	16.4	25.1	64.7	44.4
SINGLE	76.6	76.9	76.0	78.4	83.6	74.6	35.3	51.9
UNK./NS	0.9	0.9	1.0	0.4	-	0.3	-	3.7
1982								
MARRIED	21.8	21.5	22.8	19.2	30.3	21.7	57.7	37.5
SINGLE	76.4	76.8	75.3	79.5	68.1	77.2	38.5	50.0
UNK./NS	1.7	1.7	2.0	1.2	1.7	1.1	3.8	12.5
1983								
MARRIED	21.3	21.0	22.9	18.0	14.8	25.2	50.0	37.5
SINGLE	77.0	77.3	75.2	80.7	80.3	74.4	47.6	50.0
UNK./NS	1.7	1.7	1.9	1.3	4.9	0.4	2.4	12.5
1984								
MARRIED	20.3	20.0	22.4	16.6	20.2	18.9	48.2	-
SINGLE	78.5	78.8	76.3	82.4	77.9	80.7	51.8	-
UNK./NS	1.2	1.2	1.3	1.0	1.9	0.4	0	-
1985								
MARRIED	20.1	20.1	22.4	16.6	11.8	17.7	52.8	-
SINGLE	76.5	76.3	73.6	80.5	87.3	80.0	47.2	-
UNK./NS	3.4	3.6	4.0	3.0	1.0	2.3	0	-

59.5% of New Mexico resident women obtaining an abortion in 1985 were obtaining an abortion for the first time. The percentage of women who reported no previous abortion has decreased yearly since 1981 (Table 1.19, on the following page). The percentage reporting three or more has been steadily increasing - a 67% increase since 1981; because of the number of abortions performed each year, the population at risk of having more than one abortion has increased.

New Mexico resident White non-Hispanic women have shown the most steady increase in the percentage of women having had three or more previous abortions as the percentage has increased nearly 72% since 1981.

TABLE 1.19
NEW MEXICO RESIDENTS
REPORTED LEGAL INDUCED ABORTIONS
1981-1985
Percent Distribution by Previous Induced Abortions

PREVIOUS INDUCED ABORTIONS	STATE TOTAL	TOTAL WHITE	WHITE		BLACK	INDIAN	OTHER	UNKNOWN
			NON-HISPANIC	HISPANIC				
0								
1981	67.7	67.5	65.5	70.7	68.9	71.0	-	63.0
1982	67.0	66.8	63.9	71.8	69.7	71.2	42.3	75.0
1983	64.3	64.1	59.7	70.8	61.5	69.8	54.8	100.0
1984	62.5	62.3	58.2	68.3	60.6	69.9	51.8	-
1985	59.5	59.4	55.5	65.4	64.7	67.0	63.9	-
1								
1981	22.7	22.7	23.6	21.2	21.3	21.8	-	25.9
1982	23.0	23.1	24.3	21.0	25.2	21.3	11.5	-
1983	24.5	24.6	26.9	21.1	26.2	21.8	23.8	-
1984	25.1	25.1	27.0	22.3	32.7	20.5	33.9	-
1985	25.4	25.4	27.0	23.0	25.5	26.0	19.4	-
2								
1981	6.4	6.5	7.2	5.4	9.0	3.9	-	3.7
1982	6.3	6.5	7.4	4.9	2.5	4.5	23.1	-
1983	7.5	7.5	9.0	5.0	11.5	5.3	16.7	-
1984	8.1	8.2	9.5	6.2	4.8	6.8	10.7	-
1985	7.9	8.1	9.3	6.3	6.9	3.7	5.6	-
≥ 3								
1981	2.7	2.7	3.2	2.0	-	2.3	-	7.4
1982	3.0	3.1	3.8	1.9	0.8	2.2	23.1	-
1983	3.4	3.6	4.1	2.8	0.8	1.9	4.8	-
1984	3.9	4.0	5.1	2.4	1.0	2.0	3.6	-
1985	4.5	4.6	5.5	3.3	1.0	2.8	11.1	-
UNKNOWN NS								
1981	0.6	0.5	0.4	0.7	0.8	1.0	-	-
1982	0.6	0.5	0.6	0.4	1.7	0.7	-	25.0
1983	0.3	0.3	0.3	0.2	0.2	1.1	-	-
1984	0.4	0.4	0.2	0.8	1.0	0.8	-	-
1985	2.3	2.4	2.7	2.1	2.0	0.5	-	-

Slightly less than half (49.7%) of all legal abortions reported nationally were performed in the first eight weeks of gestation, 89.3% were performed in the first twelve weeks of gestation. Nationally the percentage obtained in the first eight weeks of gestation peaked in 1980 at 51.7% then has decreased yearly to 49.7% in 1983.

In New Mexico for 1985, 47% of abortions were obtained in the first eight weeks (Table 1.20) and 87.3% in the first twelve weeks. The percentage during the first eight weeks increased 9% over the prior year in 1985 after having decreased for three years.

Over half of abortions obtained by New Mexico resident women age 25 and older were obtained in the first eight weeks, younger women appear more likely to obtain abortions at a later gestation than do older women.

TABLE 1.20
NEW MEXICO RESIDENT ABORTIONS
PERCENTAGE DISTRIBUTION
BY AGE GROUP AND GESTATION PERIOD
1981-1985

GESTATION PERIOD	1981	1982	1983	1984	1985
< 9 Weeks: All Ages	45.8	44.5	44.4	43.2	47.0
< 15	25.6	42.1	25.6	30.0	29.7
15-19	36.6	35.3	33.1	33.3	37.1
20-24	43.1	43.3	42.1	39.1	42.7
25-29	54.2	49.9	51.1	50.4	55.2
30-34	51.1	54.3	57.2	55.0	57.6
35-39	57.2	54.3	56.7	57.4	59.3
> 39	50.0	59.2	53.1	47.8	59.7
Unknown/NS	50.0	62.5	60.0	65.0	50.0
9-10 Weeks: All Ages	28.8	29.2	28.4	30.2	27.5
< 15	32.6	18.4	28.2	17.5	29.7
15-19	30.3	30.3	27.8	30.6	30.6
20-24	30.2	29.0	29.6	32.6	28.4
25-29	25.9	31.2	27.2	29.6	24.9
30-34	25.0	26.6	28.8	28.1	25.4
35-39	24.7	27.7	26.7	25.5	23.4
> 39	25.0	19.7	32.8	24.6	23.9
Unknown/NS	37.5	6.3	30.0	20.0	25.0
11-12 Weeks: All Ages	13.3	13.6	13.6	14.1	12.8
< 15	20.9	13.2	28.2	12.5	10.8
15-19	15.9	17.2	18.7	17.2	15.0
20-24	13.9	14.0	13.8	15.4	15.8
25-29	10.7	10.1	11.5	11.2	10.2
30-34	9.5	11.4	8.1	10.7	8.5
35-39	11.6	10.8	9.3	10.6	8.8
> 39	11.8	9.9	7.8	15.9	3.0
Unknown/NS	11.8	18.8	5.0	10.0	8.3
13-15 Weeks: All Ages	8.4	7.7	7.6	6.3	7.0
< 15	16.3	21.1	5.1	20.0	16.2
15-19	11.3	10.7	10.7	9.1	9.4
20-24	8.0	7.9	8.3	6.9	7.8
25-29	6.3	5.4	6.6	4.4	5.5
30-34	6.5	5.5	3.3	2.9	3.9
35-39	4.7	3.0	3.7	2.8	3.7
> 39	7.4	4.2	1.6	4.3	7.5
Unknown/NS	6.3	6.3	-	5.0	4.2
16-20 Weeks: All Ages	3.7	4.3	5.6	5.3	4.8
< 15	4.7	5.3	12.8	17.5	13.5
15-19	5.1	5.9	9.1	8.6	6.7
20-24	4.0	5.0	5.8	4.9	4.5
25-29	1.9	2.5	3.1	3.6	3.9
30-34	3.2	1.7	2.6	2.6	3.7
35-39	1.9	2.6	3.7	3.2	2.7
> 39	2.9	7.0	4.7	7.2	6.0
Unknown/NS	-	6.3	-	-	8.3
> 21 Weeks: All Ages	0.1	-	0.3	0.6	0.4
< 15	-	-	-	2.5	-
15-19	0.1	-	0.4	0.8	0.4
20-24	0.1	-	0.3	0.8	0.5
25-29	-	-	0.2	0.5	0.1
30-34	-	-	-	0.5	0.7
35-39	-	-	-	-	-
> 39	1.5	-	-	-	-
Unknown/NS	-	-	5.0	-	-
Unknown Weeks: All Ages	0.8	0.7	0.1	0.3	0.5
< 15	-	-	-	-	-
15-19	0.8	0.6	0.2	0.5	0.8
20-24	0.8	0.8	0.1	0.2	0.3
25-29	1.0	0.9	0.3	0.4	0.2
30-34	0.7	0.5	-	0.2	0.2
35-39	-	1.3	-	0.4	2.0
> 39	1.5	-	-	-	-
Unknown/NS	6.3	-	-	-	4.2

TABLE 1.21
NEW MEXICO RESIDENTS 1980-1985 REPORTED LEGAL INDUCED ABORTIONS PERCENTAGE DISTRIBUTION BY WEEKS GESTATION/RACE/ETHNIC

ETHNIC/RACE	TOTAL	PERCENTAGE DISTRIBUTION BY WEEKS GESTATION					
		< 9	9-10	11-12	13-15	16-20+	UNKNOWN
1985							
STATE TOTAL	4651	47.0%	27.5%	12.8%	7.0%	5.2%	0.5%
WHITE	4298	47.8	27.5	12.5	6.9	4.8	0.5
Non-Hispanic	2608	52.3	26.5	10.9	5.9	3.8	0.6
Hispanic	1690	40.8	29.1	15.0	8.4	6.3	0.4
BLACK	102	37.3	31.4	16.7	7.8	6.9	--
INDIAN	215	34.4	25.1	17.2	9.3	13.0	0.9
OTHER	36	63.9	19.4	8.3	5.6	2.8	--
UNKNOWN	--	--	--	--	--	--	--
1984							
STATE TOTAL	5034	43.2	30.2	14.1	6.3	5.9	0.3
WHITE	4625	43.8	30.3	13.9	5.9	5.8	0.3
Non-Hispanic	2765	48.5	29.7	12.8	4.6	4.2	0.2
Hispanic	1860	36.8	31.2	15.7	7.7	8.1	0.5
BLACK	104	36.5	31.7	10.6	11.5	9.6	--
INDIAN	249	29.7	31.3	17.3	12.9	8.0	0.8
OTHER	56	60.7	17.9	16.1	1.8	3.6	--
UNKNOWN	--	--	--	--	--	--	--
1983							
STATE TOTAL	5010	44.5	28.4	13.6	7.6	5.9	0.1
WHITE	4576	45.3	28.9	13.1	7.1	5.4	0.2
Non-Hispanic	2774	50.2	27.4	11.4	6.0	4.8	0.2
Hispanic	1802	37.7	31.1	15.8	8.9	6.4	0.1
BLACK	122	36.9	20.5	22.1	12.3	8.2	--
INDIAN	262	32.1	24.8	19.1	11.8	12.2	--
OTHER	42	45.2	23.8	11.9	16.7	2.4	--
UNKNOWN	8	37.5	25.0	12.5	--	25.0	--
1982							
STATE TOTAL	5213	44.5	29.2	13.6	7.7	4.3	0.7
WHITE	4797	46.0	29.0	13.2	7.3	3.8	0.7
Non-Hispanic	3027	51.4	28.0	11.2	5.8	2.8	0.8
Hispanic	1770	36.6	30.7	16.7	9.9	5.6	0.6
BLACK	119	30.3	33.6	15.1	10.1	10.1	0.8
INDIAN	267	24.3	31.5	19.5	14.6	9.7	0.4
OTHER	26	42.3	30.8	15.4	3.8	7.7	--
UNKNOWN	4	75.0	--	--	--	--	25.0
1981							
STATE TOTAL	5194	45.5	28.5	13.2	8.3	3.7	0.8
WHITE	4742	46.5	28.7	12.7	7.7	3.5	0.9
Non-Hispanic	2905	50.7	28.5	10.8	6.5	2.6	0.9
Hispanic	1816	39.9	29.0	15.8	9.6	4.9	0.8
BLACK	122	37.7	26.2	14.8	16.4	4.1	0.8
INDIAN	207	29.0	28.7	20.2	15.0	7.2	--
OTHER	17	70.6	17.6	5.9	--	5.9	--
UNKNOWN	27	63.0	18.5	14.8	3.7	--	--
1980							
STATE TOTAL	5373	45.6	29.3	14.6	8.0	2.2	0.4
WHITE	4718	46.6	29.6	14.1	7.4	2.0	0.3
Non-Hispanic	2917	51.3	27.5	13.2	5.9	1.7	0.3
Hispanic	1801	38.9	33.1	15.7	9.7	2.3	0.3
BLACK	133	41.4	27.8	14.3	11.3	4.5	0.8
INDIAN	393	30.5	28.5	20.4	15.8	4.6	0.3
OTHER	61	54.1	24.6	14.8	4.9	1.6	--
UNKNOWN	68	64.7	17.6	11.8	4.4	--	1.5

TABLE 1.22
NEW MEXICO RESIDENTS REPORTED LEGAL INDUCED ABORTIONS 1978-1985
Percentage by Facility

YEAR	PHYSICIAN'S OFFICE	CLINIC	HOSPITAL
1985	62.4%	36.5%	1.1%
1984	65.3	33.2	1.4
1983	67.9	30.6	1.4
1982	66.6	30.3	2.8
1981	64.4	31.5	3.9
1980	59.8	33.6	6.5
1979	50.5	36.6	10.9
1978	39.8	43.8	16.4

Over half of White non-Hispanic women who obtain abortions do so in the first eight weeks of gestation while only about a third of Indian women obtaining abortions do so before the ninth week (Table 1.21).

62.4% of New Mexico women obtaining abortions in 1985 did so in a physicians office; this is a decrease from the previous year, but is 57% higher than the 1978 level. Hospital facility use has decreased over 90% (Table 1.22).

Over the four year period 1982-1985, Santa Fe County residents had the highest abortion ratio with one abortion for every two live births. Bernalillo County was second highest (Figure 1.17).

Very low abortion ratios for some counties may be indicative of a reporting problem, also residents who go to facilities in other states, for example, Texas, would not be counted in these data.

Of reported abortions performed in the State, 69% are performed in Bernalillo County and another 23% in Santa Fe County (Figure 1.17).

Curettage has been increasingly the most common method of abortion nationally, reported for almost 97% of abortions in 1983. New Mexico 1985 data shows 99% reporting curettage as the procedure of termination.

Nationally in 1981, seven deaths were associated with legal abortion. In New Mexico, thru 1985, no deaths have been associated with induced abortion.

FIGURE 1.17
NEW MEXICO RESIDENTS ABORTION RATIOS 1982-1985
(Per 1,000 Live Births)
PERCENTAGE BY COUNTY OF OCCURRENCE
1985


SAN JUAN 37.1
RIO ARRIBA 252.1
TAOS 286.3
COLFAX 181.2
UNION 91.2
LOS ALAMOS 282.5
MORA 91.2
HARDING 109.1
McKINLEY 47.9
SANDOVAL 154.6
SANTA FE 517.5
SAN MIGUEL 215.5
BERNALILLO 346.9
CIBOLA 106.0
QUAY 148.2
GUADALUPE 83.1
VALENCIA 153.5
TORRANCE 117.6
CURRY 31.0
DE BACA 25.6
CATRON 191.5
SOCORRO 185.7
ROOSEVELT 69.6
LINCOLN 140.1
CHAVES 165.0
SIERRA 43.2
GRANT 6.5
OTERO 15.6
LEA 63.2
LUNA 8.2
DONA ANA 13.3
EDDY 98.6
HIDALGO 0
68.7%
22.8%
2.0%-6.4%
Less Than 1%
None

NATALITY

FERTILITY

Over the years New Mexico's birth rate trend has quite closely paralleled that for the United States, though the State's rate has consistently been higher than the national rate (Figure 2.1).

FIGURE 2.1
LIVE BIRTHS PER 1,000 POPULATION
NEW MEXICO RESIDENTS AND UNITED STATES
1950-1985



Beginning in 1970, the State's birth rate decreased to reach its lowest level in 1975, then starting with the next year, the birth rate increased annually until reaching 20.4 in 1980, the highest rate since 1971. The rates for the next three years fluctuated slightly as 1981 followed with a slight decrease to 19.9, followed then by a 1½% increase to 20.2 in 1982, and then dropped three consecutive years: in 1983 by 2½% to 19.7, in 1984 by 2.6% to 19.2, and in 1985 by 1.6% to 18.9.

Nationally in the years following 1970 the birth rate fell to all time lows in 1975 and 1976 with subsequent years showing a general increase reaching 15.9 in 1980 and again in 1982, its highest level since 1971 and 9% higher than the low rates of 1975 and 1976. The national birth rate fell by 3.1% to 15.5 in 1983 and 1984, then increased by 1.3% to 15.7 in 1985. The 1985 New Mexico birth rate exceeded the national rate by 20%.

New Mexico's birth rate for 1985 represents 27,449 births. Births occurring in 1985 and the previous three years represent highest number of resident annual births since 1963. The State's 1985 birth rate of 18.9 ranked fourth highest among the States. Alaska ranked highest with 24.1, Utah second highest with 23.4, and Texas third highest with 19.2. Connecticut reported the lowest rate (11.6).

The New England and Middle Atlantic divisions recorded the lowest birth rates nationally in 1985 (Figure 2.2). The birth rate decreased from the prior year only for the East North Central Division. No change was seen in the New England and West North Central Divisions. Increases were seen in the Middle Atlantic, East South Central, South Atlantic, West South Central, Mountain, and Pacific Divisions. The birth rate decreased in 25 states, but remained steady in 2 and increased in 23.

Of the fourteen states recording the highest 1985 birth rates, six are those in the Mountain Division and the remainder in the Pacific (including Alaska and Hawaii), West South Central and West North Central divisions. It is not surprising then that the Mountain Division registered the highest rate of any division in the nation (Figure 2.2).


FIGURE 2.2
UNITED STATES
LIVE BIRTH RATES
BY DIVISION
1985

RATE PER 1,000 POPULATION

Of New Mexico's 33 counties, 13 showed increases in birth rate during 1985, 2 showed no change, while the remainder experienced decreases. (Table 2.1). The birth rate in 26 of the counties exceeded the national rate by up to 73%.

In 1985 the birth rates ranged from a high of 27.2 births per thousand population in McKinley County to a low of 10.3 in Sierra County. Bernalillo County, which accounted for 29% of the State's total births, exceeded the national birth rate by 9%. The other six counties registering more than a thousand births during the year experienced birth rates exceeding the national rates by 9 to 73 percent. Rates by county for several years are listed in Table 2.1 on the following page.

Two of the three counties with the highest rates in the State in 1985 experienced decreases in rates from the prior year (Rio Arriba and San Juan), while one showed no change (McKinley).

An examination of the State's birth rate by ethnicity and race shows that the decrease in the New Mexico birth rate in 1985 may be attributed to a decreased rate for Whites (Figure 2.3).

The rate for the non-Hispanic White population in the State dropped by 3% in 1985 to reach an eight year low. The birth rate for non-Hispanic Whites has been the lowest of the State's various racial and ethnic groups, being in 1985, 29% lower than the overall State rate and 15% lower than the national rate for all races.

In 1985 the birth rate for Hispanic Whites was 77% higher than for non-Hispanic Whites in the State and 51% higher than the total U.S. rate. The rate for Hispanic Whites fell by 1% in 1985. The 1984 rate for New Mexico Hispanic Whites was 65% higher than the birth rate for Whites nationally, while the rate for New Mexico non-Hispanic Whites was 5% lower.

TABLE 2.1
NEW MEXICO BIRTH RATE BY COUNTY OF RESIDENCE
1974-1985
(Rates Per 1,000 Population)

	COUNTY	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985
	UNITED STATES	14.8	14.6	14.6	15.1	15.0	15.6	15.9	15.8	15.9	15.5	15.5	15.7
	NEW MEXICO	18.9	18.1	18.5	18.8	19.1	19.3	20.4	19.9	20.2	19.7	19.2	18.9
	CIBOLA	*	*	*	*	*	*	*	*	22.3	20.9	20.8	20.8
	1 MCKINLEY	29.4	29.6	29.0	29.5	29.2	28.2	29.4	28.5	28.7	27.5	27.2	27.2
	SAN JUAN	22.4	23.4	24.1	24.9	25.0	26.3	28.4	27.8	27.6	25.0	23.4	24.5
	COLFAX	18.2	14.7	15.0	16.5	15.6	16.5	17.5	15.9	18.2	18.5	15.9	17.9
	HARDING	10.0	6.7	14.2	8.3	14.5	13.6	11.9	17.0	12.0	16.0	11.2	16.0
	LOS ALAMOS	11.0	10.6	11.6	10.5	12.4	10.9	11.4	11.5	11.1	11.1	10.5	11.1
	MORA	13.8	14.7	12.1	13.9	14.2	17.7	21.4	18.4	18.9	17.1	15.9	15.7
	2 RIO ARRIBA	23.7	21.4	21.0	21.4	22.4	21.3	22.5	21.6	21.0	21.5	22.8	23.4
	SAN MIGUEL	18.9	16.9	19.6	18.7	19.9	20.4	22.1	20.3	19.2	20.0	20.2	18.8
H	SANTA FE	16.7	15.1	16.4	16.3	16.7	16.4	16.9	17.1	15.6	17.0	17.4	17.1
E	TAOS	19.2	18.9	17.9	19.1	18.8	20.7	21.4	20.8	22.9	21.7	21.3	20.9
A	UNION	15.0	10.2	14.7	15.4	14.5	15.8	15.9	19.0	17.1	15.1	15.6	14.9
L													
T	BERNALILLO	16.4	15.4	15.9	16.5	16.9	17.5	18.3	17.2	17.7	17.2	17.3	17.1
H	SANDOVAL	22.6	19.0	18.7	18.3	17.1	16.0	19.7	20.5	20.2	22.4	25.4	22.2
	3 TORRANCE	15.9	17.4	14.5	15.8	15.4	15.1	17.4	16.7	15.9	18.0	16.5	15.8
D	VALENCIA	19.4	19.3	22.2	23.7	24.4	23.7	23.7	21.0	18.7	17.4	16.9	17.5
I													
S	CHAVES	17.4	14.5	15.7	18.2	17.4	16.7	19.7	19.3	21.3	20.9	18.1	17.9
T	CURRY	23.3	22.1	23.4	23.1	23.3	22.6	23.0	21.2	21.6	20.5	22.3	21.0
R	DE BACA	9.6	12.8	10.8	12.9	13.2	17.5	16.7	15.2	16.7	17.5	16.4	15.0
I	EDDY	16.9	18.3	18.4	18.3	19.1	18.2	20.5	19.5	22.6	19.4	17.3	16.9
C	4 GUADALUPE	16.5	14.4	16.3	15.6	19.6	15.7	21.1	18.9	20.2	20.4	17.5	17.7
T	LEA	20.8	21.3	19.9	17.4	20.5	20.6	23.4	25.5	25.8	24.7	20.8	20.9
S	LINCOLN	18.4	15.5	16.3	18.4	14.8	15.9	15.4	19.3	20.3	20.0	18.2	17.7
	QUAY	16.9	15.5	14.2	14.4	16.5	17.0	8.7	15.1	19.0	15.4	17.2	15.0
	ROOSEVELT	16.5	16.7	16.1	15.4	17.4	17.4	18.5	18.7	20.4	17.7	16.0	17.3
	CATRON	10.0	11.3	18.8	21.2	14.4	16.7	18.0	15.2	14.8	13.2	10.8	12.1
	DONA ANA	19.7	19.1	18.6	18.1	17.1	18.1	19.1	19.6	19.3	19.6	20.4	18.1
	GRANT	20.6	20.0	19.6	20.5	20.2	21.7	20.9	22.0	19.3	16.6	17.0	18.9
	5 HIDALGO	24.2	23.7	23.3	22.5	23.5	21.3	23.0	18.1	22.5	22.5	21.3	17.3
	LUNA	19.7	20.6	18.5	19.2	15.0	19.2	19.0	18.7	19.9	17.9	17.5	17.3
	OTERO	22.7	21.5	23.0	21.8	22.0	20.8	20.8	21.7	23.0	24.0	22.0	22.6
	SIERRA	10.9	11.2	11.6	12.4	8.5	11.8	10.8	18.8	10.8	14.4	10.6	10.3
	SOCORRO	19.0	20.8	18.8	20.9	20.3	18.7	19.7	20.1	16.6	19.2	16.5	18.2

* Prior to 1982, Cibola County data were included in Valencia County.

** Rates are based on revised population estimates.

The birth rate for the New Mexico Black population, which increased by 4% in 1985, was the second highest birth rate of the State's various race/ethnic groups and exceeded the total State rate by 33%. In 1984 the birth rate for New Mexico Blacks exceeded by 15% the rate for Blacks nationally.

The reported birth rate for the Indian has over the years exceeded that of all other race/ethnic populations in the State. The rate in 1985 increased by 2% over the 1984 rate, and was 71% higher than the State rate. In 1984 the rate for New Mexico Indians was double the U.S. rate and 49% higher than the birth rate for "other than white" population nationally.

The birth rate for non-Hispanic White increased from 1975 until 1979 and, except for a slight increase in 1982, declined each of the following years, dropping in 1985 to 4% less than the 1975 level. The birth rate for Hispanic White showed a general upward trend since 1975 until 1980, with the rate leveling off to remain, with slight fluctuations, at just below the 1980 rate during subsequent years through 1985. The 1985 rate was 13% higher than that for 1975.

For the State's Black population, fluctuating rates since 1975 showed a slight general decline, reaching a low point in 1979, before surging upward to a high point in 1982, then dropping for two years before increasing slightly in 1985. The 1985 rate was 12% lower than the peak 1982 rate.

FIGURE 2.3
BIRTH RATE BY RACE/ETHNIC NEW MEXICO RESIDENTS
1980-1985



Birth rates for New Mexico's Indian population fell sharply after 1976 until 1979; since then the rates have fluctuated each year within a range of 31 to 32. The 1985 rate was 9% lower than the 1976 rate.

Birth rates by New Mexico race/ethnic groups for years 1980 through 1985 are presented in Figure 2.3.

The measure of the rate of childbearing is the fertility rate. The fertility rate is the number of births per thousand women in the childbearing ages of 15-44. The purpose of using the fertility rate in comparing the fertility of two populations is to eliminate the differences created in the number of births because of differences in the age and sex compositions of the population: this results in a measure of childbearing that uses as denominators standardized populations of those "at risk" (women of childbearing age). In other words, the fertility rate enables us to distinguish whether, in a given population, women are having more babies or whether there were more babies because there were more women of childbearing age.

In 1960, the fertility rate in New Mexico was 156.2 births per every thousand women ages 15-44. By 1970, New Mexico women were having fewer babies—100.8 births per thousand women ages 15-44. The rate of childbearing in the State had by 1980 dropped further to 85.4 births per thousand women ages 15-44. Since then, the fertility rate has dropped each year, with the exception of a slight increase in 1982, reaching an all time low of 80.0 in 1985. The 1985 rate of childbearing in the State was 6% lower than in 1980, 21% lower than in 1970, and 49% lower than in 1960.

The rate of childbearing for New Mexico women has over the years exceeded the national rate and was 23% higher in 1984 (Table 2.2).

NATALITY BY AGE OF MOTHER

The age-specific birth rates in Table 2.2 show higher rates of childbearing in New Mexico than nationally for all but the 10-14 age group of mothers, comparing data for 1984.

The highest rate of childbearing for New Mexico and second highest for the nation was in the 20-24 age group. Nationally, the rate was highest for the 25-29 age group, slightly more than the rate for the 20-24 age group.

In the 20-24 age group the State's rate is 27% higher than the national rate and 15% higher than nationally for the 25-29 age group.

The rate of childbearing for New Mexico women 30 years of age and older is higher than for women nationally. A general increasing trend was noted from 1979 to 1984 for women 30 years and older in the State, reflecting national trends, then State rates dropped off in 1985.

The nation's as well as New Mexico's teenage women (15-19) had experienced a lower childbearing rate than in earlier years; the State's rate being 16% higher than nationally in 1984. Increases in the rates of childbearing for the 15-19 and 20-24 age groups were seen Statewide in 1985.

Of the 27,449 New Mexico resident births during 1985, the largest percentage was to mothers in the 20-24 age group, 33.7%. This percentage was a 2% increase from the previous year, and a 6% decrease since 1980 (Table 2.3). Women in their twenties accounted for 63% of all births in New Mexico during 1985.

The percentage of live births decreased in 1985 as compared with 1984 for New Mexico women ages 24 and younger and increased for women ages 25-39.

In 1984, the percentage of births to New Mexico women was higher than national percentages for ages 15-24 and lower for ages younger than 15 and 25-39: the percentage was the same for the 40 and older age group. The trend toward an increasing percentage of births in the 30-34 and 35-39 age groups is evident both in New Mexico and nationally (Table 2.4).

TABLE 2.2
NEW MEXICO RESIDENTS 1960-1985 BY RACE/ETHNIC UNITED STATES 1960-1984 FERTILITY RATES AND BIRTH RATES BY AGE OF MOTHER

U.S. & N.M. By Race/Ethnic	FERTILITY RATE a/	Birth Rate By Age Group								
		10-14	(15-17)	(18-19)	15-19	20-24	25-29	30-34	35-39	40-45
UNITED STATES:										
1985	NOT AVAILABLE AT THIS TIME									
1984	65.4	1.2	(31.1)	(78.3)	50.9	107.3	108.3	66.5	22.8	2.3
1983	65.8	1.1	(32.0)	(78.1)	51.7	108.3	108.7	64.6	22.1	2.2
1982	67.3	1.1	(32.4)	(80.7)	52.9	111.3	111.0	64.2	21.1	2.1
1981	67.4	1.1	(32.1)	(81.7)	52.7	111.8	112.0	61.4	20.0	2.0
1980	68.4	1.1	(32.5)	(82.1)	53.0	115.1	112.9	61.9	19.8	1.9
1979	67.2	1.2	(32.3)	(81.3)	52.3	112.8	111.4	60.3	19.5	1.9
1970	87.9	1.2	(38.8)	(114.7)	68.3	167.8	145.1	73.3	31.7	4.3
1960	118.0	.8	(31.5)	(188.5)	89.1	258.1	197.4	112.7	56.2	8.5
NEW MEXICO All Races										
1985	80.0	1.1	(42.9)	(116.7)	(72.6)	140.0	124.0	71.5	28.2	3.2
1984	80.3	1.1	(34.0)	(96.6)	59.2	136.6	124.1	76.6	30.8	3.2
1983	82.2	1.4	(39.7)	(107.1)	66.7	139.7	123.3	75.6	31.0	3.3
1982	84.6	1.0	(40.9)	(111.4)	69.2	145.0	128.3	76.4	29.0	3.3
1981	83.4	1.4	(40.3)	(110.9)	68.6	144.7	147.5	74.3	25.3	3.2
1980	85.4	1.2	(43.9)	(115.9)	72.8	151.8	126.9	72.5	25.0	3.5
1979	83.7	1.1	(43.0)	(115.4)	72.1	148.1	128.1	66.0	25.4	3.4
1970	100.8	.6	(37.2)	(145.8)	76.7	201.9	157.7	82.3	41.1	7.0
1960	156.2	.8	(64.5)	(245.1)	127.3	325.4	222.5	133.2	88.4	13.9
NON-HISPANIC-WHITE										
1985	57.7	.4	NOT AVAILABLE		40.8	99.9	98.2	60.2	19.8	2.3
1984	59.3	.4	"	"	34.3	101.5	96.2	65.7	25.6	1.5
1983	61.5	.6	"	"	40.7	106.9	99.0	63.9	21.3	1.7
1982	66.2	.5	"	"	47.4	118.9	106.4	63.6	20.1	1.8
1981	64.9	.6	"	"	47.2	115.3	107.8	60.5	17.9	1.7
1980	66.8	.5	"	"	49.8	122.7	109.6	59.2	15.3	1.7
1979	68.5	.4	"	"	51.7	124.2	117.3	55.7	18.3	1.8
HISPANIC WHITE										
1985	98.4	1.5	"	"	95.8	173.0	147.3	75.3	32.7	3.7
1984	98.0	1.5	"	"	77.6	163.7	154.4	81.5	34.0	4.4
1983	99.4	1.8	"	"	87.3	165.6	145.3	81.6	36.4	4.6
1982	98.4	1.4	"	"	84.7	163.2	143.6	85.3	35.4	4.6
1981	98.9	1.6	"	"	83.9	168.9	145.8	82.6	30.9	4.7
1980	101.7	1.3	"	"	89.2	176.4	142.9	84.9	30.6	5.0
1979	97.8	1.7	"	"	88.3	170.1	138.7	73.2	33.0	4.7
BLACK										
1985	112.2	5.2	"	"	111.3	181.1	148.8	96.7	37.8	--
1984	105.4	4.0	"	"	82.9	175.1	145.8	95.3	20.8	2.7
1983	119.3	1.6	"	"	95.8	175.7	194.0	96.9	35.9	1.9
1982	126.7	3.4	"	"	106.9	205.5	184.8	95.0	29.0	2.9
1981	113.3	.8	"	"	111.8	179.4	142.7	93.4	25.7	2.9
1980	102.3	5.3	"	"	103.6	165.5	131.1	64.6	35.0	1.0
1979	100.7	.9	"	"	101.7	177.8	119.1	58.2	18.4	5.2
INDIAN										
1985	131.7	1.9	"	"	108.0	202.9	189.5	132.1	[illegible]	9.1
1984	125.3	1.5	"	"	86.1	202.2	173.6	128.8	13.1	10.6
1983	125.9	2.7	"	"	90.1	193.8	179.0	130.3	77.4	9.4
1982	127.9	1.4	"	"	94.4	189.3	198.1	127.1	68.6	9.6
1981	122.8	1.5	"	"	90.8	192.4	174.0	132.1	57.0	9.4
1980	126.4	2.7	"	"	101.2	203.1	177.6	115.4	55.2	11.4
1979	116.9	1.0	"	"	91.0	184.7	168.1	113.2	49.2	11.0

a/ Births per 1,000 women ages 15-44

Rates per 1,000 women in age group. Also referred to as Age Specific Fertility Rates.

New Mexico rates based on 1980 Census Figures, and 1980 race and age composition as applied to 1979 and 1981-1984 total State population estimates furnished by the Bureau of the Census. 1985 based on adjusted 1985 BBER estimates by age and sex; by race based on estimates updated from 1980 census.

TABLE 2.3
NEW MEXICO RESIDENT LIVE BIRTHS
Percentage Distribution By Age of Mother
1960-1985

Age of Mother	Percentage Distribution of Birth by Age Group								
	1960	1970	1976	1980	1981	1982	1983	1984	1985
< 15	.1	.2	.4	.2	.3	.2	.3	.2	.2
15-17	5.1	5.6	8.0	6.5	6.1	6.1	6.1	5.3	5.3
18-19	11.1	12.8	12.4	11.4	11.2	11.1	11.0	10.1	10.0
20-24	35.2	38.6	36.1	35.9	35.0	34.6	34.3	34.4	33.7
25-29	23.9	24.0	27.0	27.6	28.3	28.1	27.8	28.7	28.8
30-34	14.3	12.6	11.3	13.8	14.5	14.7	14.9	15.5	15.7
35-39	8.0	5.6	3.6	3.7	3.9	4.4	4.8	4.9	5.5
40-44	2.1	1.7	1.1	.8	.8	.8	.8	.8	.8
> 45	.2	.1	.1	.1	.1	.1	.0	.0	.0
Total	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%	100.0%
(Actual Births)	30,747	22,004	22,180	26,589	26,565	27,630	27,508	27,350	27,449

TABLE 2.4
LIVE BIRTHS
Percentage Distribution By Age of Mother
NEW MEXICO RESIDENTS 1976-1985
UNITED STATES 1976-1984

	PERCENTAGE DISTRIBUTION BY AGE GROUPS								
	< 15	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40+
NEW MEXICO									
1985	.2	15.3	5.4	9.9	33.6	28.8	15.7	5.5	.8
1984	.2	15.5	5.3	10.2	34.4	28.7	15.5	4.9	.8
1983	.3	17.0	6.1	11.0	34.3	27.8	14.7	4.8	.8
1982	.2	17.2	6.1	11.1	34.6	28.1	14.7	4.4	.8
1981	.3	17.3	6.1	11.2	35.0	28.3	14.5	3.9	.8
1980	.2	17.9	6.5	11.4	35.9	27.6	13.8	3.7	.9
1979	.2	18.1	6.5	11.6	35.7	28.4	12.8	3.9	.9
1978	.2	19.0	6.9	12.0	35.7	27.7	12.7	3.8	1.0
1977	.2	18.9	7.3	11.6	35.7	28.1	12.4	3.8	.9
1976	.4	20.4	8.0	12.4	36.1	27.0	11.3	3.6	1.1
UNITED STATES									
1984	.3	12.8	4.5	8.3	31.1	31.8	17.9	5.3	.8
1983	.3	13.4	4.7	8.7	31.9	31.5	17.2	5.0	.7
1982	.3	14.0	4.9	9.0	32.8	31.3	16.4	4.6	.7
1981	.3	14.5	5.2	9.4	33.4	31.1	16.0	4.0	.7
1980	.3	15.3	5.5	9.8	33.9	30.7	15.2	3.9	.7
1979	.3	15.7	5.7	10.0	34.0	30.6	14.8	3.9	.7
1978	.3	16.3	6.1	10.2	34.2	30.5	14.2	3.8	.7
1977	.3	16.8	6.4	10.4	34.5	30.5	13.4	3.6	.8
1976	.4	17.6	6.8	10.8	34.5	30.7	12.4	3.7	.8

The percentage of births to women under 15 years of age remained unchanged from 1983 to 1984 nationally, and in the State decreased in 1984, but remained steady in 1985. Decreases were noted in the percentage of births to ages 15-19 nationally in 1984 and in the State in 1984 and 1985.

Nationally, 13.1% of all 1984 births were to teenage women; in New Mexico, this percentage was 15.7%, about 20% higher than nationally, and in 1985 had dropped 1% to 15.5%. The 1984 percentage of births to the State's non-Hispanic White teenagers was 20% lower than nationally. However, the percent of teenage births to Hispanic Whites was 48% higher than nationally. In the State, among Blacks, the national figure (for all races) was exceeded by 49%, among Indians by 38% (Figure 2.4).

The trend in births to teenagers has generally been a decreasing one over the last eight years.

Just under half (44.2%) of 1984 births in the United States were to women younger than 25. In New Mexico about half of 1984 births (50.1%) and 1985 births (49.1%) were to women in that age group.

The percentages vary markedly for the State's various ethnic/racial populations. In 1985, 58% of all births to New Mexico Black women were to women under 25 years of age. The percentage was 55% for Hispanic White women, 51% for Indian women and 41% for non-Hispanic White women.

TABLE 2.5
1985
NEW MEXICO RESIDENT LIVE BIRTHS
Percentage Distribution by Age of Mother and Race/Ethnic

RACE/ETHNIC	TOTAL BIRTHS	PERCENTAGE DISTRIBUTION BY AGE GROUPS <15	15-19	20-24	25-29	30-34	35-39	>40
ALL RACES	27,449	0.2%	15.3%	33.7%	28.8%	15.7%	5.5%	0.9%
WHITE:	22,620	0.2	15.0	33.7	29.4	15.7	5.2	0.9
Non-Hispanic	(10,250)	(0.1)	(10.1)	(31.1)	(32.5)	(19.4)	(5.9)	(1.0)
Hispanic	(12,370)	(0.3)	(19.0)	(35.8)	(26.8)	(12.7)	(4.7)	(0.8)
BLACK	675	0.9	19.3	37.8	24.9	12.9	4.3	0.0
INDIAN	3,805	0.3	17.4	33.2	25.4	15.3	7.2	1.3
ALL OTHER NON-WHITE	349	0.0	5.7	28.4	37.2	22.1	6.3	0.3

FIGURE 2.4
PERCENTAGE OF BIRTHS TO TEEN-AGED MOTHERS
NEW MEXICO RESIDENTS
1980-1985



Examining 1985 data for New Mexico's four major race/ethnic groups shows the percentage of births to mothers 25-34 years of age was highest for New Mexico non-Hispanic White women and lowest for Black women with percentages for Hispanic White and Indian women slightly higher than that for Blacks. For ages 35 and older, Indian women showed the highest percentage of births. 1985 data are presented in Table 2.5.

The higher than national rate of childbearing among the State's various ethnic/racial groups is evident in Table 2.2 on the previous page. The 1984 fertility rates for Indians were nearly double the national rates, for Blacks 61%, for Hispanics 50% higher and for non-Hispanic Whites 9% lower than nationally. The fertility rate decreased in the State in 1985 for Whites but increased for Blacks and Indians.

Generally speaking, decreases were seen in the 1985 childbearing rates in the older (30+) age groups of women among New Mexico's race/ethnic groups while increases were seen in the younger (<30) age groups, with some exceptions at various ages for the different racial and ethnic categories. Specifically, decreases were seen for ages 20-24 among non-Hispanic Whites and 25-29 among Hispanic Whites. Increases were seen for ages 40-49 among non-Hispanic Whites, 30-39 among Blacks, and 30-34 among Indians.

A percentage distribution of births by age of mother by county is presented in Table A-3 of the Appendix.

Fertility rates and age-specific birth rates vary markedly by county, year, and age. These data are presented in Table A-4 of the Appendix.

Table 2.6 compares births by age of mother and father for New Mexico (1985) and the United States (1984). Both nationally and for the State, the highest percentage of births occurred to fathers 25-29 years of age. The percentage of births to New Mexico fathers was higher than nationally for 15-24 age groups and lower for the 25-49 age groups.

The percentage of fathers with unreported or unknown ages was 58% higher for New Mexico than the United States.

Comparisons of births by age of father to age of mother show that nearly 80% of all births occurred to mothers younger than thirty, but about half of all births were to fathers of that age group, both nationally and in New Mexico. In contrast, about 4% of all births were to fathers ages forty and older, while less than 1% of births were to mothers of the same age. An important consideration in comparing the percent of births to mothers and fathers ages is the high percent of unknown and unreported ages of fathers (20.4% for New Mexico and 12.9% nationally) versus a negligible percentage of unknown or unreported ages of mothers. Throughout the nation, age of father and other paternal information is often not reported on birth records of babies born to single mothers. In New Mexico, birth certificates of infants born to single mothers, without statements of paternity, contain by law no father information.

TABLE 2.6
PERCENT OF LIVE BIRTHS BY AGE OF MOTHER AND AGE OF FATHER NEW MEXICO RESIDENTS 1985 UNITED STATES 1984

Age of Parent	New Mexico		United States	
	Mother	Father	Mother	Father
< 15	.2	.0	.3	.0
15-19	15.3	3.1	12.8	3.0
20-24	33.6	20.6	31.1	19.0
25-29	28.8	25.4	31.8	29.1
30-34	15.7	18.0	17.9	22.0
35-39	5.5	8.6	5.3	9.7
40-44	.8	2.6	.7	3.0
45-49	.0	.8	.0	.9
50 +	--	.4	N/A	.5
Unknown	N/A	20.4	N/A	12.9

LIVE BIRTH ORDER

The percentage of first live born to New Mexico women has increased substantially since the Sixties. The trend is graphically presented in Figure 2.5 and the actual percentages are presented in Table 2.7.

Between 1960 and 1980 the percentage of first live born to New Mexico women increased by 70%, from 25% of the total to 43%. The percentage of second order live births increased by 32%, third order live births decreased by 15%, fourth order births decreased by 49%, and fifth and higher order births decreased by 75%. From 1980 to 1985 the percentages of first live born dropped six percent to 40.4% of all live births. The percentages of births of second, third, and fourth order increased by 7%, fifth and higher order birth declined by nearly 20%. New Mexico has followed the national trend towards a smaller family over the past four decades, with a shift to slightly larger families during the first half of the eighties.

For women 20 to 39 years of age, the lower percentages of fourth and higher order births indicate a trend of postponement of birth. From 1960 to 1985, for women 20-24, the percentage of third order births decreased 36%, fourth order decreased 65% and fifth and higher order decreased 81%, with accompanying increases in first (49%) and second (11%) order births: the shift is toward first order births; women 25-29 account for the State's second highest portion of births, and in this group is noted the decreased percentages of third and higher order births and increases in the percentages of first (189%) and second (79%) order births. For ages 30-39 the percentages of the first and second order births have tripled since 1960 (Table 2.7).

FIGURE 2.5
NEW MEXICO RESIDENT LIVE BIRTHS PERCENTAGE DISTRIBUTION BY LIVE BIRTH ORDER 1960-1985



TABLE 2.7
NEW MEXICO RESIDENT LIVE BIRTHS PERCENTAGE DISTRIBUTION BY LIVE BIRTH ORDER BY AGE OF MOTHER
1960, 1970, 1980-1985

YEAR	NUMBER OF BIRTHS	DISTRIBUTION BY LIVE BIRTH ORDER 1	2	3	4	5
1985						
TOTAL	27,449	40.4%	32.1%	16.2%	6.7%	4.6%
< 15	63	100.0	--	--	--	--
15-19	4,194	77.6	19.1	3.1	0.2	0.0
20-24	9,236	45.1	36.9	13.4	3.6	1.0
25-29	7,911	29.5	36.0	21.9	8.5	4.1
30-34	4,298	22.2	31.9	23.1	12.3	10.4
35-39	1,506	18.1	23.5	21.0	16.7	20.7
40+	241	14.9	12.4	17.0	16.2	39.4
1984						
TOTAL	27,350	39.8%	32.2%	16.6%	6.7%	4.8%
< 15	67	97.0	3.0	--	--	--
15-19	4,230	76.9	19.5	3.2	0.4	0.1
20-24	9,396	44.2	36.7	14.1	3.9	1.1
25-29	7,846	28.9	36.3	21.9	8.4	4.4
30-34	4,246	21.5	31.3	24.3	12.5	10.5
35-39	1,338	15.1	23.0	21.7	16.3	23.8
40+	227	13.7	15.4	13.2	15.9	41.9
1983						
TOTAL	27,508	40.6%	31.6%	15.8%	6.7%	5.4%
< 15	82	95.1	3.7	1.2	--	--
15-19	4,681	77.6	18.8	3.1	0.4	0.0
20-24	9,429	45.1	36.9	13.0	3.9	1.0
25-29	7,650	28.4	36.4	21.9	8.1	5.2
30-34	4,112	20.4	29.4	23.9	13.4	12.9
35-39	1,323	12.5	21.8	20.5	18.4	26.8
40+	231	9.1	12.1	19.9	13.4	45.5
1982						
TOTAL	27,630	40.6%	31.4%	16.3%	6.6%	5.2%
< 15	61	96.7	3.3	--	--	--
15-19	4,741	77.8	19.1	2.7	0.4	0.1
20-24	9,562	44.9	36.8	14.0	3.3	1.0
25-29	7,774	28.5	36.2	21.9	8.9	4.5
30-34	4,056	18.9	28.3	25.7	13.9	13.3
35-39	1,207	13.5	20.7	21.3	16.7	27.8
40+	229	16.2	11.8	12.2	12.2	47.6
1981						
TOTAL	26,565	41.9%	30.8%	15.7%	6.3%	5.2%
< 15	78	96.2	3.8	--	--	--
15-19	4,583	78.0	19.3	2.1	0.4	0.1
20-24	9,293	46.8	35.9	13.1	3.3	0.9
25-29	7,515	29.9	35.7	21.8	8.1	4.5
30-34	3,848	19.3	27.9	25.4	13.7	13.7
35-39	1,029	12.5	18.1	21.7	18.0	29.7
40+	219	8.7	9.1	13.2	16.4	52.5
1980						
TOTAL	26,589	43.0%	29.9%	15.3%	6.2%	5.5%
< 15	66	95.5	4.5	--	--	--
15-19	4,758	79.0	18.3	2.3	0.4	0.0
20-24	9,541	47.8	35.1	13.2	3.0	0.9
25-29	7,331	30.3	34.3	22.4	8.1	4.9
30-34	3,671	18.2	28.4	23.6	14.9	14.8
35-39	992	13.7	15.1	18.1	17.1	35.9
40+	230	13.0	11.7	7.8	17.4	50.0
1970						
TOTAL	22,004	38.3%	25.8%	14.2%	8.3%	13.4%
< 15	36	96.2	3.8	--	--	--
15-19	4,026	80.9	16.5	2.4	0.1	0.1
20-24	8,486	45.2	35.8	12.6	4.4	2.0
25-29	5,271	18.7	28.2	24.4	14.7	14.0
30-34	2,550	7.5	15.2	20.2	18.2	38.9
35-39	1,234	4.4	7.6	11.4	15.5	61.1
40+	401	0.5	3.6	8.7	10.7	76.4
1960						
TOTAL	30,747	25.3%	22.6%	18.1%	12.2%	21.8%
< 15	44	100.0	--	--	--	--
15-19	4,957	68.5	24.7	5.5	1.0	0.3
20-24	10,833	30.2	33.2	21.0	10.2	5.4
25-29	7,362	10.2	20.1	25.1	19.1	25.5
30-34	4,386	5.6	10.6	17.4	17.3	49.1
35-39	2,459	4.5	6.0	13.0	15.3	61.1
40+	706	4.1	6.2	10.6	5.4	73.7

TABLE 2.8
NEW MEXICO RESIDENTS 1985
UNITED STATES 1984
LIVE BIRTHS
Percentage Distribution By Live Birth Order By Age of Mother

AGE OF MOTHER	NUMBER OF BIRTHS	PERCENTAGE DISTRIBUTION BY LIVE BIRTH ORDER 1	2	3	4	5+
UNITED STATES (All Races)						
TOTAL:	XXX	41.9	33.1	15.4	5.7	3.9
< 15	XXX	97.2	2.5	0.2	0.0	--
15-17	XXX	87.0	11.8	1.1	0.1	0.0
18-19	XXX	72.0	22.8	4.4	0.7	0.1
(15-19)	XXX	(77.3)	(18.9)	(3.2)	(0.5)	(0.1)
20-24	XXX	48.8	34.8	12.1	3.2	1.0
25-29	XXX	35.0	37.3	18.2	6.3	3.2
30-34	XXX	24.3	35.7	22.8	9.8	7.4
35-39	XXX	17.9	27.6	23.2	13.9	17.3
40 +	XXX	12.2	16.2	17.7	14.9	39.0
NEW MEXICO (All Races)						
TOTAL:	27,449	40.4	32.1	16.2	6.7	4.6
< 15	63	100.0	--	--	--	--
15-17	1,469	88.4	10.7	0.9	0.1	--
18-19	2,725	71.7	23.6	4.3	0.3	0.0
(15-19)	(4,194)	(77.6)	(19.1)	(3.1)	(0.2)	(0.0)
20-24	9,236	45.1	36.9	13.4	3.6	1.0
25-29	7,911	29.5	36.0	21.9	8.5	4.1
30-34	4,298	22.2	31.9	23.1	12.3	10.4
35-39	1,506	18.1	23.5	21.0	16.7	20.7
40 +	241	14.9	12.4	17.0	16.2	39.4
NEW MEXICO (White Non-Hispanic)						
TOTAL:	10,250	42.8	35.0	15.0	4.7	2.5
< 15	9	100.0	--	--	--	--
15-17	316	92.1	7.3	0.6	--	--
18-19	718	74.0	23.0	2.9	0.1	--
(15-19)	(1,034)	(79.5)	(18.2)	(2.2)	(0.1)	(--)
20-24	3,189	50.2	36.6	10.4	2.4	0.3
25-29	3,335	35.7	38.4	18.9	5.1	1.9
30-34	1,986	29.3	37.5	20.2	8.3	4.6
35-39	600	27.0	31.3	21.2	9.8	10.7
40 +	97	21.6	17.5	19.6	12.4	28.9
NEW MEXICO (White Hispanic)						
TOTAL:	12,370	40.2	31.5	16.6	7.0	4.7
< 15	36	100.0	--	--	--	--
15-17	881	87.5	11.6	0.8	0.1	--
18-19	1,468	70.0	24.4	5.1	0.5	0.1
(15-19)	(2,349)	(76.5)	(19.6)	(3.5)	(0.3)	(0.0)
20-24	4,431	43.7	36.9	14.2	3.9	1.3
25-29	3,313	25.6	36.0	23.7	10.0	4.7
30-34	1,565	17.3	30.2	26.3	14.4	11.8
35-39	581	13.1	21.7	21.7	20.3	23.2
40 +	95	12.6	8.4	18.9	18.9	41.1
NEW MEXICO (Black)						
TOTAL:	675	42.4	31.1	17.2	6.2	3.1
< 15	6	100.0	--	--	--	--
15-17	36	72.2	27.8	--	--	--
18-19	94	71.3	26.6	2.1	--	--
(15-19)	(130)	(71.5)	(26.9)	(1.5)	(--)	(--)
20-24	255	43.1	33.7	17.6	5.1	0.4
25-29	168	31.0	36.3	21.4	7.7	3.6
30-34	87	23.0	26.4	28.7	11.5	10.3
35-39	29	17.2	17.2	27.6	20.7	17.2
40 +	0	--	--	--	--	--
NEW MEXICO (Indian)						
TOTAL:	3,805	33.5	26.4	18.2	11.0	10.9
< 15	12	100.0	--	--	--	--
15-17	232	88.8	9.5	1.7	--	--
18-19	429	74.1	21.4	4.2	0.2	--
(15-19)	(661)	(79.3)	(17.2)	(3.3)	(0.2)	(--)
20-24	1,262	36.8	38.4	17.5	5.4	1.9
25-29	965	18.7	28.4	26.7	15.9	10.4
30-34	583	10.8	17.5	23.3	21.1	27.3
35-39	274	9.5	9.1	19.3	24.1	38.0
40 +	48	6.3	10.4	6.3	18.8	58.3
NEW MEXICO (Other)						
TOTAL:	349	43.6	34.1	16.0	4.6	1.7
< 15	0	--	--	--	--	--
15-17	4	100.0	--	--	--	--
18-19	16	75.0	25.0	--	--	--
(15-19)	(20)	(80.0)	(20.0)	(--)	(--)	(--)
20-24	99	53.5	34.3	12.1	--	--
25-29	130	46.2	31.5	16.2	5.4	0.8
30-34	77	24.7	39.0	24.7	9.1	2.6
35-39	22	18.2	45.5	13.6	9.1	13.6
40 +	1	--	--	100.0	--	--

The percentages of first and second order live births to New Mexico women are slightly below corresponding national figures and higher for third and higher order live births. The percentage of first live birth is higher for the State's non-Hispanic White and Black women than nationally, but lower for Indian and Hispanic women in the State (Table 2.8 previous page).

The variations by race and age are notable. For example, the percentage of first order live births to Indian women 25-29 years is just over half the percentage for women 25-29 nationally.

Birth order by county shows quite a variation in the State. A detailed breakdown is presented in Table A-5 of the Appendix for 1985 data.

In 1985, 16,372 of New Mexico's 27,449 live births were single births of second or higher order. Of these births, about 1% had been preceded by the birth of a sibling that same year. Sixty-five percent of second and higher order births in 1985 were to mothers who had last given birth to a live born infant during the preceding five years (1980-1984) - 1 out of 5 (21%) of these births had occurred two years prior in 1983, as shown in Figure 2.6. For 1 out of 5 births of second and higher order, the year of the last previous live birth was not stated.

FIGURE 2.6
PERCENT OF LIVE BIRTHS
BY YEAR OF LAST LIVE BIRTH
SECOND ORDER AND HIGHER SINGLE BIRTHS
NEW MEXICO RESIDENTS
1985



BIRTHS TO SINGLE MOTHERS

Continuing the upward trend of the Sixties and Seventies the percentage of births to single mothers in New Mexico rose to 19.8% in 1980, and further still to 26.4% in 1985. As noted in Figure 2.7 the percentage of births to single mothers in the State has exceeded the national rate since 1965. The national rate was 18.4% in 1980 and 21.0% in 1984. The gap in the percent of births to single mothers between New Mexico and the rest of the nation has been wider since 1981 than in prior years (Figure 2.7).

FIGURE 2.7
BIRTHS TO SINGLE MOTHERS
PERCENTAGE OF TOTAL BIRTHS
NEW MEXICO RESIDENTS AND UNITED STATES
1960-1985



★(1985 U.S. DATA NOT AVAILABLE AT PRINTING)

The percentage of births to single mothers ranged from the highest in McKinley County (52.0%) to the lowest in Los Alamos County (5.9%) in 1985 (Table 2.9).

Most of the State's counties experienced increases in the percentage of births to single mothers in 1985 as compared with 1984. Comparison by county of 1985 data with that for earlier periods generally reflects the overall increasing trend in the State.

TABLE 2.9
NEW MEXICO
BIRTHS TO SINGLE MOTHERS
PERCENTAGE OF TOTAL BIRTHS
BY COUNTY OF RESIDENCE

HEALTH DISTRICTS	COUNTY	1973-75	1976-78	1980	1981	1982	1983	1984	1985
	NEW MEXICO	14.8%	16.9%	19.8%	21.6%	22.1%	23.1%	24.6%	26.4%
	CIBOLA	*	*	*	*	27.7	34.7	36.8	36.4
1	MCKINLEY	26.0	30.3	35.8	39.6	42.3	43.6	47.4	52.0
	SAN JUAN	17.4	19.8	21.3	21.8	22.8	25.0	28.3	28.6
	COLFAX	15.5	15.8	20.9	21.1	21.0	19.2	22.2	23.8
	HARDING	10.0	9.3	7.7	11.8	8.3	18.8	9.1	12.5
	LOS ALAMOS	3.7	3.9	7.0	8.3	4.1	4.9	5.0	5.9
	MORA	25.9	22.7	34.4	25.3	30.1	24.7	19.7	37.5
2	RIO ARRIBA	18.4	19.6	27.5	27.6	29.1	28.4	31.2	36.5
	SAN MIGUEL	19.7	23.6	27.4	33.1	33.2	37.1	33.1	31.3
	SANTA FE	13.7	17.8	16.6	21.2	22.9	22.3	23.7	26.6
	TAOS	19.9	20.1	16.1	16.7	22.2	23.1	26.4	26.2
	UNION	11.5	21.5	14.7	15.4	12.4	17.5	26.2	15.8
	BERNALILLO	14.9	17.0	20.7	23.1	22.8	23.1	24.5	26.0
	SANDOVAL	29.5	29.9	34.3	32.4	32.4	30.3	32.5	28.2
3	TORRANCE	11.0	11.0	19.2	16.5	19.8	18.5	13.2	14.7
	VALENCIA	13.5	14.1	17.9	18.9	22.8	20.8	19.0	21.3
	CHAVES	14.8	18.2	19.1	22.2	18.5	17.9	20.4	24.0
	CURRY	9.7	11.9	13.1	17.3	14.2	19.4	16.9	20.6
	DE BACA	9.8	12.2	17.1	5.7	27.5	11.9	28.9	16.7
	EDDY	10.2	12.0	14.1	15.6	12.8	15.1	18.4	20.5
4	GUADALUPE	24.3	23.0	29.5	38.6	36.0	37.0	33.3	30.8
	LEA	11.4	12.0	12.6	11.5	12.7	16.8	14.5	16.5
	LINCOLN	10.1	9.9	14.8	15.3	10.9	16.1	14.6	17.1
	QUAY	12.2	13.9	20.1	26.8	19.4	15.8	22.4	33.0
	ROOSEVELT	8.1	14.2	16.2	18.2	18.8	18.0	18.7	17.8
	CATRON	3.3	5.0	8.2	4.9	15.0	16.2	3.3	8.8
	DONA ANA	11.8	12.2	15.1	15.5	18.1	17.9	19.6	23.8
	GRANT	9.3	13.6	17.0	15.4	18.1	21.8	22.8	24.1
5	HIDALGO	13.1	14.5	15.8	18.8	16.2	20.8	21.3	26.2
	LUNA	12.8	14.3	13.5	20.3	20.3	24.3	24.4	26.8
	OTERO	8.4	9.6	11.2	13.8	15.6	16.0	17.4	15.3
	SIERRA	11.0	15.7	20.9	18.3	20.8	19.5	24.2	18.4
	SOCORRO	11.7	12.8	15.0	23.6	16.0	18.1	21.3	20.6

*Prior to 1982, Cibola County data were included with Valencia County.

As Figure 2.8 shows, in 1985 the percentage of births to single mothers increased over the previous year for the State's race/ethnic groups, except for Indian women, which remained unchanged from 1984. The percentage of births to single women was highest for Indian women (just over half of all births) and second highest for Black women (slightly more than one-third of all births) in 1985. For Hispanic White women the figure was more than one in four, for non-Hispanic White women the figure was one in seven births to single mothers.

As shown in Figure 2.9, it is for women 19 and younger that the highest percentages of births to single mothers is noted. The percentage has been lowest since 1978 for women in the 30-34 age group.

The percentage of births to single mothers increased in 1985 over the previous year for all age groups of women in the State, except for the less than 15 age group.

Because there are less than one hundred births a year to New Mexico women under fifteen years of age, the percentages show marked fluctuation from year to year, yet the percentage of births to single mothers in this age group remains the highest of any age group (Figure 2.9).

Among the State's four largest race/ethnic categories the 1985 percentage of births to single mothers was lowest for the State's non-Hispanic White women in most age groups, reaching a minimum for the 30-34 age group (Table 2.10).

FIGURE 2.8
BIRTHS TO SINGLE MOTHERS
PERCENTAGE BY ETHNIC/RACE
NEW MEXICO RESIDENTS
1980-1985



In 1985 for Indians the 35-39 age group had the lowest percentage births to single mothers; for Blacks and Hispanic Whites the 30-34 age group had lowest percentages.

The highest percentage of births to single mothers for each age group, most notably for ages under 20, occurred to Indian women, especially the Pueblo.

FIGURE 2.9
BIRTHS TO SINGLE MOTHERS
PERCENTAGE BY AGE GROUP, BY YEAR
NEW MEXICO RESIDENTS
1970-1985



TABLE 2.10
NEW MEXICO BIRTHS TO SINGLE MOTHERS, 1985
Percentage by Age of Mother and Ethnic/Race

ETHNIC/RACE	ALL AGES	PERCENTAGE BY AGE OF MOTHER < 15	15-19	20-24	25-29	30-34	35-39	40+
ALL RACES	26.4	90.5	57.1	28.7	16.5	12.6	17.4	21.2
WHITE:	22.2	88.9	52.0	24.1	12.7	9.5	14.8	17.2
Non-Hispanic	13.7	88.9	41.2	16.1	7.6	6.0	11.3	13.4
Hispanic	29.3	88.9	56.7	29.9	17.9	14.1	18.4	21.1
BLACK:	38.1	83.3	73.1	38.0	24.4	14.9	20.7	--
INDIAN:	51.1	100.0	80.5	55.6	42.7	32.4	28.8	35.4
Pueblo	59.0	100.0	84.9	64.0	47.8	36.8	27.5	28.6
Navajo	50.3	100.0	81.0	56.0	42.4	32.3	30.0	35.9
Apache	46.9	100.0	72.1	42.9	47.2	16.7	27.3	--
Other	34.1	--	60.0	33.7	27.7	30.0	15.4	50.0
OTHER NON-WHITE	8.0	--	50.0	9.1	4.6	0.0	9.1	100.0

Though overall the percentage of births to single mothers in New Mexico was about 17% higher than nationally in 1984, it is interesting to note that it is not among teenage women, but rather for women 20 years of age and older, for whom the State's figures for births to single mothers are higher than nationally (Table 2.11).

TABLE 2.11
BIRTHS TO SINGLE MOTHERS
Percentage By Age Of Mother
NEW MEXICO RESIDENTS 1976-1985
UNITED STATES 1976-1984 ESTIMATES

	PERCENTAGE BY AGE OF MOTHER											
	< 15	15-19						20-24	25-29	30-34	35-39	40+
		TOTAL	15	16	17	18	19					
NEW MEXICO												
1985	90.5	57.1	80.6	70.9	64.0	57.1	45.8	28.7	16.5	12.6	17.4	21.2
1984	92.5	53.0	79.0	68.1	59.4	50.5	43.8	26.6	15.4	11.6	14.3	18.1
1983	86.6	49.5	72.5	65.3	53.3	49.5	39.2	23.9	13.7	11.4	12.1	13.0
1982	88.5	46.6	67.6	65.6	54.2	45.4	34.8	22.5	13.4	10.5	13.5	21.0
1981	78.2	44.6	74.1	57.8	49.9	44.7	33.8	22.1	12.6	11.1	15.5	16.9
1980	84.8	42.7	76.1	56.0	48.0	40.4	32.9	19.8	10.7	9.5	12.0	16.1
1979	73.7	38.7	61.1	49.9	43.6	37.9	30.4	17.9	9.5	9.4	10.7	16.6
1978	78.2	37.1	63.5	52.0	42.8	33.4	28.5	16.9	9.4	8.8	11.3	12.8
1977	71.9	37.1	59.8	51.6	39.2	35.3	28.9	16.2	8.6	9.1	10.4	14.2
1976	74.4	34.8	53.7	50.9	39.3	31.0	25.7	14.8	7.6	7.7	10.0	7.5
UNITED STATES												
1984	91.1	55.6	82.6	72.9	63.3	53.9	43.7	24.5	11.7	9.0	10.7	13.8
1983	90.4	53.4	81.3	71.6	61.5	51.6	41.1	22.9	11.0	8.6	10.1	13.4
1982	89.2	50.7	80.3	69.3	58.6	48.6	38.6	21.4	10.3	8.2	9.8	13.4
1981	89.2	49.2	79.2	67.4	56.8	47.0	37.1	20.4	9.7	7.8	9.8	12.7
1980	88.7	47.6	77.7	65.5	54.9	45.1	35.7	19.3	9.0	7.4	9.4	12.1
1979	88.8	46.1	76.8	64.6	52.8	43.3	34.0	17.7	7.5	6.1	7.9	10.3
1978	87.3	44.1	74.1	61.9	50.4	41.0	32.2	16.4	6.9	5.6	7.5	9.6
1977	88.2	42.9	74.3	60.5	49.2	39.6	30.2	14.7	6.4	5.3	7.3	9.3
1976	86.4	40.3	71.9	58.6	46.2	36.4	27.8	13.3	5.7	5.4	7.4	8.9

Percentages of births to single mothers by age group are presented in Table 2.12 for the State's counties. Over half of all births to women 20-24 years old in McKinley County were to single mothers; in McKinley and Sandoval counties more than three-fourths of births to women ages 15-19 were to single mothers.

TABLE 2.12
NEW MEXICO BIRTHS TO SINGLE MOTHERS
Percentage by County & Age
1985

HEALTH DISTRICTS

	COUNTY	TOTAL	PERCENTAGE BY AGE GROUP <15	15-19	20-24	25-29	30-34	35-39	>40
	NEW MEXICO	26.4%	90.5%	57.1%	28.7%	16.5%	12.6%	17.4%	21.2%
	CIBOLA	36.4	100.0	60.4	41.2	23.6	22.5	28.1	0.0
1	MCKINLEY	52.0	100.0	84.2	58.2	42.6	29.5	31.4	34.9
	SAN JUAN	28.6	100.0	58.8	30.3	18.6	18.7	18.7	26.3
	COLFAX	23.8	100.0	63.8	22.1	9.5	7.7	--	50.0
	HARDING	12.5	--	0.0	20.0	14.3	0.0	--	--
	LOS ALAMOS	5.9	--	42.9	6.5	5.3	4.0	0.0	0.0
	MORA	37.5	--	66.7	38.5	20.0	37.5	0.0	--
2	RIO ARRIBA	36.5	100.0	68.6	37.2	24.5	17.2	17.5	50.0
	SAN MIGUEL	31.3	100.0	71.7	35.8	18.9	14.3	19.2	25.0
	SANTA FE	26.6	100.0	68.8	31.9	18.0	10.8	12.0	17.4
	TAOS	26.2	100.0	54.1	29.6	22.1	7.8	27.6	37.5
	UNION	15.8	--	50.0	10.0	4.0	0.0	50.0	0.0
	BERNALILLO	26.0	92.3	59.8	31.9	15.5	10.2	17.9	16.7
	SANDOVAL	28.2	100.0	78.1	32.2	18.2	14.4	16.3	33.3
3	TORRANCE	14.7	--	33.3	20.0	6.8	5.9	0.0	0.0
	VALENCIA	21.3	100.0	58.0	21.8	12.4	11.7	9.8	14.3
	CHAVES	24.0	83.3	54.7	22.3	11.5	12.8	16.7	0.0
	CURRY	20.6	100.0	44.1	20.7	11.2	4.4	5.0	0.0
	DE BACA	16.7	--	45.5	9.1	0.0	0.0	0.0	0.0
	EDDY	20.5	100.0	47.4	17.4	10.4	11.1	16.2	14.3
4	GUADALUPE	30.8	--	73.7	25.0	0.0	22.2	0.0	100.0
	LEA	16.5	50.0	34.6	16.6	8.6	7.2	16.7	44.4
	LINCOLN	17.1	--	46.4	16.7	13.9	5.6	11.1	0.0
	QUAY	33.0	--	52.7	22.2	24.4	25.0	40.0	0.0
	ROOSEVELT	17.6	100.0	41.5	16.0	12.2	2.3	15.4	0.0
	CATRON	8.8	--	0.0	27.3	0.0	0.0	0.0	0.0
	DONA ANA	23.8	75.0	49.6	25.7	15.6	11.2	11.0	31.3
	GRANT	24.1	100.0	52.4	21.8	10.6	11.6	27.6	20.0
5	HIDALGO	26.2	--	42.9	26.8	16.1	11.1	66.7	0.0
	LUNA	26.8	--	50.0	22.2	20.5	13.2	16.7	25.0
	OTERO	15.3	100.0	37.8	13.5	9.4	7.7	16.7	0.0
	SIERRA	18.4	--	37.5	23.5	13.8	0.0	0.0	--
	SOCORRO	20.6	--	56.5	17.7	9.2	11.1	0.0	0.0

-- No births in this age group.
0.0 No births to single mothers.

LOW BIRTH WEIGHT

For many years the term "premature" was applied to all small newborns even though it was recognized that size at birth was an uncertain indicator of the duration of gestation. In 1961, the Expert Committee on Maternal and Child Health of the World Health Organization urged that the term "Low Birth Weight" be used only for infants weighing 2500 grams or less (5½ pounds or less) at birth, and the adjective "premature" be reserved for infants born before the 37th week of gestation. In 1980 New Mexico adopted the slightly revised definition established in the Ninth Revision of the International Classification of Diseases, in which low birth weight is defined as *less than 2500 grams.*

Numerous studies indicate that low birth weight is a risk factor in growth and development. The low birth weight newborn is subject to a relatively high risk of early death, along with an increased vulnerability to neurological damage.

As shown in Figure 2.10, the incidence of low birth weight babies in New Mexico dropped sharply from 1978 to 1981, rose slightly in 1982 to 7.7%, and in 1983 dropped back to the 1981 level of 7.5% where it remained in 1984. The 1984 New Mexico figure was 12% higher than nationally (6.7%). In 1985, the State incidence of low birth weight dropped to 6.9%, its lowest ever.

FIGURE 2.10
PERCENTAGE LOW BIRTH WEIGHT
NEW MEXICO RESIDENTS & UNITED STATES
1960-1985



✪ 1985 U.S. data not available at printing

The incidence of low birth weight was lowest for mothers in the 25 to 29 and the 30 to 34 age groups nationally and in New Mexico. Higher than national rates are noted in New Mexico for women in all age groups except the <15, 15-19, and 40+ age groups for which the New Mexico rates were only slightly lower than the U.S. rates (Figure 2.11).

The incidence of low birth weight by county and age of mother for 1983-1985 is presented in Table 2.13.

FIGURE 2.11
LOW BIRTH WEIGHT BY MOTHER'S AGE
NEW MEXICO 1983-1985 AGGREGATE
UNITED STATES 1984



TABLE 2.13
NEW MEXICO RESIDENTS
PERCENT LOW BIRTH WEIGHT BY COUNTY
(By Age of Mother)
1983-1985 AGGREGATE

HEALTH DISTRICTS	COUNTY	ALL AGES	<15	15-19	20-24	25-29	30-34	35-39	>40
			PERCENTAGE BY AGE GROUP						
	NEW MEXICO	7.3	12.3	8.9	7.4	6.7	6.4	7.9	7.9
	CIBOLA	8.4	10.0	10.4	7.2	8.5	6.0	12.2	16.7
1	MCKINLEY	6.8	5.9	6.2	6.8	6.6	7.5	6.6	9.7
	SAN JUAN	6.3	9.1	8.3	5.9	6.0	5.1	8.4	4.4
	COLFAX	6.5	0.0	9.1	7.8	3.8	3.8	14.7	0.0
	HARDING	9.3	--	0.0	23.1	0.0	0.0	50.0	--
	LOS ALAMOS	5.4	--	4.2	8.2	3.1	5.3	8.6	9.1
	MORA	12.4	--	7.5	14.5	13.3	13.3	10.0	0.0
2	RIO ARRIBA	9.7	0.0	9.6	10.6	9.9	7.3	12.4	0.0
	SAN MIGUEL	9.3	60.0	14.6	7.2	6.7	9.8	16.7	7.7
	SANTA FE	7.9	0.0	10.0	8.3	7.4	7.4	5.7	8.6
	TAOS	9.3	25.0	13.5	9.6	8.8	7.3	7.3	0.0
	UNION	8.3	--	8.9	8.5	4.9	10.0	25.0	20.0
	BERNALILLO	7.8	15.5	10.0	8.2	7.1	6.6	7.8	6.3
	SANDOVAL	6.3	0.0	7.6	6.6	6.4	5.3	4.8	9.7
3	TORRANCE	8.4	--	6.1	8.3	11.3	6.1	8.0	0.0
	VALENCIA	8.3	0.0	9.7	9.0	7.9	7.6	6.0	0.0
	CHAVES	6.2	15.4	7.8	6.0	5.3	4.9	9.8	11.1
	CURRY	6.2	12.5	7.7	5.3	7.1	4.2	4.9	7.7
	DE BACA	8.6	--	6.7	13.5	9.4	0.0	0.0	0.0
	EDDY	7.8	37.5	8.1	7.6	7.8	8.3	7.0	0.0
4	GUADALUPE	7.7	--	8.6	8.1	11.1	3.3	0.0	0.0
	LEA	7.9	5.6	9.4	8.8	7.0	5.1	5.4	3.8
	LINCOLN	8.0	0.0	9.6	8.1	7.1	9.6	0.0	18.2
	QUAY	6.3	0.0	7.4	3.4	8.2	5.8	16.7	0.0
	ROOSEVELT	5.5	0.0	8.8	4.8	5.6	2.9	2.9	14.3
	CATRON	14.9	--	13.3	20.7	4.5	9.1	36.4	0.0
	DONA ANA	6.1	10.0	8.0	6.5	4.9	5.9	5.1	7.1
	GRANT	7.7	0.0	7.9	7.1	5.2	9.7	12.2	40.0
5	HIDALGO	7.5	0.0	8.0	7.9	4.0	8.1	20.0	0.0
	LUNA	6.7	0.0	7.1	6.1	4.4	8.2	14.3	18.8
	OTERO	5.7	20.0	7.6	5.1	5.4	4.4	11.2	0.0
	SIERRA	8.1	--	5.8	8.6	8.4	8.6	7.7	20.0
	SOCORRO	6.8	0.0	9.7	6.6	8.0	1.5	8.3	25.0

-- No Births in This Age Group
0.0 No Low Birth Weight

The general decrease in the incidence of low birth weight in the State is graphically presented in Figure 2.12. The incidence in New Mexico dropped from 8.7% in 1978 to 6.9% in 1985; a drop of 21%. The overall decrease for 1985 was noted for all race/ethnic groups except Blacks.

FIGURE 2.12
LOW BIRTH WEIGHT BY ETHNIC/RACE
NEW MEXICO RESIDENTS
1980-1985



The incidence of low birth weight is higher for female infants than for male infants in the State, and is highest among Blacks and lowest among all Indians for both male and female infants.

The rate decreased by 10% in 1985 for males, but decreased only 5% for females, resulting overall in an 8% decrease from 1984 (Table 2.14).

TABLE 2.14
NEW MEXICO RESIDENTS
Percentage Low Birth Weight
By Race/Ethnic and Sex
1983-1985

RACE/ETHNIC	MALE			FEMALE			TOTAL		
	1983	1984	1985	1983	1984	1985	1983	1984	1985
ALL RACES	6.8%	6.9%	6.2%	8.3%	8.1%	7.7%	7.5%	7.5%	6.9%
WHITE:	6.8	7.0	6.2	8.4	8.3	8.1	7.6	7.7	7.1
Non-Hispanic	6.1	6.7	5.5	7.6	7.8	7.4	6.8	7.2	6.4
Hispanic	7.4	7.4	6.9	9.0	8.8	8.7	8.2	8.1	7.7
BLACK	12.7	9.7	9.4	12.5	9.0	10.4	12.6	9.4	9.9
INDIAN:	5.7	5.6	5.3	6.7	6.3	5.0	6.2	6.0	5.2
Pueblo	5.6	4.8	4.8	7.5	6.8	6.7	6.5	5.9	5.7
Navajo	5.9	5.5	5.1	6.5	6.1	4.5	6.2	5.8	4.8
Apache	4.9	9.3	4.4	6.2	6.7	2.2	5.5	8.0	3.4
All Other	3.9	7.2	9.2	6.1	5.5	6.4	4.9	6.4	7.9
OTHER NON-WHITE	8.1	8.4	7.1	11.4	7.1	6.1	9.6	7.7	6.6

The incidence of low birth weight by county from 1973 through 1985 is shown below in Table 2.15. Colfax, Harding, Los Alamos, Sandoval, Chaves, Curry, Lincoln, Quay, Grant, Otero, and Sierra counties showed decreases of over thirty percent from 1973 to 1985. Bernalillo County showed a decrease of nineteen percent. Rates in counties with small numbers of births often fluctuated from year to year, most likely as a result of random changes.

TABLE 2.15
NEW MEXICO
PERCENT LOW BIRTH WEIGHT
BY COUNTY OF RESIDENCE
1973-1985

HEALTH DISTRICTS	COUNTY	1973-75	1976-78	1980	1981	1982	1983	1984	1985
	NEW MEXICO	9.0%	8.7%	7.6%	7.5%	7.7%	7.5%	7.5%	6.9%
	CIBOLA	*	*	*	*	6.4	9.4	9.4	6.3
1	MCKINLEY	9.5	9.2	7.5	8.1	7.4	6.6	6.9	6.9
	SAN JUAN	7.4	6.9	6.1	6.0	6.6	7.0	5.9	5.8
	COLFAX	10.0	9.6	9.6	8.7	7.7	7.2	8.0	4.7
	HARDING	13.3	11.6	15.4	17.6	8.3	12.5	9.1	6.3
	LOS ALAMOS	11.0	9.9	7.5	5.4	11.7	5.8	4.0	6.4
	MORA	13.9	8.6	11.1	10.1	12.0	11.7	10.5	15.3
2	RIO ARRIBA	10.2	9.9	9.3	10.9	10.9	10.2	9.7	9.4
	SAN MIGUEL	13.2	12.1	9.3	10.6	11.4	8.9	8.9	10.1
	SANTA FE	10.7	10.3	7.2	8.1	8.5	7.8	8.5	7.5
	TAOS	11.9	10.5	6.5	8.7	11.0	8.0	10.6	9.2
	UNION	6.5	8.5	10.7	3.3	4.5	10.0	2.4	13.2
	BERNALILLO	9.4	9.0	8.5	7.4	8.3	8.2	7.7	7.6
	SANDOVAL	9.3	8.9	8.0	7.2	5.8	7.3	5.7	6.1
3	TORRANCE	11.0	7.0	10.0	5.5	4.0	11.6	4.4	8.8
	VALENCIA	8.5	8.7	8.1	7.0	6.4	8.1	8.7	7.9
	CHAVES	9.3	8.7	7.1	7.4	7.0	6.0	7.0	5.7
	CURRY	8.8	9.0	7.2	6.5	7.9	7.1	6.2	5.2
	DE BACA	4.3	0.0	9.8	2.9	7.5	9.5	10.5	5.6
	EDDY	7.1	7.5	6.6	6.3	6.1	7.9	8.4	7.1
4	GUADALUPE	4.6	7.0	11.6	9.4	5.6	7.6	9.0	6.4
	LEA	8.5	9.3	8.9	8.7	7.5	7.7	8.4	7.6
	LINCOLN	10.3	8.1	7.7	11.6	10.0	10.6	6.5	6.9
	QUAY	9.6	7.8	6.1	7.0	7.0	7.3	5.5	6.3
	ROOSEVELT	6.3	8.3	4.5	6.2	7.3	4.8	5.6	6.3
	CATRON	10.0	8.5	6.1	12.2	10.0	16.2	20.0	8.8
	DONA ANA	7.6	7.2	6.2	6.4	7.3	5.8	6.7	5.7
	GRANT	9.3	10.9	7.3	9.4	7.9	8.2	8.7	6.4
5	HIDALGO	7.3	8.4	6.5	5.4	5.6	8.3	7.4	6.5
	LUNA	6.9	8.0	7.1	4.4	9.5	4.7	7.2	8.3
	OTERO	7.6	7.1	5.7	6.5	6.3	6.1	6.6	4.4
	SIERRA	7.6	10.4	8.8	3.8	6.3	9.4	12.6	2.0
	SOCORRO	6.4	9.0	8.5	9.5	7.4	5.8	9.2	5.6

*Prior to 1982, Cibola County data were included with Valencia County

Figure 2.13 shows the percent low birth weight by county of residence for 1983-1985. Many of the counties with percentage of birth weight above the state average were in the north central portion of the State.

FIGURE 2.13
NEW MEXICO RESIDENT LIVE BIRTHS PERCENT LOW BIRTH WEIGHT BY COUNTY 1983-1985 AGGREGATE


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
LOS ALAMOS
MORA
HARDING
McKINLEY
SANDOVAL
SANTA FE
SAN MIGUEL
CIBOLA
BERNALILLO
GUADALUPE
QUAY
VALENCIA
TORRANCE
CURRY
DE BACA
CATRON
SOCORRO
ROOSEVELT
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
PERCENT LOW BIRTH WEIGHT
≥9.0%
8.0-8.9%
7.0-7.9%
6.0-6.9%
<6.0%

Birth weight distribution for New Mexico and the United States is shown in Table 2.16. New Mexico closely follows the United States in the percentage distribution by birth weight, with a larger proportion of births in the 2000-3499 gram range and a correspondingly smaller proportion of births with 3500 grams and higher birth weight.

TABLE 2.16
PERCENT OF LIVE BIRTHS BY BIRTH WEIGHT DISTRIBUTION NEW MEXICO RESIDENTS 1985 UNITED STATES 1984

BIRTH WEIGHT (IN GRAMS)	NEW MEXICO	UNITED STATES
< 1000	0.4	0.6%
1000-1499	0.6	0.6
1500-1999	1.1	1.3
2000-2499	4.8	4.2
All < 2500 gms.	6.9	6.7
2500-2999	20.3	15.9
3000-3499	39.9	36.7
3500-3999	24.6	29.5
4000-4499	5.7	9.1
All 2500-4499 gms	90.5	91.2
4500-4999	0.9	1.7
5000 +	0.2	0.2
All 4500+ gms.	1.1	1.9
UNSTATED	1.5	0.1

Infant mortality, especially during the first month of life, varies with birth weight, with weights of 3000-4499 grams associated with the best chance for a healthy survival: this included in New Mexico 69% of all live births and 75% nationally in 1984. For New Mexico in 1985 this percentage was 70%.

Birth weight distribution by race and ethnicity for New Mexico is shown in Figure 2.14. The largest category of 1985 New Mexico births (3000-3499 grams) included, from among the States four major ethnic groups, from 39.1% of all births to non-Hispanic Whites to 41.9% of all births to Indians; the second largest category (3500-4499 grams birth weight) included from 23.8% of births to Blacks to 34.3% to non-Hispanic Whites. In contrast for the United States, the largest category was 3500-4499 grams birth weight with 38.7% and the second largest category was 3000-3499 grams with 36.7% in 1984.

FIGURE 2.14
PERCENTAGE DISTRIBUTION OF LIVE BIRTHS BY BIRTH WEIGHT 1985 NEW MEXICO RESIDENTS BY RACE/ETHNICITY 1984 UNITED STATES



GESTATIONAL AGE

Gestation-the period of the development of the fetus from conception until birth-normally takes place in 37 to 41 weeks. Most live births were "term" (born at 37 to 41 weeks gestation) both nationally (73%) and statewide (61%) in 1984: 16% fewer term births in the state than nationally. In 1985 the percentage of term births in the State decreased to 55%.

TABLE 2.17
PERCENT OF LIVE BIRTHS BY WEEKS OF GESTATION NEW MEXICO RESIDENTS 1984-1985 UNITED STATES 1983-1984

LENGTH OF GESTATION	NEW MEXICO		UNITED STATES	
(In Weeks)	1984	1985	1983	1984
< 28	.4	.4	.7	.7
28-31	1.0	.9	1.1	1.1
32-36	11.2	7.4	7.4	7.3
ALL =< 36	12.7	8.6	9.3	9.1
37-39	23.0	23.0	36.5	36.2
40	28.0	17.8	21.3	21.6
41	9.8	14.0	14.9	15.0
ALL 37-41	60.9	54.8	73.0	72.8
=> 42	9.9	14.6	14.4	14.6
UNSTATED/UNKNOWN	16.5	22.0	3.6	3.6

The ten percent decrease in the reported percentage of term births from 1984 to 1985 in the State was probably due to an increase of one third in the percentage of births with unknown/unstated weeks gestation. The 1984 New Mexico percentage unknown/unstated was nearly five times greater than that for the U.S. (Table 2.17).

CONGENITAL ANOMALIES

According to information reported on birth certificates, congenital anomalies occurred with 1.35% of 1981-1985 New Mexico resident live births and 0.90% of 1983 United States live births (1983 data on live births with congenital anomalies is the most recently available for the U.S.).

Anomalies of the musculo-skeletal system—including congenitally dislocated hip, deformities of the feet, polydactyly (extra toes or fingers) syndactyly (webbed toes or fingers), and several other related deformities—accounted for over a third of all anomalies occurring in 1981-1985 New Mexico live births. As shown in Figure 2.15, other organ systems were affected much less frequently by congenital anomalies.

FIGURE 2.15
NEW MEXICO RESIDENT
LIVE BIRTHS WITH CONGENITAL ANOMALIES
PERCENT BY TYPE OF ANOMALY
1981-1985 AGGREGATE



The reported overall rate as well as the rates of certain types of congenital anomalies were higher for New Mexico (1981-1985) resident live births than for live births nationally (1983). Most notably, rates were two times greater in New Mexico than nationally for congenital anomalies of the ear, four-and-one-half times greater for patent ductus arteriosus, and three times greater for congenital dislocation of the hip (Table 2.18). Conversely, New Mexico's rates for births reporting anencephalus, spina bifida, and syndactyly were twenty to forty percent lower than national rates. The New Mexico rate for congenital anomalies of the integument was much higher than the national rates.

In examining the data on live births with reported congenital anomalies, the reader should note that congenital anomalies are under-reported by up to eighty percent, depending on the research cited. However, the available data still provide useful and interesting information.

TABLE 2.18
LIVE BIRTHS WITH CONGENITAL ANOMALIES
RATE PER 100,000 LIVE BIRTHS
BY SELECTED TYPES OF ANOMALIES*
NEW MEXICO RESIDENT 1981-1985 AGGREGATE
UNITED STATES 1983

TYPE OF CONGENITAL ANOMALY (ICD-9 CATEGORIES 740-759)	NEW MEXICO	UNITED STATES
CONGENITAL ANOMALIES OF THE NERVOUS SYSTEM	68.1	82.3
Anencephalus and Similar Anomalies	(8.8)	(15.2)
Spina Bifida	(19.8)	(31.9)
Congenital Hydrocephulus	(16.8)	(19.5)
CONGENITAL ANOMALIES OF THE EAR, FACE, NECK	65.9	31.0
Congenital Anomalies of the Ear	(54.9)	(24.0)
CONGENITAL ANOMALIES OF THE HEART	56.4	55.2
OTHER CONGENITAL ANOMALIES OF THE CIRCULATORY SYSTEM	43.2	23.9
Patent Ductus Arteriosis	(23.4)	(5.1)
Absence or Hypoplasia of Umbilical Artery	(12.5)	(11.9)
CLEFT PALATE AND CLEFT LIP	101.1	84.9
Cleft Palate	(31.5)	(26.3)
Cleft Lip	(29.3)	(21.0)
Cleft Palate with Cleft Lip	(40.3)	(37.6)
OTHER CONGENITAL ANOMALIES OF THE DIGESTIVE SYSTEM**	43.2	47.1
CONGENITAL ANOMALIES OF THE GENITAL ORGANS	110.6	107.8
Undescended Testicle	(35.2)	(26.4)
Hypospadias and Epispadias	(52.7)	(63.3)
MUSCULOSKELETAL DEFORMITIES	501.1	367.6
Congenital Dislocation of the Hip	(101.8)	(34.9)
Deformities of the Feet	(94.5)	(74.6)
Polydactyly	(77.7)	(91.6)
Syndactlyly	(15.4)	(20.7)
Reduction Deformities of the Limbs	(20.5)	(22.6)
CONGENITAL ANOMALIES OF THE INTEGUMENT	143.6	81.7
Congenital Anomalies of the Skin	(132.6)	(70.4)
CHROMOSOMAL ANOMALIES	53.5	47.1
Down's Syndrome	(46.2)	(35.9)
MULTIPLE CONGENITAL ANOMALIES, SO DESCRIBED	86.4	24.4
OTHER AND UNSPECIFIED CONGENITAL ANOMALIES***	79.9	70.1
ALL LIVE BIRTHS WITH CONGENITAL ANOMALIES REPORTED	1 ,353.1	898.6

* Individual categories for the U.S. exceed the total for U.S., since each condition listed on a birth certificate was coded. In contrast, space limitations on the computerized record permitted only one entry for Congenital Anomaly data for New Mexico births. In the event that more than one condition was listed for a New Mexico birth, the code for "multiple congenital anomalies, so described" (759.7) would be assigned if more than one system was affected; if only one system was affected the code used would reflect the most severe condition.

** Includes ICD-9 categories 750 and 751

*** Includes ICD-9 categories 743, 748, 753, 759.0-759.6, 759.8-759.9

The incidence of congenital anomalies in live births to each of New Mexico's ethnic and racial groups exceeded the national rates. The percent of live births with reported congenital anomalies was higher among Blacks (1.75%) than any other race in New Mexico during 1981-1985, and nearly double the incidence nationally among Blacks (0.95%) during 1983 (Figure 2.16).

The percentage of live births with congenital anomalies was higher for all age groups of mothers in New Mexico than nationally, especially for mothers 19 and younger. As shown in Figure 2.17, both the State and the nation reported the highest percentages for births to mothers 35 years and older. The lowest percentages of births with congenital anomalies occurred to mothers 30 to 34 years of age in New Mexico (1.19%) during 1981-1985, and to U.S. (.78%) mothers 25-29 years of age during 1973-1974 (1973-74 U.S. data by age of mother for live births with congenital anomalies is the most recently available).

FIGURE 2.16
LIVE BIRTHS WITH CONGENITAL ANOMALIES PERCENT REPORTED BY RACE/ETHNIC GROUP NEW MEXICO RESIDENT 1981-1985 AGGREGATE UNITED STATES 1983



FIGURE 2.17
LIVE BIRTHS WITH CONGENITAL ANOMALIES PERCENT REPORTED BY AGE OF MOTHER NEW MEXICO RESIDENT 1981-1985 AGGREGATE UNITED STATES 1973-1974 AGGREGATE



Both nationally and Statewide the incidence of live births with congenital anomalies was higher for male births than for female births. The percent of male births with congenital anomalies was 1.46% for New Mexico in 1981-1985 and 1.01% for the United States in 1983; the percent of female births with congenital anomalies was 1.24% for New Mexico and 0.78% nationally for the same reporting periods.

Table 2.19 presents by county of residence the number and percent of live births with congenital anomalies. These figures vary each year from county to county. In counties averaging over a thousand births annually from 1981-1985, the lowest reported percentage occurred to Dona Ana County residents with less than .3% in any year. Chaves, San Juan, and Bernalillo counties showed some of the highest percentages. Bernalillo County accounted for 35% of the State's live births with congenital anomalies in 1981 and 31% in 1985.

TABLE 2.19
LIVE BIRTHS WITH CONGENITAL ANOMALIES NUMBER AND PERCENT REPORTED BY COUNTY OF RESIDENCE 1981-1985

	1981		1982		1983		1984		1985	
COUNTY	NUMBER	PERCENT	NUMBER	PERCENT	NUMBER	PERCENT	NUMBER	PERCENT	NUMBER	PERCENT
NEW MEXICO	332	1.25%	373	1.35%	449	1.63%	369	1.35%	324	1.18%
CIBOLA	**	**	9	1.36	2	.35	4	.75	9	1.71
McKINLEY	18	1.08	19	1.10	16	.96	20	1.20	9	.52
SAN JUAN	26	1.47	26	1.45	30	1.30	40	1.79	47	2.11
COLFAX	2	.92	1	.48	3	1.20	1	.47	1	.39
HARDING	0	—	1	8.33	0	—	0	—	0	—
LOS ALAMOS	0	—	4	2.02	1	.49	0	—	1	.49
MORA	2	2.53	2	2.61	2	2.60	0	—	2	2.78
RIO ARRIBA	9	1.56	16	2.47	15	2.22	12	1.64	9	1.19
SAN MIGUEL	7	1.48	13	2.79	5	1.04	7	1.41	6	1.28
SANTA FE	14	1.08	22	1.60	16	1.17	10	.70	13	.90
TAOS	1	.24	2	.45	4	.89	8	1.81	8	1.75
UNION	2	3.33	1	1.52	0	—	1	1.19	2	2.63
BERNALILLO	117	1.59	123	1.60	206	2.59	155	1.99	101	1.27
SANDOVAL	12	1.66	10	1.25	13	1.62	7	.78	9	.92
TORRANCE	1	.79	0	—	2	1.27	1	.74	3	2.21
VALENCIA	19	1.45	13	2.19	13	2.02	10	1.59	7	1.11
CHAVES	22	2.13	4	.35	31	2.64	5	.47	4	.40
CURRY	11	1.22	5	.57	12	1.33	18	1.84	6	.67
DE BACA	0	—	1	3.50	2	4.76	0	—	1	2.78
EDDY	14	1.45	30	2.58	20	1.96	17	1.83	27	3.05
GUADALUPE	0	—	2	2.25	1	1.09	1	1.28	1	1.28
LEA	10	.66	12	.74	11	.69	8	.58	18	1.32
LINCOLN	4	1.85	6	2.51	11	4.23	6	2.44	5	2.02
QUAY	2	1.41	0	—	3	1.82	5	2.73	0	—
ROOSEVELT	15	5.15	8	2.42	2	.68	5	1.87	7	2.45
CATRON	0	—	0	—	0	—	2	6.67	1	2.94
DONA ANA	3	.15	1	.05	5	.24	3	.13	2	.09
GRANT	4	.67	2	.37	3	.43	6	1.29	8	1.55
HIDALGO	0	—	0	—	0	—	0	—	1	.92
LUNA	2	.68	6	1.85	4	1.25	2	.69	3	.99
OTERO	8	.91	11	1.13	12	1.02	8	.74	11	.99
SIERRA	1	1.88	0	—	2	1.56	1	1.05	1	1.02
SOCORRO	2	.73	2	.87	4	1.44	6	2.50	1	.40

* ICD Categories 740-759 only.
** Prior to 1982, data for Cibola County were included with Valencia County.

Combining several years of data can provide some stability to small numbers as in the case with congenital anomalies reported for New Mexico live births by county. Figure 2.18 shows by county of residence the percent of births with congenital anomalies, averaged over a five year period (1981-1985). Reporting the highest percentage of births was Lincoln County (2.66%), followed by Roosevelt (2.52%), Mora (2.33%) and Eddy (2.18%) counties. Of counties with more than 500 births each year, the highest incidence was reported for Eddy County followed by Bernalillo (1.83%) and San Miguel (1.78%) counties. The lowest percentages were reported in Dona Ana (.13%), Hidalgo (.16%) and Los Alamos (.59%) counties.

FIGURE 2.18
NEW MEXICO RESIDENT
LIVE BIRTHS WITH CONGENITAL ANOMALIES
PERCENT REPORTED BY COUNTY
1981-1985 AGGREGATE



DATA FOR CIBOLA AND VALENCIA COUNTIES 1982-1985
(PRIOR TO 1982 CIBOLA COUNTY WAS INCLUDED WITH VALENCIA COUNTY)

PARENT'S LEVEL OF EDUCATION

As shown in Figure 2.19, the majority of 1985 New Mexico resident live births were to parents with high school (12 years) and higher educational attainment. Twenty-four percent of all births were to mothers with less than a high school education (11 years or less), while about 13% of all births were to fathers of that same educational level.

FIGURE 2.19
PERCENT OF LIVE BIRTHS
BY EDUCATION OF MOTHER AND FATHER
NEW MEXICO RESIDENCE
1985



Differences in educational attainment may be related partly to the higher percentage of unknowns for fathers than for mothers. These differences may also result from the interruption in education of a sizeable number of women by early motherhood, especially for those with less than high school educations.

MULTIPLE BIRTHS

In 1985, there were 418 live births in multiple deliveries. The majority of these multiple deliveries (99%) were twin deliveries. One set of triplets was delivered in the state in 1985.

The multiple birth ratio, defined as the number of live births in multiple deliveries per 1,000 total live births was 17.9 in 1984 for New Mexico, and 20.3 nationally. In 1985, this ratio was 15.2 for New Mexico.

HOSPITAL/ATTENDANT

Of all births reported in New Mexico during 1985, 97.9% took place in hospitals, just slightly above the 1984 level of 97.5% (Table 2.20). The State remains lower than the national level of 99.0%.

In examining the data by county, in Table 2.20 it should be kept in mind that drastic changes in the percent of hospital births over the years in some counties, reflect the opening and closing of facilities and ensuing migration across county boundaries in facility utilization.

TABLE 2.20
NEW MEXICO
PERCENTAGE OF LIVE BIRTHS IN HOSPITAL
1956, 1966, 1976, 1980, 1982-1985
By County of Occurrence

HEALTH DISTRICTS	COUNTY	1956	1966	1976	1980	1982	1983	1984	1985
	NEW MEXICO	90.3	97.1	97.8	98.1	97.3	97.4	97.5	97.9
	CIBOLA	*	*	*	*	98.8	98.0	99.0	100.0
1	MCKINLEY	88.4	96.9	98.8	99.5	99.7	99.7	99.6	99.5
	SAN JUAN	94.2	98.4	99.5	99.0	99.3	99.5	98.7	99.4
	COLFAX	95.0	97.2	99.1	99.5	99.6	100.0	96.5	100.0
	HARDING	0.0	0.0	0.0	--	--	--	--	--
	LOS ALAMOS	100.0	100.0	100.0	99.8	99.8	100.0	99.8	99.5
	MORA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
2	RIO ARRIBA	80.7	97.6	88.7	92.4	90.3	88.1	92.3	93.5
	SAN MIGUEL	60.5	77.6	92.4	94.8	92.0	96.3	94.1	94.9
	SANTA FE	91.3	99.2	95.8	94.8	95.2	95.4	96.1	95.9
	TAOS	46.7	90.8	90.2	89.5	81.5	82.6	85.8	81.3
	UNION	68.2	92.1	96.5	100.0	98.7	97.0	95.2	98.5
	BERNALILLO	96.7	99.5	99.2	98.7	97.2	97.5	98.2	98.6
	SANDOVAL	0.0	4.8	0.0	87.4	89.0	88.1	87.3	88.5
3	TORRANCE	57.3	90.4	0.0	0.0	0.0	0.0	0.0	0.0
	VALENCIA	67.2	98.4	98.0	98.5	98.3	96.2	97.0	96.7
	CHAVES	98.8	99.7	99.3	99.4	98.9	99.5	99.7	99.7
	CURRY	94.2	95.9	99.9	99.8	99.2	99.4	99.5	99.5
	DE BACA	97.7	100.0	100.0	100.0	100.0	100.0	96.9	100.0
	EDDY	96.6	98.5	99.6	98.8	99.3	99.4	99.8	99.3
4	GUADALUPE	58.2	97.5	100.0	90.9	95.2	100.0	95.0	100.0
	LEA	99.3	98.1	98.1	98.1	99.2	98.9	98.8	98.8
	LINCOLN	97.7	98.4	91.9	98.1	98.4	98.4	98.2	98.4
	QUAY	99.0	95.4	98.4	95.9	97.8	97.3	96.4	98.7
	ROOSEVELT	97.2	98.6	98.7	99.0	97.9	97.0	97.8	99.2
	CATRON	75.0	75.0	0.0	0.0	0.0	0.0	0.0	0.0
	DONA ANA	81.7	97.1	96.7	97.6	97.3	97.5	94.9	96.3
	GRANT	97.8	99.1	99.0	98.1	92.5	90.4	94.0	96.3
5	HIDALGO	84.9	100.0	60.0	0.0	0.0	0.0	0.0	--
	LUNA	96.9	99.4	98.3	99.3	98.1	98.3	99.6	98.2
	OTERO	92.9	88.5	99.0	98.4	96.6	95.2	94.6	95.7
	SIERRA	90.4	90.4	97.5	93.7	83.9	100.0	33.3	87.5
	SOCORRO	56.4	97.5	99.4	96.0	93.5	97.4	95.5	99.5

* Prior to 1982, Cibola County data were included with Valencia County.

As seen in Table 2.21, residents of many counties go elsewhere to give birth, usually in other counties in the State, but occasionally outside of New Mexico.

TABLE 2.21
1985 NEW MEXICO RESIDENT BIRTHS BY STATE/COUNTY OF OCCURRENCE

HEALTH DISTRICTS	COUNTY	TOTAL RESIDENT BIRTHS	BIRTHS OCCURRING IN: COUNTY OF RESIDENCE NUMBER	PERCENT	OTHER COUNTIES NUMBER	PERCENT	OUT OF STATE NUMBER	PERCENT
	NEW MEXICO	27,449	22,439	81.75%	3,974	14.48%	1,036	3.77%
	CIBOLA	525	258	49.14	267	50.86	0	--
1	MCKINLEY	1,720	1,561	90.76	50	2.91	109	6.34
	SAN JUAN	2,228	1,859	83.44	130	5.83	239	10.73
	COLFAX	256	203	79.30	34	13.28	19	7.42
	HARDING	16	0	--	13	81.25	3	18.75
	LOS ALAMOS	204	181	88.73	21	10.29	2	.98
	MORA	72	2	2.78	70	97.22	0	--
2	RIO ARRIBA	758	294	38.79	457	60.29	7	.92
	SAN MIGUEL	467	406	86.94	60	12.85	1	.21
	SANTA FE	1,449	1,283	88.54	161	11.11	5	.35
	TAOS	458	317	69.21	120	26.20	21	4.59
	UNION	76	49	64.47	10	13.16	17	22.37
	BERNALILLO	7,937	7,846	98.85	78	.98	13	.16
	SANDOVAL	981	92	9.38	885	90.21	4	.41
3	TORRANCE	136	2	1.47	134	98.53	0	--
	VALENCIA	629	138	21.94	489	77.74	2	.32
	CHAVES	1,012	968	95.65	42	4.15	2	.20
	CURRY	893	657	73.57	213	23.85	23	2.58
	DE BACA	36	13	36.11	22	61.11	1	2.78
	EDDY	885	820	92.66	62	7.01	3	.34
4	GUADALUPE	78	30	38.46	48	61.54	0	--
	LEA	1,367	1,200	87.78	32	2.34	135	9.88
	LINCOLN	246	218	88.62	27	10.98	1	.41
	QUAY	176	145	82.39	20	11.36	11	6.25
	ROOSEVELT	286	241	84.27	40	13.99	5	1.75
	CATRON	34	2	5.88	18	52.94	14	41.18
	DONA ANA	2,140	1,766	82.52	30	1.40	344	16.07
	GRANT	515	488	94.76	19	3.69	8	1.55
5	HIDALGO	107	0	--	96	89.72	11	10.28
	LUNA	302	238	78.81	62	20.53	2	.66
	OTERO	1,114	980	87.97	103	9.25	31	2.78
	SIERRA	98	8	8.16	87	88.78	3	3.06
	SOCORRO	248	174	70.16	74	29.84	0	--

Of all births reported in New Mexico during 1985, 91.8% were attended by physicians, a slight decrease from the previous year's 92.7%. The percentage of births attended by nurse-midwives increased from 5.3% in 1984 to 6.3% in 1985, representing a nineteen percent increase (Table 2.22).

TABLE 2.22
NEW MEXICO LIVE BIRTHS PERCENTAGE OCCURRING IN HOSPITAL AND BIRTHS BY ATTENDANT
Distribution By Occurrence, 1965, 1971-1985

YEAR	PERCENTAGE OF BIRTHS: HOSPITAL IN	HOSPITAL OUT OF	ATTENDANT AT BIRTH: PHYSICIAN	LAY MIDWIFE	NURSE MIDWIFE	R.N.	Other
1985	97.9%	2.1%	91.8%	1.0%	6.3%	.1%	.8%
1984	97.5	2.5	92.7	1.1	5.3	.1	.8
1983	97.4	2.6	93.1	.9	4.7	.1	1.3
1982	97.3	2.7	93.7	.8	3.9	.1	1.6
1981	97.8	2.2	92.8	.5	3.9	.9	2.6
1980	98.1	1.9	92.4	.0	5.6	.0	1.9
1979	98.5	1.5	93.3	.0	5.6	.0	1.1
1978	96.7	3.3	94.4	.3	4.5	.0	.8
1977	97.1	2.9	94.8	.4	4.1	.0	.7
1976	97.7	2.3	94.4	.4	4.5	.1	.6
1975	97.8	2.2	94.1	.4	4.9	.0	.6
1974	97.7	2.3	93.5	.7	5.1	.1	.6
1973	98.2	1.8	94.6	.7	4.1	.1	.6
1972	98.1	1.9	94.8	.6	4.0	.1	.5
1971	97.6	2.4	95.9	.7	2.9	.0	.6
1965	96.7	3.3	96.9	1.0	1.1	.1	.9

.0 Equals less than a tenth of a percent.

A detailed listing by county of occurrence, facility, and attendant is presented in Table A-6 of the Appendix.

LEVELS OF PRENATAL CARE

Of the 27,449 New Mexico resident women giving birth to a live born infant in 1985, at least 3,851 (14.0%) received no prenatal care, or very low levels of prenatal care; that is almost 1 out of every 7 of these women. We say "at least," because there were another 4,828 women for whom the level of prenatal care received was unknown (Table 2.23). For women whose level of care was known, only about half received a high or optimum level of care.

In sixteen of the State's counties, at least fifteen percent of women giving birth in 1985 received a low (or no) level of prenatal care. Because of poor reporting in some counties, resulting in a high percentage of unknown, the reader should carefully consider the skewness created by the unknown factor when examining individual counties (Table 2.23).

The level of prenatal care is of grave concern as it impacts so dramatically on the outcome of pregnancy. Mortality during the first twenty seven days of life (neonatal period for infants whose mothers received low levels of prenatal care is six times greater than the mortality noted for infants whose mothers received a high level of care. Overall, infant mortality (all deaths less than one year of age) is over three times higher for infants whose mothers received low levels of prenatal care as compared to those whose mothers received high (optimum) levels of care (refer to section on infant mortality).

TABLE 2.23
NEW MEXICO RESIDENT LIVE BIRTHS BY COUNTY AND LEVEL OF PRENATAL CARE
1985

HEALTH DISTRICTS	COUNTY	LIVE BIRTHS NUMBER	HIGH NO.	%	MID NO.	%	NON/LOW NO.	%	UNKNOWN NO.	%
	NEW MEXICO	27449	10917	39.8	7853	28.6	3851	14.0	4828	17.6
	CIBOLA	525	161	30.7	168	32.0	103	19.6	93	17.7
1	MCKINLEY	1720	594	34.5	579	33.7	363	21.1	184	10.7
	SAN JUAN	2228	1037	46.5	542	24.3	305	13.7	344	15.4
	COLFAX	256	113	44.1	64	25.0	19	7.4	60	23.4
	HARDING	16	6	37.5	5	31.3	1	6.3	4	25.0
	LOS ALAMOS	204	151	74.0	39	19.1	5	2.5	9	4.4
	MORA	72	24	33.3	33	45.8	10	13.9	5	6.9
2	RIO ARRIBA	758	297	39.2	234	30.9	150	19.8	77	10.2
	SAN MIGUEL	467	197	42.2	163	34.9	70	15.0	37	7.9
	SANTA FE	1449	630	43.5	459	31.7	263	18.2	97	6.7
	TAOS	458	152	33.2	102	22.3	54	11.8	150	32.8
	UNION	76	30	39.5	11	14.5	8	10.5	27	35.5
	BERNALILLO	7937	2899	36.5	2296	28.9	886	11.2	1856	23.4
	SANDOVAL	981	437	44.5	287	29.3	140	14.3	117	11.9
3	TORRANCE	136	40	29.4	43	31.6	22	16.2	31	22.8
	VALENCIA	629	229	36.4	202	32.1	87	13.8	111	17.6
	CHAVES	1012	312	30.8	376	37.2	253	25.0	71	7.0
	CURRY	893	308	34.5	162	18.1	70	7.8	353	39.5
	DE BACA	36	6	16.7	9	25.0	9	25.0	12	33.3
	EDDY	885	382	43.2	315	35.6	152	17.2	36	4.1
4	GUADALUPE	78	16	20.5	40	51.3	14	17.9	8	10.3
	LEA	1367	570	41.7	370	27.1	212	15.5	215	15.7
	LINCOLN	246	129	52.4	82	33.3	28	11.4	7	2.8
	QUAY	176	52	29.5	56	31.8	46	26.1	22	12.5
	ROOSEVELT	286	92	32.2	103	36.0	47	16.4	44	15.4
	CATRON	34	3	8.8	9	26.5	4	11.8	18	52.9
	DONA ANA	2140	1121	52.4	325	15.2	250	11.7	444	20.7
	GRANT	515	155	30.1	146	28.3	45	8.7	169	32.8
5	HIDALGO	107	24	22.4	26	24.3	11	10.3	46	43.0
	LUNA	302	78	25.8	82	27.2	68	22.5	74	24.5
	OTERO	1114	531	47.7	401	36.0	102	9.2	80	7.2
	SIERRA	98	48	49.0	25	25.5	16	16.3	9	9.2
	SOCORRO	248	93	37.5	99	39.9	38	15.3	18	7.3

Note should be made of the percentage of "unknown level of care", since a high percentage of unknown (e.g. Curry County) will greatly skew reporting in the other categories.

The percentage of New Mexico women giving birth and having received low levels of prenatal care (examining 1985 data) is high for young mothers, decreasing to its lowest percentage for women in the 30-34 age group (Figure 2.20) and increasing once again with increasing age.

Subdividing the 15-19 age group shows about the same percentage of low level prenatal care for 15-17 year olds as for 18-19 year olds (Table 2.24). This table further details the age breakdown by county, presenting also actual numbers to aid in maintaining a proper perspective. County comparisons are interesting-seven counties exceeded by more than 50% the State percentage for the 30 to 34 age group.

In the 20-24 age group, 340 Bernalillo County resident women who gave birth in 1985 reported having received little or no prenatal care, as did over a hundred women in this age group in McKinley as well as in San Juan and Chaves counties. Of these, the percentage was highest in Chaves County where over a fourth of the women in the 20-24 age group received inadequate care prior to their 1985 delivery.

FIGURE 2.20
NEW MEXICO RESIDENTS
PERCENT BIRTHS WITH LOW LEVEL PRENATAL CARE BY AGE GROUP OF MOTHER
1984 AND 1985



TABLE 2.24
NEW MEXICO
RESIDENT LIVE BIRTHS
WITH LOW LEVEL PRENATAL CARE
NUMBER AND PERCENT BY AGE OF MOTHER
1985

HEALTH DISTRICTS	COUNTY	TOTAL		Number and Percent by Age of Mother																			
				<15		15-17		18-19		(15-19)		20-24		25-29		30-34		35-39		40-44		>45	
		NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%
	NEW MEXICO	3851	14.0	17	27.0	323	22.0	603	22.1	(926)	(22.1)	1399	15.1	848	10.7	437	10.2	191	12.7	29	12.6	4	36.4
	CIBOLA	103	19.6	1	33.3	13	36.1	12	20.0	(25)	(26.0)	32	18.8	27	19.3	13	16.3	5	15.6	0	0.0	0	0.0
1	MCKINLEY	363	21.1	4	66.7	33	28.7	47	26.7	(80)	(27.5)	117	20.6	83	19.2	51	18.3	24	19.8	4	18.2	0	0.0
	SAN JUAN	305	13.7	1	16.7	22	21.8	47	21.5	(69)	(21.6)	107	14.1	67	10.5	37	10.2	21	17.1	3	16.7	0	0.0
	COLFAX	19	7.4	0	0.0	1	4.8	1	3.8	(2)	(4.3)	7	8.1	4	5.4	6	15.4	0	0.0	0	0.0	0	0.0
	HARDING	1	6.3	0	0.0	0	0.0	0	0.0	(0)	(0.0)	1	20.0	0	0.0	0	0.0	0	0.0	0	0.0	0	0.0
	LOS ALAMOS	5	2.5	0	0.0	0	0.0	1	16.7	(1)	(14.3)	1	2.2	2	2.7	1	2.0	0	0.0	0	0.0	0	0.0
	MORA	10	13.9	0	0.0	1	33.3	1	8.3	(2)	(13.3)	3	11.5	2	10.0	2	25.0	1	33.3	0	0.0	0	0.0
2	RIO ARRIBA	150	19.8	0	0.0	12	22.6	30	30.0	(42)	(27.5)	59	22.9	24	13.0	15	15.2	8	14.0	1	20.0	1	100.0
	SAN MIGUEL	70	15.0	0	0.0	8	34.8	11	29.7	(19)	(31.7)	25	15.4	20	14.0	6	8.6	0	0.0	0	0.0	0	0.0
	SANTA FE	263	18.2	1	50.0	20	26.3	41	36.3	(61)	(32.3)	84	20.4	50	12.7	41	13.9	24	18.0	2	8.7	0	0.0
	TAOS	54	11.8	0	0.0	6	27.3	2	5.1	(8)	(13.1)	16	11.9	15	12.3	12	11.8	2	6.9	0	0.0	1	100.0
	UNION	8	10.5	0	0.0	3	50.0	1	10.0	(4)	(25.0)	1	5.0	1	4.0	1	11.1	0	0.0	1	33.3	0	0.0
	BERNALILLO	886	11.2	3	23.1	63	17.9	122	17.1	(185)	(17.4)	340	13.5	213	8.9	113	7.9	30	6.4	2	4.2	0	0.0
	SANDOVAL	140	14.3	1	50.0	10	27.8	16	23.2	(26)	(24.8)	46	15.9	38	11.5	17	8.8	10	20.4	2	16.7	0	0.0
3	TORRANCE	22	16.2	0	0.0	3	37.5	5	38.5	(8)	(38.1)	9	20.0	2	4.5	1	5.9	2	28.6	0	0.0	0	0.0
	VALENCIA	87	13.8	0	0.0	9	25.7	12	26.1	(21)	(25.9)	31	14.7	21	10.8	8	8.5	5	12.2	1	16.7	0	0.0
	CHAVES	253	25.0	2	33.3	26	33.8	36	35.3	(62)	(34.6)	111	28.5	51	19.0	20	16.0	6	16.7	1	14.3	0	0.0
	CURRY	70	7.8	1	33.3	8	15.7	13	11.8	(21)	(13.0)	29	7.6	11	4.7	6	6.6	1	5.0	1	25.0	0	0.0
	DE BACA	9	25.0	0	0.0	2	66.7	0	0.0	(2)	(18.2)	3	27.3	3	27.3	0	0.0	0	0.0	1	100.0	0	0.0
	EDDY	152	17.2	1	100.0	12	20.0	31	27.9	(43)	(25.1)	59	19.0	29	10.8	10	11.1	7	18.9	3	42.9	0	0.0
4	GUADALUPE	14	17.9	0	0.0	1	14.3	7	58.3	(8)	(42.1)	3	10.7	1	6.3	1	11.1	1	20.0	0	0.0	0	0.0
	LEA	212	15.5	1	16.7	18	18.8	38	26.4	(56)	(23.3)	83	16.8	49	11.4	13	8.6	9	25.0	1	11.1	0	0.0
	LINCOLN	28	11.4	0	0.0	3	33.3	6	31.6	(9)	(32.1)	11	11.5	4	5.6	1	2.8	2	22.2	1	25.0	0	0.0
	QUAY	46	26.1	0	0.0	7	41.2	13	34.2	(20)	(26.4)	14	25.9	8	17.8	2	12.5	2	40.0	0	0.0	0	0.0
	ROOSEVELT	47	16.4	0	0.0	1	6.3	10	27.0	(11)	(20.8)	14	14.9	11	14.9	7	15.9	3	23.1	1	16.7	0	0.0
	CATRON	4	11.8	0	0.0	0	0.0	0	0.0	(0)	(0.0)	1	9.1	1	10.0	0	0.0	1	33.3	0	0.0	1	100.0
	DONA ANA	250	11.7	0	0.0	18	16.5	45	20.3	(63)	(19.0)	97	12.8	52	8.4	22	7.5	13	11.0	3	20.0	0	0.0
	GRANT	45	8.7	0	0.0	5	12.5	11	16.9	(16)	(15.2)	11	6.7	13	9.2	3	4.3	1	3.4	0	0.0	1	50.0
5	HIDALGO	11	10.3	0	0.0	1	9.1	2	20.0	(3)	(14.3)	4	9.8	3	9.7	1	11.1	0	0.0	0	0.0	0	0.0
	LUNA	68	22.5	0	0.0	7	24.1	13	31.7	(20)	(28.6)	22	22.2	7	9.6	13	34.2	5	27.8	1	25.0	0	0.0
	OTERO	102	9.2	1	50.0	7	18.4	18	15.3	(25)	(16.0)	40	8.4	23	7.4	7	5.4	6	16.7	0	0.0	0	0.0
	SIERRA	16	16.3	0	0.0	1	50.0	3	21.4	(4)	(25.0)	5	14.7	4	13.8	2	12.5	1	33.3	0	0.0	0	0.0
	SOCORRO	38	15.3	0	0.0	2	12.5	8	26.7	(10)	(21.7)	13	16.5	9	13.8	5	11.1	1	8.3	0	0.0	0	0.0

Percentage is based on births reporting little or no care as a portion of all births within the County/age category.
The number of births with low or no prenatal care may be signifcantly undercounted in counties within a large proportion of births reporting an "unknown" level of care.

Table 2.25 and Figure 2.21 give data on live births to mothers who reported receiving little or no prenatal care by race and ethnicity of the infant. Examination of this data shows that among Indians, mothers were nearly three times as likely to have received low levels of prenatal care than among non-Hispanic Whites, while among Hispanic Whites and Blacks, mothers were nearly twice as likely to have received low levels of care in 1985.

Nearly one-third of births to Hispanic Whites in Chaves, De Baca and Quay counties were to women who received low levels of prenatal care. In actual numbers Bernalillo and Dona Ana counties recorded among Hispanic Whites the highest number of births to women receiving low levels of prenatal care, while McKinley, San Juan and Sandoval counties were highest among Indians (Table 2.25).

For the purpose of this report, a high level of care was defined to be care that began in the first to third month of pregnancy with nine or more visits. Mid-levels of care were defined to be care that began in the first to third month with 5-8 visits, or care beginning in the fourth to sixth month with five or more visits. The various combinations are graphically depicted in Figure 2.22. The chart allows for the categorization of inconsistent entries such as "no visits with care beginning in a given month."

TABLE 2.25
NEW MEXICO
RESIDENT LIVE BIRTHS
PERCENT REPORTING LOW LEVEL PRENATAL CARE BY RACE/ETHNIC GROUP
1985

HEALTH DISTRICTS	COUNTY	TOTAL		WHITE Non-Hispanic		WHITE Hispanic		BLACK		INDIAN		OTHER	
		NO.	%	NO.	%	NO.	%	NO.	%	NO.	%	NO.	%
	NEW MEXICO	3851	14.0	946	9.2	1946	15.7	111	16.4	811	21.3	37	10.6
	CIBOLA	103	19.6	14	15.9	34	17.3	1	12.5	53	23.1	1	25.0
1	MCKINLEY	363	21.1	11	6.9	24	13.2	1	12.5	327	23.9	0	0.0
	SAN JUAN	305	13.7	51	6.0	37	10.5	3	16.7	213	21.5	1	11.1
	COLFAX	19	7.4	5	5.5	14	8.6	0	0.0	0	0.0	0	0.0
	HARDING	1	6.3	0	0.0	1	11.1	0	0.0	0	0.0	0	0.0
	LOS ALAMOS	5	2.5	3	1.9	2	5.1	0	0.0	0	0.0	0	0.0
	MORA	10	13.9	2	33.3	8	12.3	0	0.0	0	0.0	0	0.0
2	RIO ARRIBA	150	19.8	7	12.1	113	21.0	2	33.3	28	17.9	0	0.0
	SAN MIGUEL	70	15.0	5	10.9	64	15.4	0	0.0	1	33.3	0	0.0
	SANTA FE	263	18.2	68	15.5	183	20.0	3	25.0	5	7.2	4	26.7
	TAOS	54	11.8	8	9.1	35	10.9	0	0.0	10	22.2	1	33.3
	UNION	8	10.5	1	3.1	7	15.9	0	0.0	0	0.0	0	0.0
	BERNALILLO	886	11.2	297	8.9	468	12.4	40	15.3	61	16.1	20	12.1
	SANDOVAL	140	14.3	32	8.9	41	15.6	2	11.8	64	19.3	1	10.3
3	TORRANCE	22	16.2	4	6.9	16	21.9	0	0.0	2	50.0	0	0.0
	VALENCIA	87	13.8	22	10.3	56	14.8	0	0.0	8	25.8	1	25.0
	CHAVES	253	25.0	95	20.3	150	29.4	6	26.1	1	25.0	1	16.7
	CURRY	70	7.8	27	5.3	35	13.0	6	7.8	1	25.0	1	2.7
	DE BACA	9	25.0	4	21.1	5	29.4	0	0.0	0	0.0	0	0.0
	EDDY	152	17.2	64	14.5	85	19.9	3	27.3	0	0.0	0	0.0
4	GUADALUPE	14	17.9	1	11.1	13	20.0	0	0.0	0	0.0	0	0.0
	LEA	212	15.5	59	7.7	127	24.7	25	33.8	1	16.7	0	0.0
	LINCOLN	28	11.4	16	10.6	11	12.9	0	0.0	0	0.0	1	50.0
	QUAY	46	26.1	15	19.5	29	30.2	1	100.0	1	50.0	0	0.0
	ROOSEVELT	47	16.4	23	13.5	21	20.4	2	50.0	1	25.0	0	0.0
	CATRON	4	11.8	1	5.0	3	23.1	0	0.0	0	0.0	0	0.0
	DONA ANA	250	11.7	33	5.5	207	14.1	7	19.4	1	4.5	2	11.8
	GRANT	45	8.7	12	7.0	31	9.5	0	0.0	1	20.0	1	25.0
5	HIDALGO	11	10.3	7	16.3	4	6.3	0	0.0	0	0.0	0	0.0
	LUNA	68	22.5	12	15.4	53	24.5	3	42.9	0	0.0	0	0.0
	OTERO	102	9.2	35	5.7	41	14.9	6	6.3	19	21.6	1	2.3
	SIERRA	16	16.3	6	12.2	9	19.1	0	0.0	1	100.0	0	0.0
	SOCORRO	38	15.3	6	9.1	19	14.4	0	0.0	12	26.1	1	33.3

Categorization by child's race/ethnic group.
Percentage is based on births reporting little or no prenatal care as a portion of all births with the county/race/ethnic category.
The number of births with little or no prenatal care may be significantly undercounted in counties with a large proportion of births reporting an "unknown" level of care.

FIGURE 2.21
NEW MEXICO RESIDENT LIVE BIRTHS PERCENT REPORTING LOW LEVEL PRENATAL CARE BY RACE/ETHNIC GROUP
1984 AND 1985



Inconsistent entries on the birth certificate were a relatively minor problem; a greater problem was the number of certificates on which the information was not reported at all.

FIGURE 2.22
NEW MEXICO
CLASSIFICATION OF LEVELS OF PRENATAL CARE USING AS CRITERIA THE MONTH CARE BEGAN AND THE NUMBER OF PRENATAL VISITS



NUMBER OF PRENATAL VISITS	MONTH CARE BEGAN				
	1-3	4-6	7-9	NONE	UNKNOWN
9+	H	M	L	L	?
5-8	M	M	L	L	?
1-4	L	L	L	L	L
NONE	L	L	L	(NONE) L	L
UNKNOWN	?	?	L	L	?

HIGH (OPTIMUM) LEVEL
LOW LEVEL
UNKNOWN LEVEL ?
MID LEVEL
NONE (LOW LEVEL)

A breakdown of resident births by county and selected cities is presented in Table A-7 and births by county for the years 1970-1985 in Table A-2 of the Appendix.

A summarization of natality data characteristics for 1985 is presented in Table 2.26, on the following page.

TABLE 2.26
NATALITY CHARACTERISTICS BY COUNTY
NEW MEXICO RESIDENTS 1985

HEALTH DISTRICTS	COUNTY	LIVE BIRTHS		SEX	LOW BIRTH	% BIRTHS	BIRTHS TO	RACE OF CHILD				
			RATE	RATIO	WEIGHT	SINGLE	MOTHERS=<19	WHITE		NON-WHITE		
		NUMBER	PER 1,000 POP	M/F	%	MOTHERS	%	NON-HISPANIC	HISPANIC	BLACK	INDIAN	OTHER
	NEW MEXICO	27449	18.9	1.07	6.9	26.4	15.5	10250	12370	675	3805	349
	CIBOLA	525	20.8	1.00	6.3	36.4	18.9	88	196	8	229	4
1	MCKINLEY	1720	27.2	1.04	6.9	52.0	17.3	159	182	8	1370	1
	SAN JUAN	2228	24.5	1.06	5.8	28.6	14.6	855	353	18	993	9
	COLFAX	256	17.9	1.21	4.7	23.8	18.8	91	163	0	0	2
	HARDING	16	16.0	1.29	6.3	12.5	12.5	7	9	0	0	0
	LOS ALAMOS	204	11.1	1.17	6.4	5.9	3.4	159	39	0	3	3
	MORA	72	15.7	.89	15.3	37.5	20.8	6	65	0	1	0
2	RIO ARRIBA	758	23.4	1.09	9.4	36.5	20.3	58	538	6	156	0
	SAN MIGUEL	467	18.8	.96	10.1	31.3	13.3	46	416	1	3	1
	SANTA FE	1449	17.1	1.00	7.5	26.6	13.2	439	914	12	69	15
	TAOS	458	20.9	.94	9.2	26.2	13.5	88	320	2	45	3
	UNION	76	14.9	1.05	13.2	15.8	21.1	32	44	0	0	0
	BERNALILLO	7937	17.1	1.09	7.6	26.0	13.6	3348	3784	261	379	165
	SANDOVAL	981	22.2	1.17	6.1	28.2	10.9	360	263	17	331	10
3	TORRANCE	136	15.8	1.16	8.8	14.7	15.4	58	73	1	4	0
	VALENCIA	629	17.5	1.09	7.9	21.3	13.0	213	379	2	31	4
	CHAVES	1012	17.9	1.17	5.7	24.0	18.3	469	510	23	4	6
	CURRY	893	21.0	1.06	5.2	20.6	18.4	505	270	77	4	37
	DE BACA	36	15.0	2.00	5.6	16.7	30.6	19	17	0	0	0
	EDDY	885	16.9	.97	7.1	20.5	19.4	442	428	11	1	3
4	GUADALUPE	78	17.7	1.00	6.4	30.8	24.4	9	65	0	1	3
	LEA	1367	20.9	1.01	7.6	16.5	18.0	764	514	74	6	9
	LINCOLN	246	17.7	1.00	6.9	17.1	11.4	151	85	2	6	2
	QUAY	176	15.0	1.12	6.3	33.0	31.3	77	96	1	2	0
	ROOSEVELT	286	17.3	.97	6.3	17.8	19.2	171	103	4	4	4
	CATRON	34	12.1	.89	8.8	8.8	8.8	20	13	0	1	0
	DONA ANA	2140	18.1	1.07	5.7	23.8	15.7	596	1469	36	22	17
	GRANT	515	18.9	.97	6.4	24.1	20.6	172	328	6	5	4
5	HIDALGO	107	17.3	1.14	6.5	26.2	19.6	43	64	0	0	0
	LUNA	302	17.3	1.22	8.3	26.8	23.2	78	216	7	0	1
	OTERO	1114	22.6	1.15	4.4	15.3	14.2	612	275	96	88	43
	SIERRA	98	10.3	1.39	2.0	18.4	16.3	49	47	1	1	0
	SOCORRO	248	18.2	.89	5.6	20.6	18.5	66	132	1	46	3

MORTALITY

ALL CAUSES

The trend in crude death rates (number dying during the year expressed per 1,000 population) for New Mexico and the United States is shown in Figure 3.1. The lower death rates for New Mexico are the result of a younger population rather than a more favorable mortality. As noted earlier in Figure 1.4, only 9.6% of New Mexico's population is age 65 or older, compared with 12.0% for the U.S.

FIGURE 3.1
CRUDE DEATH RATE
NEW MEXICO RESIDENTS AND UNITED STATES
1960-1985



To control the effect created upon the rates by a difference in age composition, age-adjusted methods are employed to yield a death rate which would be expected if New Mexico had the same age structure population as that of the United States.

In Table 3.1, crude death rates by sex for New Mexico and the various race/ethnic populations are compared with national data. The apparent lower death rates for New Mexico are deceiving.

The favorable comparison caused by the State's young population structure disappears upon age-adjusting the rates. Age adjusted mortality rates for New Mexico were slightly lower than national rates for males and females; not as great as was indicated by the crude rates. Examination of age-adjusted mortality rates for the State's various race/ethnic populations show the highest male mortality in the State as well as the highest female mortality to be for Blacks. Mortality rates for Indian and Black males in the State exceed that for males nationally; for females the national rate was exceeded by the rate for New Mexico Black women.

Comparison of adjusted mortality rates in the State for the periods 1980-1982, 1981-1983, and 1983-1985 show an overall mortality decrease for both males and females.

TABLE 3.1
NEW MEXICO CRUDE AND ADJUSTED DEATH RATES NEW MEXICO RESIDENTS 1983-85, 1981-83, 1980-82 AVERAGES
Compared with the United States 1984

RATE PER 100,000 POPULATION	1983-85	1983-85	1981-83	1980-82
	CRUDE DEATH RATE	NEW MEXICO AGE-ADJUSTED DEATH RATE	NEW MEXICO AGE-ADJUSTED DEATH RATE	NEW MEXICO AGE-ADJUSTED DEATH RATE
UNITED STATES				
MALE	935.9	XXX	XXX	XXX
FEMALE	792.7	XXX	XXX	XXX
NEW MEXICO ALL RACES:				
MALE	752.9	828.5	890.5	915.3
FEMALE	552.7	693.7	747.7	737.0
NEW MEXICO WHITE NON-HISPANIC:				
MALE	874.3	824.0	893.0	910.9
FEMALE	684.5	687.5	753.6	742.9
NEW MEXICO WHITE HISPANIC:				
MALE	607.5	785.0	850.4	878.5
FEMALE	421.8	707.3	737.2	722.0
NEW MEXICO BLACK:				
MALE	700.7	1033.6	974.2	985.3
FEMALE	477.1	815.4	778.6	780.9
NEW MEXICO INDIAN:				
MALE	703.3	1020.1	1021.3	1097.4
FEMALE	354.1	673.6	750.3	757.2

In comparing mortality rates by age and sex for New Mexico and the United States (Table 3.2), variations are quite apparent. Interestingly, New Mexico's mortality is higher than nationally for age groups under 45 and lower for age groups 45 and older with the exception for males in the 35-44 age bracket for whom the mortality rate in New Mexico is 2% lower than nationally.

TABLE 3.2
AGE SPECIFIC DEATH RATES NEW MEXICO RESIDENTS 3 YEAR AVERAGE 1983-85 AND UNITED STATES 1984

	MALE		FEMALE	
AGE GROUPS	NEW MEXICO	UNITED STATES	NEW MEXICO	UNITED STATES
1 - 4	73.8	57.0	54.2	46.5
5 -14	41.1	32.1	26.1	21.1
15 -24	216.9	141.6	62.8	51.1
25 -34	252.8	173.9	82.3	68.5
35 -44	265.7	270.2	143.2	141.5
45 -54	616.0	676.6	321.6	374.6
55 -64	1378.4	1705.2	778.0	918.4
65 -74	3095.8	3813.0	1738.6	2096.4
75 -84	7134.0	8445.9	4221.3	5188.2
85+	15117.3	18119.5	12962.7	14053.9

RATES PER 100,000 POPULATION IN SPECIFIED GROUP.

Age specific rates by county and sex are presented in Table A-8 of the Appendix.

LIFE EXPECTANCY

Life expectancy for New Mexico males is the same (70.3 years) as for the U.S. male population. However, for the State's female population, life expectancy is 1.1 years more than for the U.S. female population.

For New Mexico males, life expectancy is highest for the non-Hispanic White, while for females, it is highest for Hispanic White. (Figure 3.2)

New Mexico life expectancy data is based on a 3-year (1979-1981) average mortality and race/ethnic population based on 1980 Census. (Table 3.3)

FIGURE 3.2
AVERAGE LIFETIME IN YEARS NEW MEXICO 1979-1981 UNITED STATES 1981



LEADING CAUSES OF DEATH

The leading causes of death in 1985 in New Mexico and the United States are ranked in Table 3.4.

TABLE 3.3
AVERAGE YEARS OF LIFE REMAINING NEW MEXICO RESIDENTS BASED ON THREE YEAR AVERAGE (1979-81) MORTALITY RATES

Sex/Race	All Races	Non-Hispanic White	Hispanic White	Black	Indian
Male:					
0 - 1	70.3	71.0	70.6	68.2	64.2
1 - 4	70.2	70.9	70.5	68.8	64.2
5 - 14	66.4	67.1	66.7	65.0	60.6
15 - 24	56.7	57.1	57.0	55.6	51.0
25 - 34	48.2	48.6	48.4	46.5	43.9
35 - 44	39.4	39.5	39.8	37.5	36.5
45 - 54	30.6	30.4	31.2	28.7	29.8
55 - 64	22.5	22.1	23.2	21.5	23.1
65 - 74	15.4	15.0	16.1	15.6	17.1
75 - 84	9.7	9.3	10.3	10.8	11.0
85 +	5.6	5.3	5.9	6.0	6.1
Female:					
0 - 1	79.0	79.0	79.5	77.4	77.7
1 - 4	78.8	78.8	79.2	77.6	77.7
5 - 14	75.0	75.0	75.4	74.0	74.1
15 - 24	65.3	65.2	65.6	64.2	64.5
25 - 34	55.7	55.6	55.9	54.6	55.1
35 - 44	46.1	46.0	46.4	45.2	46.3
45 - 54	36.8	36.7	37.0	35.9	37.6
55 - 64	28.0	27.9	27.9	27.4	29.7
65 - 74	19.9	19.8	19.8	20.2	22.3
75 - 84	12.8	12.6	12.8	14.4	14.9
85 +	7.4	7.1	7.6	10.3	10.9

TABLE 3.4
NEW MEXICO RESIDENTS, 1985 UNITED STATES 1985, PROVISIONAL DEATHS RATES FROM LEADING CAUSES

NEW MEXICO				UNITED STATES		
CRUDE DEATH RATE*	PERCENT OF TOTAL DEATHS	RANK	CAUSE OF DEATH	CRUDE DEATH RATE*	PERCENT OF TOTAL DEATHS	RANK
664.6	100.0%	xxx	ALL CAUSES	874.8	100.0%	xxx
186.2	28.0	1	Diseases of the Heart	325.0	37.2	1
132.5	19.9	2	Malignant Neoplasms	191.7	22.0	2
39.7	6.0	4	Cerebrovascular Disease	64.0	7.3	3
59.4	8.9	3	Accidents	38.6	4.4	4
33.3	5.0	5	Chronic Obstructive Pulmonary Disease and Allied Conditions	31.2	3.6	5
18.6	2.8	6	Influenza & Pneumonia	27.9	3.2	6
17.3	2.6	8	Diabetes Mellitus	16.2	1.9	7
17.7	2.7	7	Suicide	12.0	1.4	8
13.7	2.1	9	Chronic Liver Disease and Cirrhosis	11.2	1.3	9
6.1	0.9	14	Atherosclerosis	9.9	1.1	10
7.3	1.1	13	Nephritis, Nephrotic Syndrome and Nephrosis	9.4	1.1	11
11.0	1.7	10	Homicide	8.1	0.9	12
8.4	1.3	11	Certain Conditions Originating in the Perinatal Period	7.6	0.9	13
3.1	0.5	16	Septicemia	7.1	0.8	14
7.7	1.2	12	Congenital Anomalies	5.5	0.6	15
5.4	0.8	15	Alcoholism**	1.7	0.2	xxx

---

*Rate per 100,000 population

**Includes only deaths coded 303, "Alcohol dependence syndrome", according to the Ninth Revision International Classification of Diseases. Deaths due to alcoholic cirrhosis are included in the category "Chronic Liver Disease and Cirrhosis." U.S. figure for category 303 based on 1984 data.

A detailed listing of causes of death in New Mexico is found in Table A-9 of the Appendix. The percentage of deaths in 1985 for leading causes are displayed graphically in Figure 3.3.

FIGURE 3.3
LEADING CAUSES OF DEATH
NEW MEXICO RESIDENTS 1985
UNITED STATES 1985



Diseases of the heart and malignant neoplasms rank one and two for both New Mexico and the United States, although these two causes account for lower percentages of deaths in New Mexico than nationally.

Nationally, stroke ranks as the third major cause of death, but in New Mexico, accidents rank third, accounting for twice the percentage of deaths than for the nation as a whole.

New Mexico's fatality rate for accidental deaths by place of occurrence ranks third highest among the states, as the State ranks highest in motor vehicle accident fatalities per 100,000 population. If, the motor vehicle fatality rate is taken with regard to the number of deaths per vehicle miles, New Mexico ranks first among the states in 1985, and third in traffic deaths per registered automobiles.

FIGURE 3.4
PRINCIPAL CLASSES OF ACCIDENTAL DEATHS
NEW MEXICO AND UNITED STATES
1985
(Death Rate Per 100,000 Population)



SOURCE: Accident Facts 1986 Edition
National Safety Council

As noted in Table 3.5, the State's crude death rate fluctuated over the last few years with the result that the State's crude death rate in 1985 (664.6 deaths per 100,000 population) was 1% lower than the 1982 rate (672.0). Increases in rates were noted for the leading causes in 1985 over the prior year, with the following exceptions: malignant neoplasms decreased 5%, accidents decreased 4%, cerebrovascular disease decreased 2%, influenza and pneumonia decreased 8%, certain causes of mortality in early infancy decreased 10%, atherosclerosis decreased 2% and septicemia decreased 29%. Heart disease, which is the leading cause of death, decreased by a tenth of a percent.

The ranking for the top 11 leading causes remained the same as for the previous year.

The percentage of deaths for leading causes showed marked variation among New Mexico's various ethnic/racial groups Table 3.6.

Accidents are by far the leading cause of death among Indians. Alcoholism, suicide and cirrhosis were also notably high, while the percentage of deaths due to diseases of the heart and neoplasms were comparatively low. The percentage of homicides is high among Indians and Blacks. The Hispanic showed the second highest percentage of deaths due to cirrhosis of the liver. The non-Hispanic White showed

TABLE 3.5
NEW MEXICO RESIDENTS DEATH RATES FROM LEADING CAUSES
1982 TO 1985

CAUSE OF DEATH	1985 CRUDE DEATH RATE	RANK	1984 CRUDE DEATH RATE	RANK	1983 CRUDE DEATH RATE	RANK	1982 CRUDE DEATH RATE	RANK
ALL CAUSES	664.6	XX	666.5	XX	653.2	XX	672.0	XX
DISEASES OF HEART	186.2	1	186.5	1	174.9	1	160.7	1
MALIGNANT NEOPLASMS	132.5	2	139.1	2	136.5	2	142.2	2
ACCIDENTS	59.4	3	61.6	3	59.8	3	62.0	3
CEREBROVASCULAR DISEASE	39.7	4	40.3	4	39.2	4	39.1	4
CHRONIC OBSTRUCTIVE PULMONARY DISEASE AND ALLIED CONDITIONS	33.3	5	31.3	5	29.6	5	33.0	5
INFLUENZA AND PNEUMONIA	18.6	6	20.3	6	17.2	8	16.8	8
SUICIDE	17.7	7	17.5	7	18.2	7	19.7	7
DIABETES MELLITUS	17.3	8	15.8	8	20.5	6	23.5	6
CHRONIC LIVER DISEASE AND CIRRHOSIS	13.7	9	11.7	9	12.9	9	11.6	9
HOMICIDE	11.0	10	9.5	10	7.7	10	9.9	10
CERTAIN CAUSES OF MORTALITY IN EARLY INFANCY	8.4	11	9.3	11	7.4	11	8.5	12
CONGENITAL ANOMALIES	7.7	12	5.5	14	6.6	13	6.6	14
NEPHRITIS, NEPHROTIS SYNDROME AND NEPHROSIS	7.3	13	7.2	12	3.6	15	5.3	15
ATHEROSCLEROSIS	6.1	14	6.2	13	6.9	12	6.9	13
ALCOHOLISM *	5.4	15	5.3	15	5.1	14	9.9	10
SEPTICEMIA	3.1	16	4.4	16	1.9	16	2.7	16

* Includes only deaths coded 303, "Alcohol Dependence Syndrome", according to the Ninth Revision International Classification of Diseases. Deaths due to alcoholic cirrhosis are included in the category "Chronic Liver Disease and Cirrhosis".

CRUDE DEATH RATES PER 100,000 POPULATION.

TABLE 3.6
NEW MEXICO RESIDENT DEATHS PERCENTAGE DISTRIBUTION OF DEATHS BY CAUSE, ETHNIC/RACE
1983-1985 AGGREGATE

CAUSE	PERCENTAGE DISTRIBUTION FOR EACH ETHNIC/RACE				
	ALL RACES	White		BLACK	INDIAN
		Non-Hispanic	Hispanic		
ALL CAUSES	100.0%	100.0%	100.0%	100.0%	100.0%
Diseases of the Heart	27.6	30.2	25.4	27.9	12.1
Malignant Neoplasms	20.6	22.7	18.0	20.3	11.2
Cerebrovascular Disease	6.0	6.4	5.8	8.0	2.8
Accidents	9.1	6.9	10.5	6.9	24.9
Chronic Obstructive Pulmonary Diseases and Allied Conditions	4.7	6.0	3.0	2.3	1.0
Influenza & Pneumonia	2.8	2.8	2.8	1.7	3.4
Diabetes Mellitus	2.7	2.2	3.5	4.8	3.4
Suicide	2.7	2.6	2.7	0.8	4.1
Chronic Liver Disease and Cirrhosis	1.9	1.3	2.9	2.3	4.1
Atherosclerosis	1.0	1.1	0.7	0.6	0.4
Nephritis, Nephrotic Syndrome and Nephrosis	0.9	0.9	1.0	1.5	0.9
Homicide	1.4	0.7	2.5	3.1	3.4
Certain Conditions Originating in the Perinatal Period	1.3	0.8	2.0	2.7	2.4
Septicemia	0.5	0.5	0.5	1.0	0.7
Congenital Anomalies	1.0	0.6	1.6	0.6	1.8
Alcoholism*	0.8	0.5	1.0	0.2	2.9
All Other	15.0	13.9	16.1	15.1	20.6

the highest percentage of deaths due to heart disease and malignant neoplasms.

As already noted, diseases of the heart, malignant neoplasms, accidents and stroke ranked as the four leading causes of death in New Mexico. New Mexico's crude death rate for diseases of the heart, malignant neoplasms and stroke were much lower than corresponding national rates.

The reader is reminded of the "bias" produced by the "youngness" of New Mexico's population. To obtain more accurate comparisons of New Mexico's mortality rates to national figures, New Mexico's rates must be age-adjusted in order to remove the difference caused by population age structure.

Table 3.7 on the following page presents the leading causes of death ranked by crude death rates for the United States and New Mexico. Also presented are New Mexico age-adjusted rates. The reader will note the difference made by the age-adjustment of the rates. New Mexico data are based on a three year, 1983-85, average.

For example, comparison of crude death rates for influenza/pneumonia shows New Mexico's rate 26% lower than the national rate, but upon age-adjustment of New Mexico's rate, the State's mortality rate from influenza/pneumonia is only 7% lower than the national rate.

Heart disease and cancer ranked first and second as the leading causes of death for all of New Mexico's race/ethnic groups except for the Indian for whom accidents were the leading cause of death. Accidents ranked third for the State's other race/ethnic populations, with the exception for the Black for whom cerebrovascular disease ranked third and accidents fourth. The percentage of accidental deaths involving motor vehicles were generally higher for the State's various race/ethnic populations than nationally. Suicide rates were notably high for the White and Indian. Homicide rates for the Hispanic White, Black and Indian were also high. Alcoholism and cirrhosis of the liver rates were notably high for some of New Mexico's population.

Mortality due to malignant neoplasms was highest among Black males and non-Hispanic Whites for females, exceeding the corresponding national rates in the case of males. COPD mortality rates were substantially higher for New Mexicans of both sexes. Diabetes mellitus rates were particularly high for the Hispanic, Black and Indian populations.

Variations are pronounced and more detailed comparisons are left to the interest of the reader (Table 3.7).

Trends in death rates for the seven leading causes of death over a 25 year span are presented in Figure 3.5.

The death rate from diseases of the heart peaked in the early Seventies and had since shown a general decline up through 1981 when the mortality rate was lower than it had been in 1960; however the rate has increased notably since 1981.

The State's mortality rate from cancer has shown a general decrease since 1982.

Since peaking in the late Sixties and early Seventies accident mortality in New Mexico has dropped markedly, leveling off somewhat the last few years—dropping 4% in 1985 compared to 1984. Non-motor vehicle accident mortality decreased and motor vehicle mortality increased over the previous year.

Stroke mortality has dropped dramatically since the mid Seventies, leveling off somewhat the last three years.

Influenza/pneumonia mortality fluctuated showing an 8% decrease in 1985.

Suicide mortality in New Mexico in 1985 increased 1% from the 1984 rate, however suicide mortality in the State has increased 58% since 1960 and presently ranks as the 7th leading cause of death in the State.

TABLE 3.7
NEW MEXICO RESIDENTS
CRUDE AND AGE-ADJUSTED DEATH RATES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

RATES PER 100,000 POPULATION	CRUDE DEATH RATE	NEW MEXICO ADJUSTED DEATH RATE		
		TOTAL	MALE	FEMALE
UNITED STATES:				
All Causes	862.3	XXX	935.9	792.7
Heart Disease	323.5	XXX	345.2	303.0
Malignant Neoplases	191.8	XXX	211.1	173.5
Cerebrovascular	65.3	XXX	53.6	76.3
Accidents	39.3	XXX	55.7	23.8
(% Motor Vehicle)	(49.9)	XXX	(51.3)	(46.2)
COPD	29.2	XXX	38.3	20.7
Influenza & Pneumonia	24.9	XXX	25.6	24.2
Diabetes Mellitus	15.1	XXX	12.9	17.2
Suicide	12.4	XXX	19.7	5.4
Cirrhosis of Liver	11.6	XXX	5.3	8.0
Atherosclerosis	10.3	XXX	8.0	12.5
Nephritis, Nephrotic Syndrome and Nephrosis	8.5	XXX	8.7	8.3
Homicide	8.4	XXX	13.1	3.9
NEW MEXICO-ALL RACES:				
All Causes	651.1	766.2	828.5	693.2
Heart Disease	179.7	220.1	236.6	199.9
Malignant Neoplases	133.8	157.2	162.8	148.5
Accidents	59.4	63.2	90.9	36.0
(% Motor Vehicle)	(56.1)	(53.8)	(56.3)	(48.6)
Cerebrovascular	39.0	48.8	37.4	60.1
COPD	30.9	37.5	44.3	29.2
Influenza & Pneumonia	18.4	23.2	21.2	24.3
Diabetes Mellitus	17.6	21.0	17.2	24.7
Suicide	17.6	18.5	29.9	7.4
Cirrhosis of Liver	12.6	13.9	19.7	8.2
Homicide	9.3	9.5	15.2	4.1
Certain Causes of Mortality in Early Infancy*	8.2	6.7	8.4	5.1
Congenital Anomalies	6.5	5.6	5.9	5.2
NON-HISPANIC WHITE:				
All Causes	777.7	757.2	824.0	687.5
Heart Disease	234.8	228.7	251.4	204.8
Malignant Neoplases	176.2	165.6	178.7	151.6
Accidents	53.7	53.9	73.6	34.4
(% Motor Vehicle)	(48.4)	(48.8)	(51.0)	(44.8)
Cerebrovascular	49.7	49.5	37.8	60.9
COPD	46.7	44.4	55.8	32.9
Influenza & Pneumonia	21.9	22.3	20.5	23.8
Suicide	20.0	19.4	30.3	8.7
Diabetes	17.3	16.6	14.1	19.0
Cirrhosis of Liver	9.9	9.0	12.2	5.9
Atherposclerosis	8.7	8.9	6.2	11.5
Nephritis, Nephrotic Syndrome and Nephrosis	6.7	6.6	7.8	5.2
Homicide	5.6	5.6	8.1	3.2
HISPANIC WHITE:				
All Causes	513.4	754.4	785.0	707.3
Heart Disease	130.6	212.8	219.5	198.0
Malignant Neoplases	92.4	143.6	133.0	149.6
Accidents	54.1	62.0	89.7	34.9
(% Motor Vehicle)	(62.3)	(58.1)	(62.0)	(49.0)
Cerebrovascular	29.6	48.6	36.3	61.4
Diabetes Mellitus	17.8	28.5	23.3	33.9
COPD	15.3	25.4	24.8	24.2
Cirrhosis of Liver	14.8	20.6	31.1	10.2
Influenza & Pneumonia	14.6	23.6	19.9	26.4
Suicide	14.1	14.8	24.7	5.4
Homicide	12.7	13.5	23.1	4.3
Certain Causes of Mortality in Early Infancy*	10.4	6.8	8.4	5.3
Congenital Anomalies	8.1	6.1	6.1	6.0
BLACK:				
All Causes	593.4	934.0	1033.6	815.4
Heart Disease	165.5	283.1	292.2	268.0
Malignant Neoplases	120.7	201.2	254.7	134.6
Cerebrovascular	47.3	84.1	58.2	114.6
Accidents	41.1	47.6	81.1	12.2
(% Motor Vehicle)	(54.5)	(50.2)	(49.9)	(55.7)
Diabetes Mellitus	28.6	46.2	21.5	73.5
Homicide	18.7	20.6	36.3	3.6
Certain Causes of Mortality in Early Infancy*	16.2	10.1	16.1	4.4
COPD	13.7	22.2	29.2	12.2
Cirrhosis of Liver	13.7	21.7	30.5	13.5
Influenza & Pneumonia	10.0	16.3	22.3	10.8
Nephritis, Nephrotic Syndrome and Nephrosis	8.7	16.7	4.1	30.6
Septicemia	6.2	8.5	9.8	7.2
INDIAN:				
All Causes	522.0	844.9	1020.1	673.6
Accidents	130.1	157.7	259.2	64.4
(% Motor Vehicle)	(65.0)	(60.0)	(58.3)	(68.0)
Heart Disease	63.1	130.9	141.2	116.8
Malignant Neoplases	58.5	114.1	111.4	112.8
Cirrhosis of Liver	21.5	31.6	41.2	23.3
Suicide	21.2	23.1	45.5	3.9
Diabetes	17.8	37.0	26.0	49.1
Influenza & Pneumonia	17.5	33.2	36.5	26.5
Homicide	17.5	19.1	28.8	11.0
Alcoholism	15.0	19.9	35.8	6.1
Cerebrovascular	14.4	31.1	36.6	25.4
Certain Causes of Mortality of Early Infancy*	12.7	6.2	6.9	5.5
Congenital Anomalies	9.6	5.0	5.3	4.8

*Certain Causes of Perinatal Mortality ICDA 760-769.4-772, 774-778

COPD - Chronic Obstructive Pulmonary Disease

FIGURE 3.5
NEW MEXICO RESIDENT DEATHS
CRUDE DEATH RATES PER 100,000 POPULATION
SELECTED CAUSES
1960-1985



* *C.O.P.D. – CHRONIC OBSTRUCTIVE PULMONARY DISEASE*
Reporting under this category began in 1979

LEADING CAUSES BY AGE

Nearly twice the percentage of deaths in New Mexico occurred to persons under 35 years of age than did nationally. Of the deaths occurring nationally, 7.0% were persons under 35 years. In New Mexico, 12.6% were under age 35.

Among the Indian population, the percentage of deaths for males under 35 was 38.7%, and for females, 25.7%, the highest for this age group as compared to the State's other ethnic populations.

Nationally, 77.5% of female deaths were 65 or older. In New Mexico, the highest percentage of deaths for women 65 or older were for non-Hispanic White females at 75.6%. Only 44.0% of the deaths for Indian women occurred at age 65 or older.

TABLE 3.8
PERCENTAGE OF DEATHS BY AGE GROUPS SEX AND RACE/ETHNIC GROUPS NEW MEXICO RESIDENTS 1983-1985 UNITED STATES 1985

SEX/RACE	TOTAL	PERCENTAGE DISTRIBUTION BY AGE GROUPS				
		< 1	1-14	15-34	35-64	65+
NEW MEXICO						
TOTAL	28,322	2.9%	1.6%	8.1%	23.9%	63.5%
Male						
All Races	16,095	3.1	1.7	10.9	26.3	58.0
Non-Hispanic White	9,880	2.2	1.1	6.7	25.7	64.3
Hispanic White	4,694	4.3	2.3	15.1	26.4	52.0
Black	293	5.8	2.7	9.2	34.5	47.8
Indian	1,196	5.5	4.3	28.9	28.6	32.7
Female						
All Races	12,227	2.7	1.5	4.4	20.7	70.7
Non-Hispanic White	8,023	1.5	1.0	2.7	19.2	75.6
Hispanic White	3,343	4.4	2.1	6.7	22.0	64.8
Black	184	3.3	1.6	3.8	28.8	62.5
Indian	650	8.0	4.5	13.2	30.3	44.0
UNITED STATES						
TOTAL	XXX	1.9	0.8	4.3	22.6	70.5
Male						
Total	XXX	2.1	0.9	6.0	26.9	64.1
White	XXX	1.7	0.8	5.3	25.5	66.6
Other	XXX	4.3	1.6	10.4	36.1	48.4
Female						
Total	XXX	1.7	0.7	2.3	17.8	77.5
White	XXX	1.4	0.6	2.0	16.2	79.8
Other	XXX	4.2	1.6	4.9	29.0	60.3

One factor contributing to these high percentages of younger deaths in New Mexico is of course the State's young population structure. Also, the high rate of mortality in the State for causes such as accidents, normally associated with younger age groups, would lead one to suspect higher rates of mortality within these age groups. Rates of mortality by cause of death are presented by age group in Tables 3.9 to 3.16. Comparison of mortality rates per 100,000 population in the age groups will provide insight into levels of mortality for the various age groups in the State.

AGES LESS THAN ONE

Mortality for infants less than one year of age is lower for the State than for the Nation for New Mexico White and Indian, but higher for Black. The death rate is higher for males than for females. Causes are similar although the actual rates differ in comparing State and national data. Accidental mortality is higher in New Mexico, with a much higher percentage involving motor vehicles. Further analysis of infant deaths per live births is presented in the section on infant mortality.

TABLE 3.9
AGES UNDER 1 YEAR
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

NEW MEXICO ETHNIC/RACE	RATE/100,000 POPULATION TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	1085.6	1198.2	967.5
Certain Causes of Mortality in Early Infancy*	512.4	565.2	457.0
Congenital Anomalies	234.4	245.2	223.1
Sudden Infant Death Syndrome (SIDS)	144.2	170.7	116.6
Heart Diseases	26.1	28.1	24.1
All Accidents	23.0	24.1	21.9
(% Motor Vehicle)	(19.1)	(17.4)	(21.0)
Influenza & Pneumonia	18.8	21.4	16.0
Septicemia	8.1	9.2	7.0
Nephritis, Nephrotic Syndrome and Nephrosis	7.5	8.7	6.2
Meningitis	7.4	8.3	6.3
NEW MEXICO-ALL RACES			
All Causes:	1007.3	1191.0	815.6
Certain Causes of Mortality in Early Infancy*	422.2	500.2	340.7
Congenital Anomalies	247.0	259.6	233.7
Sudden Infant Death Syndrome (SIDS)	148.4	219.1	74.6
All Accidents	26.8	31.0	22.4
(% Motor Vehicle)	(40.7)	(38.4)	(44.2)
Influenza & Pneumonia	24.3	31.0	17.4
Heart Disease	14.6	21.4	7.5
Malignant Neoplases	9.7	9.5	9.9
Homicide	7.3	2.4	12.4
WHITE-Non-Hispanic			
All Causes:	1078.6	1342.9	796.7
Certain Causes of Mortality in Early Infancy*	427.6	520.8	328.0
Congenital Anomalies	272.1	307.5	234.3
Sudden Infant Death Syndrome (SIDS)	197.6	301.2	87.0
All Accidents	32.4	37.7	26.8
(% Motor Vehicle)	(29.9)	(33.4)	(25.0)
Influenza & Pneumonia	25.9	31.4	20.1
Malignant Neoplases	13.0	12.6	13.4
Heart Disease	9.7	12.6	6.7
Nephritis, Nephrotic Syndrome and Nephrosis	6.5	6.3	6.7
Homicide	6.5	--	13.4
Septicemia	6.5	6.3	6.7
WHITE-Hispanic			
All Causes:	934.5	1058.0	806.3
Certain Causes of Mortality in Early Infancy*	428.2	497.3	356.5
Congenital Anomalies	234.3	232.8	235.9
Sudden Infant Death Syndrome (SIDS)	118.5	164.0	71.3
All Accidents	16.2	15.9	16.5
(% Motor Vehicle)	(33.3)	(33.3)	(33.3)
Influenza & Pneumonia	16.2	21.2	11.0
Heart Disease	16.2	26.5	5.5
Malignant Neoplases	10.8	10.6	11.0
BLACK			
All Causes:	1168.7	1696.6	621.1
Certain Causes of Mortality in Early Infancy*	609.8	998.0	207.0
Sudden Infant Death Syndrome (SIDS)	254.1	299.4	207.0
Congenital Anomalies	101.6	199.6	--
Homicide	50.8	--	103.5
Septicemia	50.8	99.8	--
INDIAN			
All Causes:	1054.8	1171.5	936.4
Certain Causes of Mortality in Early Infancy*	384.4	390.5	378.2
Congenital Anomalies	232.4	230.7	234.1
Sudden Infant Death Syndrome (SIDS)	107.3	177.5	36.0
Influenza & Pneumonia	53.6	71.0	36.0
All Accidents	44.7	53.2	36.0
(% Motor Vehicle)	(60.0)	(33.3)	(100.0)
Heart Disease	26.8	35.5	18.0
Homicide	17.9	17.7	18.0
Cerebrovascular	8.9	--	18.0

---

*Certain Causes originating in the Perinatal Period, ICDA 760-779

AGES 1 TO 4

Accidents are the leading cause of death for this age group. Rates for New Mexicans were higher than nationally with a larger percentage involving motor vehicles. Especially high were the mortality rates for Blacks and Indians. Mortality rates in this age group were higher for males.

TABLE 3.10
AGES 1-4
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

NEW MEXICO ETHNIC/RACE	RATE/100,000 POPULATION TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	51.9	57.0	46.5
All Accidents	19.8	23.9	15.5
(% Motor Vehicle)	(34.8)	(33.5)	(36.8)
Congenital Anomalies	6.7	6.7	6.6
Malignant Neoplasms	4.0	4.3	3.6
Heart Diseases	2.4	2.5	2.3
Homicide	2.4	2.4	2.4
Influenza & Pneumonia	1.5	1.4	1.5
Meningitis	1.2	1.4	1.0
NEW MEXICO-ALL RACES			
All Causes:	64.2	73.8	54.2
All Accidents	29.0	36.0	21.7
(% Motor Vehicle)	(49.0)	(41.9)	(60.8)
Congenital Anomalies	8.9	7.6	10.2
Homicide	3.3	5.8	0.6
Influenza & Pneumonia	3.0	2.9	3.0
Malignant Neoplasms	2.7	2.9	2.4
Heart Disease	2.7	2.3	3.0
WHITE-Non-Hispanic			
All Causes:	53.5	60.3	46.4
All Accidents	24.1	29.4	18.5
(% Motor Vehicle)	(53.1)	(35.0)	(83.8)
Congenital Anomalies	6.8	5.9	7.7
Malignant Neoplasms	4.5	5.9	3.1
Certain Causes of Mortality in Early Infancy*	2.3	4.4	--
Influenza & Pneumonia	1.5	1.5	1.5
Heart Disease	0.8	--	--
Homicide	0.8	1.5	--
Nephritis, Nephrotic Syndrome and Nephrosis	0.8	--	1.5
WHITE-Hispanic			
All Causes:	52.9	58.1	47.5
All Accidents	18.7	26.4	10.9
(% Motor Vehicle)	(39.6)	(40.2)	(37.6)
Congenital Anomalies	10.7	9.2	12.2
Influenza & Pneumonia	4.0	2.6	5.4
Homicide	4.0	6.6	1.4
Malignant Neoplasms	2.0	1.3	2.7
Heart Disease	2.0	1.3	2.7
Cerebrovascular	1.3	1.3	1.4
Certain Causes of Mortality in Early Infancy*	1.3	2.6	--
BLACK			
All Causes:	84.8	141.3	24.5
All Accidents	36.3	70.7	--
(% Motor Vehicle)	(66.7)	(66.6)	--
Heart Disease	12.1	23.6	--
Diabetes	12.1	23.6	--
Homicide	12.1	23.6	--
Certain Causes of Mortality in Early Infancy*	12.1	--	24.5
INDIAN			
All Causes:	137.4	164.2	110.3
All Accidents	80.2	86.7	73.6
(% Motor Vehicle)	(51.4)	(47.4)	(56.3)
Congenital Anomalies	11.5	9.1	13.8
Heart Disease	9.2	9.1	9.2
Homicide	6.9	13.7	--
Certain Causes of Mortality in Early Infancy*	4.6	9.1	--
Influenza & Pneumonia	4.6	9.1	--

COPD - Chronic Obstructive Pulmonary Disease
*SIDS (Sudden Infant Death Syndrome) death rates not ranked for this age group.
** Certain Causes originating in the Perinatal Period, ICDA 760-779

AGES 5 TO 14

As for the previous age group, accidents are the leading cause of death for this age group. The accidental mortality rate and overall mortality rates in the State were highest for White non-Hispanics for both females and males. Motor vehicles were involved in a greater percentage of New Mexico accident fatalities than nationally. Overall State mortality rates were higher for males than females in this age group.

TABLE 3.11
AGES 5-14
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

NEW MEXICO ETHNIC/RACE	RATE/100,000 POPULATION TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	26.7	32.1	21.1
All Accidents	12.4	16.5	8.0
(% Motor Vehicle)	(54.0)	(50.3)	(61.3)
Malignant Neoplasms	3.6	4.0	3.0
Congenital Anomalies	1.4	1.5	1.4
Homicide	1.3	1.3	1.3
Heart Disease	1.0	1.0	1.0
Suicide	0.7	1.0	0.3
Influenza & Pneumonia	0.5	0.5	0.5
NEW MEXICO-ALL RACES			
All Causes:	33.7	41.1	26.1
All Accidents	16.9	22.1	11.6
(% Motor Vehicle)	(66.9)	(61.5)	(77.6)
Malignant Neoplasms	5.0	5.7	4.3
Suicide	1.6	1.7	1.4
Homicide	1.6	1.4	1.7
Congenital Anomalies	1.3	1.4	1.2
Heart Disease	1.0	1.1	0.9
Influenza & Pneumonia	0.7	0.6	0.9
WHITE-Non-Hispanic			
All Causes:	38.1	42.8	33.3
All Accidents	21.1	25.4	16.7
(% Motor Vehicle)	(66.4)	(60.6)	(74.9)
Malignant Neoplasms	4.1	4.0	4.2
Suicide	2.0	2.7	1.4
Congenital Anomalies	2.0	2.7	1.4
Homicide	1.7	1.3	2.1
Influenza & Pneumonia	1.0	0.7	1.4
WHITE-Hispanic			
All Causes:	32.6	41.6	23.3
All Accidents	15.1	22.1	8.0
(% Motor Vehicle)	(69.5)	(64.7)	(83.8)
Malignant Neoplasms	6.2	7.2	5.3
Homicide	1.6	1.3	2.0
Heart Disease	1.3	1.3	1.3
Congenital Anomalies	1.0	0.7	1.3
Suicide	1.0	0.7	1.3
BLACK			
All Causes:	28.1	27.7	28.6
All Accidents	14.1	13.9	14.3
(% Motor Vehicle)	(100.0)	(100.0)	(100.0)
INDIAN			
All Causes:	25.5	38.7	12.6
All Accidents	10.2	12.9	7.6
(% Motor Vehicle)	(50.0)	(40.3)	(65.8)
Malignant Neoplasms	5.1	7.7	2.5
Suicide	2.6	2.6	2.5
Heart Disease	1.3	2.6	--
Homicide	1.3	2.6	--

AGES 15 TO 24

This age group could be termed "the age of violent death." Nationally, accidents were the leading cause of death in this age group, followed by suicide and homicide. In New Mexico, the suicide rate was almost twice the national level. Mortality rates were much higher for males than for females, and New Mexico rates were generally higher than nationally for all race/ethnic groups. Accidents, suicides, and homicides ranked as leading causes of death among Indians. Homicide ranked second for Blacks.

TABLE 3.12
AGES 15-24
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE

New Mexico Residents 1983-1985
United States 1984

	RATE/100,000 POPULATION		
NEW MEXICO ETHNIC/RACE	TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	96.8	141.6	51.1
All Accidents	49.4	75.7	22.5
(% Motor Vehicle)	(74.3)	(72.5)	(81.3)
Suicide	12.5	20.5	4.4
Homicide	12.0	18.1	5.8
Malignant Neoplasms	5.5	6.7	4.2
Heart Disease	2.7	3.3	2.1
Congenital Anomalies	1.3	1.5	1.0
Cerebrovascular	0.8	0.9	0.8
NEW MEXICO-ALL RACES			
All Causes:	140.7	216.9	62.8
All Accidents	77.8	125.8	28.6
(% Motor Vehicle)	(76.0)	(74.1)	(84.6)
Suicide	24.8	41.1	8.1
Homicide	13.4	21.6	5.1
Malignant Neoplasms	5.6	7.4	3.8
Heart Disease	3.6	4.5	2.7
Congenital Anomalies	1.3	1.8	0.8
Diabetes	0.9	1.1	0.8
WHITE-Non-Hispanic			
All Causes:	105.4	156.3	52.4
All Accidents	59.0	91.0	25.6
(% Motor Vehicle)	(77.8)	(74.8)	(88.7)
Suicide	19.7	30.1	8.7
Homicide	8.0	12.3	3.5
Malignant Neoplasms	5.1	6.7	3.5
Heart Disease	3.7	5.0	2.3
Congenital Anomalies	1.1	1.1	1.2
Diabetes	0.9	1.1	0.6
WHITE-Hispanic			
All Causes:	150.7	228.6	71.8
All Accidents	78.0	126.1	29.3
(% Motor Vehicle)	(76.3)	(75.5)	(79.5)
Suicide	27.4	47.3	7.3
Homicide	15.5	25.0	6.0
Malignant Neoplasms	5.6	7.2	4.0
Heart Disease	3.6	3.9	3.3
Congenital Anomalies	1.3	2.0	0.7
Cerebrovascular	1.0	0.7	1.3
BLACK			
All Causes:	63.6	94.0	25.9
All Accidents	28.9	41.8	12.9
(% Motor Vehicle)	(40.1)	(24.9)	(100.0)
Homicide	11.6	10.4	12.9
Suicide	5.8	10.4	--
Heart Disease	5.8	10.4	--
Malignant Neoplasms	5.8	10.4	--
Diabetes	5.8	10.4	--
INDIAN			
All Causes:	299.0	526.1	89.1
All Accidents	185.3	336.1	45.9
(% Motor Vehicle)	(73.5)	(71.3)	(88.2)
Suicide	42.1	76.0	10.8
Homicide	33.7	61.4	8.1
Malignant Neoplasms	8.4	11.7	5.4
Heart Disease	2.8	2.9	2.7
Influenza & Pneumonia	1.4	--	2.7
Alcoholism	1.4	--	2.7
Cerebrovascular	1.4	2.9	--

AGES 25 TO 44

For ages 25-44, accidents remained the leading cause of death. Nationally, malignant neoplasms and heart disease moved into the next two leading causes of death. In New Mexico, accidents, malignant neoplasms, suicide and homicide made up the four leading causes of death. Alcoholism ranked third for the Indian population. Homicide ranked second for the Black population. Mortality rates were higher for males than females in this age group. This pattern was true for most causes of death with the exception of malignant neoplasms. Again, motor vehicles were involved in a higher percentage of accident fatalities in the State than nationally.

TABLE 3.13
AGES 25-44
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE

New Mexico Residents 1983-1985 Average
United States 1984

	RATE/100,000 POPULATION		
NEW MEXICO ETHNIC/RACE	TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	156.9	215.0	100.0
All Accidents	35.5	56.4	14.9
(% Motor Vehicle)	(58.9)	(56.7)	(66.4)
Malignant Neoplasms	27.3	25.0	29.6
Heart Disease	21.1	31.7	10.8
Suicide	15.3	23.9	6.9
Homicide	13.3	21.3	5.2
Cirrhosis of Liver	5.9	8.4	3.6
Cerebrovascular	4.5	4.8	4.4
NEW MEXICO-ALL RACES			
All Causes:	183.0	258.4	108.6
All Accidents	57.1	90.2	24.6
(% Motor Vehicle)	(70.2)	(67.5)	(79.3)
Malignant Neoplasms	23.9	19.9	27.8
Suicide	22.7	35.7	9.8
Homicide	15.2	24.9	5.6
Heart Disease	13.7	20.3	7.3
Cirrhosis of Liver	9.1	13.9	4.4
Alcoholism	6.4	10.8	2.0
WHITE-Non-Hispanic			
All Causes:	141.6	192.0	90.5
All Accidents	43.2	65.8	20.3
(% Motor Vehicle)	(65.0)	(63.4)	(70.9)
Malignant Neoplasms	25.0	22.5	27.5
Suicide	21.8	34.0	9.4
Heart Disease	12.6	18.4	6.7
Homicide	8.0	12.1	3.9
Cirrhosis of Liver	3.3	4.4	2.2
Alcoholism	2.2	3.8	0.6
WHITE-Hispanic			
All Causes:	201.6	290.8	115.3
All Accidents	57.3	90.8	24.8
(% Motor Vehicle)	(72.1)	(69.7)	(80.6)
Homicide	24.1	42.7	6.2
Malignant Neoplasms	23.5	18.8	27.9
Suicide	22.1	33.5	11.1
Heart Disease	14.0	21.1	7.1
Cirrhosis of Liver	12.0	21.1	3.1
Alcoholism	6.8	11.5	2.2
BLACK			
All Causes:	198.7	275.3	108.3
All Accidents	45.2	75.1	9.8
(% Motor Vehicle)	(69.9)	(66.7)	(100.0)
Homicide	36.1	66.7	--
Heart Disease	27.1	41.7	9.8
Suicide	13.5	16.7	9.8
Malignant Neoplasms	9.0	8.3	9.8
Cerebrovascular	9.0	8.3	9.8
INDIAN			
All Causes:	441.5	680.8	229.1
All Accidents	177.2	308.4	60.8
(% Motor Vehicle)	(77.0)	(72.6)	(96.7)
Cirrhosis of Liver	44.0	61.7	28.4
Alcoholism	39.7	70.8	12.2
Suicide	36.5	70.8	6.1
Malignant Neoplasms	25.8	11.4	38.5
Homicide	25.8	34.3	18.2
Heart Disease	19.3	25.1	14.2

AGES 45 TO 64

Nationally, cancer, heart disease and stroke are the leading causes of death for this age group; the same ranking applies to New Mexicans in general.

Mortality rates for New Mexico Blacks and Indians were higher than national rates. Malignant neoplasms, heart disease and accidents were the top 3 leading causes for the non-Hispanic White population in the State. The same was true for the Hispanic White with the exception of cirrhosis of the liver as the number three cause of death. Accidents, malignant neoplasms and heart disease were the top three leading causes of death among Indians. The percentage of motor vehicle accidents remained higher for the State than the nation. Alcoholism and cirrhosis appeared in the leading causes for various race/ethnic groups.

TABLE 3.14
AGES 45-64
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

NEW MEXICO ETHNIC/RACE	RATE/100,000 POPULATION TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	902.9	1180.7	650.0
Malignant Neoplasms	308.9	350.5	271.6
Heart Disease	302.9	425.5	166.4
Cerebrovascular	39.1	44.0	35.1
All Accidents	34.1	51.0	18.8
(% Motor Vehicle)	(45.5)	(43.1)	(50.0)
Cirrhosis of Liver	29.1	40.7	18.8
C.O.P.D.	27.7	35.1	21.0
Diabetes	16.7	17.3	16.3
Suicide	16.7	25.3	9.0
NEW MEXICO-ALL RACES			
All Causes:	748.5	974.4	542.0
Malignant Neoplasms	230.4	250.3	212.2
Heart Disease	199.5	303.9	104.1
All Accidents	52.2	84.0	23.2
(% Motor Vehicle)	(49.2)	(47.9)	(53.9)
Cirrhosis of Liver	37.0	54.4	21.2
C.O.P.D.	28.5	31.8	25.5
Diabetes	26.4	27.1	25.8
Cerebrovascular	26.0	24.6	27.3
WHITE-Non-Hispanic			
All Causes:	746.5	966.5	542.0
Malignant Neoplasms	253.9	282.3	227.4
Heart Disease	209.7	327.1	100.6
All Accidents	44.2	69.9	20.2
(% Motor Vehicle)	(41.6)	(40.3)	(46.0)
C.O.P.D.	36.8	41.3	32.7
Suicide	27.0	40.0	14.9
Cirrhosis of Liver	25.5	35.2	16.6
Cerebrovascular	24.1	20.4	27.5
WHITE-Hispanic			
All Causes:	712.0	930.2	515.1
Malignant Neoplasms	186.9	187.4	186.5
Heart Disease	185.6	267.6	111.6
Cirrhosis of Liver	55.4	87.9	26.1
All Accidents	49.9	81.1	21.8
(% Motor Vehicle)	(63.3)	(64.4)	(60.1)
Diabetes	35.3	37.7	33.1
Cerebrovascular	28.4	29.0	27.9
Alcoholism	17.0	29.9	5.2
BLACK			
All Causes:	1198.5	1597.9	822.4
Heart Disease	369.5	445.9	297.5
Malignant Neoplasms	342.4	445.9	245.0
Diabetes	72.1	55.7	87.5
All Accidents	63.1	130.1	--
(% Motor Vehicle)	(42.8)	(42.8)	--
Cirrhosis of Liver	54.1	92.9	17.5
C.O.P.D.	45.1	55.7	35.0
Cerebrovascular	36.0	55.7	17.5
INDIAN			
All Causes:	933.4	1253.8	670.7
All Accidents	172.2	286.6	78.3
(% Motor Vehicle)	(51.6)	(45.8)	(68.8)
Malignant Neoplasms	161.4	143.3	176.2
Heart Disease	129.1	203.0	68.5
Diabetes	78.0	71.6	83.2
Cirrhosis of Liver	75.3	101.5	53.9
Cerebrovascular	37.7	47.8	29.4
Alcoholism	29.6	59.7	4.9

AGES 65 TO 74

By this age, accidents has dropped to the 7th ranked cause of death nationally. It is important to note that this drop was not due to a lower accident mortality rate for this age group, but rather to the higher mortality rates for heart disease, cancer, etc., which appear with advancing age.

In New Mexico, accidents ranked 5th. Mortality rates for many causes notably exceeded national levels among the various race/ethnic groups. Mortality from heart disease and cancer are much lower in the State than nationally.

Higher than national percentages of accidents involved motor vehicles. Accident mortality rates were highest for Indians.

TABLE 3.15
AGES 65-74
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE
New Mexico Residents 1983-1985 Average
United States 1984

NEW MEXICO ETHNIC/RACE	RATE/100,000 POPULATION TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	2848.1	3813.0	2096.4
Heart Disease	1102.7	1537.2	764.3
Malignant Neoplasms	835.1	1088.0	638.1
Cerebrovascular	177.0	205.4	154.9
C.O.P.D.	141.4	208.4	89.3
Diabetes	59.4	59.2	59.5
Influenza & Pneumonia	53.7	76.2	36.1
All Accidents	50.3	71.5	33.9
(% Motor Vehicle)	(35.8)	(34.8)	(37.5)
NEW MEXICO-ALL RACES			
All Causes:	2362.9	3095.8	1738.6
Heart Disease	741.9	1064.6	467.0
Malignant Neoplasms	688.6	838.5	560.9
C.O.P.D.	170.2	236.8	113.4
Cerebrovascular	124.8	135.6	115.5
All Accidents	85.1	127.4	49.0
(% Motor Vehicle)	(40.9)	(38.1)	(47.1)
Diabetes	78.3	74.0	81.9
Influenza & Pneumonia	43.1	50.1	37.1
WHITE-Non-Hispanic			
All Causes:	2435.3	3263.3	1736.9
Heart Disease	773.0	1145.1	459.1
Malignant Neoplasms	746.0	921.2	598.3
C.O.P.D.	211.0	297.4	138.1
Cerebrovascular	126.7	143.3	112.7
All Accidents	68.9	99.9	42.7
(% Motor Vehicle)	(36.0)	(31.3)	(45.2)
Diabetes	55.6	62.6	49.8
Influenza & Pneumonia	42.4	44.6	40.6
WHITE-Hispanic			
All Causes:	2227.1	2739.5	1781.3
Heart Disease	714.8	957.4	503.7
Malignant Neoplasms	579.4	650.1	517.8
Diabetes	124.9	103.5	143.5
Cerebrovascular	117.4	106.7	126.6
All Accidents	111.4	165.0	64.7
(% Motor Vehicle)	(48.7)	(49.0)	(47.9)
C.O.P.D.	90.3	119.7	64.7
Cirrhosis of Liver	52.7	84.1	25.3
BLACK			
All Causes:	3067.1	4336.6	2040.8
Heart Disease	925.9	1294.5	627.9
Malignant Neoplasms	723.4	1165.0	366.3
Cerebrovascular	376.2	388.3	366.3
Diabetes	202.5	64.7	314.0
All Accidents	86.8	194.2	--
(% Motor Vehicle)	(66.7)	(66.6)	--
Cirrhosis of Liver	86.8	129.4	52.3
Influenza & Pneumonia	57.9	129.4	--
C.O.P.D.	57.9	64.7	52.3
Septicemia	57.9	129.4	--
INDIAN			
All Causes:	1998.3	2539.2	1514.4
Malignant Neoplasms	467.7	630.3	322.2
Heart Disease	416.7	504.2	338.3
All Accidents	195.6	324.1	80.6
(% Motor Vehicle)	(39.1)	(33.3)	(59.9)
Diabetes	136.1	90.0	177.2
Cerebrovascular	68.0	126.1	16.1
C.O.P.D.	42.5	54.0	32.2
Nephritis, Nephrotic Syndrome and Nephrosis	42.5	18.0	64.4

TABLE 3.16
AGES 75 AND OVER
AGE-SPECIFIC DEATH RATES
LEADING CAUSES BY NEW MEXICO ETHNIC/RACE

New Mexico Residents 1983-1985 Average
United States 1984

	RATE/100,000 POPULATION		
NEW MEXICO ETHNIC/RACE	TOTAL	MALE	FEMALE
UNITED STATES			
All Causes:	8488.6	10321.5	7493.5
Heart Disease	3814.4	4431.0	3480.1
Malignant Neoplases	1351.2	1941.7	1029.7
Cerebrovascular	924.4	896.8	939.4
Influenza & Pneumonia	373.6	472.2	320.0
C.O.P.D.	285.1	522.5	154.5
Atherosclerosis	183.3	175.8	187.3
Diabetes	147.9	145.7	147.7
All Accidents	142.6	188.4	117.6
(% Motor Vehicle)	(19.2)	(22.7)	(16.2)
NEW MEXICO-ALL RACES			
All Causes:	7170.0	8689.4	6191.9
Heart Disease	2520.1	3009.3	2205.2
Malignant Neoplases	1140.0	1552.2	874.7
Cerebrovascular	687.3	663.0	702.9
C.O.P.D.	404.0	658.3	241.0
Influenza & Pneumonia	349.5	424.3	301.3
All Accidents	245.6	315.1	200.8
(% Motor Vehicle)	(15.4)	(19.3)	(11.5)
Diabetes	197.9	184.1	206.9
WHITE-Non-Hispanic			
All Causes:	7290.0	9207.8	6208.2
Heart Disease	2632.2	3214.9	2303.5
Malignant Neoplases	1155.8	1702.1	847.6
Cerebrovascular	714.1	697.0	723.7
C.O.P.D.	460.0	830.9	250.7
Influenza & Pneumonia	344.3	439.0	290.6
All Accidents	251.4	323.3	210.8
(% Motor Vehicle)	(14.5)	(18.0)	(11.5)
Diabetes	179.4	164.2	188.0
WHITE-Hispanic			
All Causes:	7115.2	8081.1	6329.2
Heart Disease	2454.4	2904.9	2087.7
Malignant Neoplases	1153.3	1343.5	998.5
Cerebrovascular	675.6	654.0	693.2
Influenza & Pneumonia	361.7	390.4	338.3
C.O.P.D.	329.8	420.8	255.8
Diabetes	236.6	233.2	239.3
All Accidents	213.8	253.5	181.5
(% Motor Vehicle)	(14.9)	(18.0)	(11.3)
BLACK			
All Causes:	7688.3	8478.5	7056.6
Heart Disease	2683.2	2903.6	2507.0
Malignant Neoplases	1599.6	2439.0	928.5
Cerebrovascular	980.4	580.7	1299.9
Diabetes	309.6	116.1	464.3
C.O.P.D.	154.8	348.4	--
Influenza & Pneumonia	154.8	116.1	185.7
All Accidents	154.8	232.3	92.9
(% Motor Vehicle)	--	--	--
Atherosclerosis	154.8	232.3	92.9
INDIAN			
All Causes:	6195.7	7140.8	5284.9
Heart Disease	1373.7	1571.0	1183.6
Malignant Neoplases	813.0	971.2	660.6
Influenza & Pneumonia	446.6	571.3	330.3
All Accidents	392.5	599.8	192.7
(% Motor Vehicle)	(28.6)	(33.3)	(14.3)
Cerebrovascular	336.4	399.9	275.3
Diabetes	224.3	171.4	275.3
C.O.P.D.	126.2	199.9	55.1

AGES 75 AND OVER

Heart disease, cancer and stroke ranked in that order as leading causes of death nationally as well as in New Mexico; however, the rates were lower than nationally for heart disease, cancer, stroke and influenza/pneumonia, but higher for C.O.P.D., diabetes and accidents. Overall, mortality rates were lower than nationally, and lowest for the Indian.

Accident mortality rates in the 75 and older age group are the highest of any age group, but the lower ranking as a cause of death results from higher rates for those causes usually associated with old age. Accident mortality is highest for the Indian. Accident mortality in this age group is high for both males and females, while for younger ages, rates for males had been much higher.

In Tables 3.18 and 3.19 on Pages 48 and 49, Age and Cause Specific Death Rates for the United States and New Mexico's various populations are presented for males and females respectively. The comparisons to be drawn are many and are left to the interest of the reader.

CAUSES BY COUNTY

Age-adjusted mortality rates by sex and county are presented in Table 3.17.

The rates have been age-adjusted to eliminate the effect created by the variations in age structure in the county populations.

The highest overall mortality rates in the State are found in De Baca County for males and in Harding County for females.

De Baca County males registered the highest accident mortality rate, while Catron County had the highest accident mortality rate among females.

The highest mortality rate for alcoholism was that for San Miguel County males. Also notably high were the rates for males in McKinley, Taos, Rio Arriba, and Socorro counties. For females, the rate in Quay County was highest.

Guadalupe County registered the highest homicide fatality rate for males. For females the homicide rate was highest in Cibola County.

High suicide rates for males were noted in several of the State's counties, highest was Harding County. The highest overall suicide rate for females was in Sierra County.

Cirrhosis of the liver mortality rates were exceptionally high for Rio Arriba County males, and for females the rate was the highest for Sierra County.

Stroke mortality rates are generally higher for females than for males. The highest cancer mortality in the State was for De Baca County males. For females, the rate was highest in Quay County.

Further breakdown of selected causes of death by county is presented in Table A-10 through Table A-12 of the Appendix. A breakdown of deaths by county and city is presented in Table A-7.

TABLE 3.17
AGE-ADJUSTED MORTALITY RATES BY SELECTED CAUSES NEW MEXICO RESIDENTS BY COUNTY 1983-85 AVERAGE
(Rates Per 100,000 Population)

COUNTY	ALL CAUSES	Malignant Neoplasms 140-209	Diabetes Mellitus 250	Alcoholism 303	Diseases of the Heart 390-398 402, 404 410-429	Cerebro-vascular Disease 430-438	Athero-sclerosis 440	Influenza and Pneumonia 480-487	Chronic Obstructive Pulmonary Diseases 490-496	Chronic Liver Disease and Cirrhosis 571	Accidents E800-E949	Suicide E950-E959	Homicide E960-E978	Nephritis Nephrotic Syndrome Nephrosis 580-589
DISTRICT I														
Cibola:														
Male	709.3	123.7	16.3	8.7	172.3	31.7	2.8	9.7	11.8	17.7	128.9	50.3	30.2	7.6
Female	764.1	145.6	24.0	--	203.5	79.3	14.6	32.6	13.7	9.0	63.9	--	24.5	13.6
McKinley:														
Male	1101.9	148.5	25.9	27.8	236.5	23.5	4.4	41.5	46.3	33.3	216.7	44.5	18.6	10.3
Female	808.3	124.8	51.8	5.2	164.1	50.7	6.6	32.7	23.8	21.7	54.2	6.3	5.5	10.3
San Juan:														
Male	866.4	148.1	23.8	11.1	195.6	47.1	5.5	19.6	64.5	11.8	143.8	25.3	16.0	4.3
Female	613.1	134.7	27.6	4.9	151.3	42.3	5.8	13.9	25.9	5.1	61.6	13.8	4.9	6.9
DISTRICT II														
Colfax:														
Male	859.0	175.0	20.8	--	237.8	21.6	--	11.5	51.4	18.6	118.7	22.3	32.1	13.3
Female	619.2	121.2	12.2	--	248.6	56.5	13.2	17.8	17.2	5.4	25.5	3.7	--	--
Harding:														
Male	681.9	198.2	--	--	140.2	24.8	--	24.8	--	--	--	85.8	--	--
Female	921.0	114.2	114.2	--	415.0	--	40.2	--	--	--	47.1	--	--	40.2
Los Alamos:														
Male	665.3	110.0	7.6	--	173.3	78.0	--	12.2	33.0	8.5	58.7	17.1	--	12.2
Female	546.9	81.3	38.1	--	98.4	50.2	21.8	--	18.1	9.5	60.8	11.3	--	6.5
Mora:														
Male	781.2	148.9	7.1	13.5	201.1	5.7	7.1	7.1	11.5	47.6	121.2	35.3	16.5	--
Female	590.2	180.8	57.6	--	141.1	31.3	--	12.7	--	--	--	--	--	--
Rio Arriba:														
Male	1081.5	174.5	20.7	22.0	276.1	32.8	7.5	24.4	36.8	49.4	183.5	37.8	40.6	9.6
Female	884.8	177.6	33.0	6.0	219.8	78.6	15.4	20.7	29.5	22.6	80.1	7.7	6.4	-
San Miguel:														
Male	863.3	104.7	26.0	42.7	250.3	35.6	7.0	42.4	38.6	36.5	75.6	42.8	18.9	2.7
Female	644.6	123.7	27.6	--	207.4	26.1	16.7	32.1	40.9	6.0	43.3	5.7	13.9	5.5
Santa Fe:														
Male	694.7	140.4	11.5	5.7	179.5	25.5	2.7	22.9	35.9	23.4	76.1	38.8	12.8	3.7
Female	644.2	164.9	22.8	5.0	165.4	50.3	1.9	23.4	29.1	13.8	39.5	3.6	4.8	4.6
Taos:														
Male	857.9	162.1	25.2	23.9	176.5	20.4	3.3	20.6	17.5	30.2	127.9	32.5	28.3	7.8
Female	714.1	138.7	24.1	3.3	170.4	66.7	4.4	41.4	19.7	3.1	49.2	10.1	3.1	16.8
Union:														
Male	915.8	220.7	9.4	--	293.0	45.4	9.8	28.6	32.1	11.8	109.9	58.2	17.9	19.5
Female	743.4	191.0	8.7	--	240.6	55.8	18.1	8.7	8.7	--	29.0	--	--	--
DISTRICT III														
Bernalillo:														
Male	851.8	173.4	13.1	5.6	259.0	42.8	5.5	22.5	47.6	18.5	64.2	28.5	15.2	8.2
Female	688.7	155.7	21.8	1.9	196.6	59.1	11.2	22.7	33.0	8.0	28.9	10.3	3.5	9.0
Sandoval:														
Male	741.2	147.4	12.0	17.3	185.4	23.1	1.2	18.3	26.3	30.1	117.8	28.3	6.8	1.3
Female	600.7	152.1	21.6	--	163.4	46.4	3.1	12.5	19.0	4.5	35.8	7.9	3.3	6.1
Torrance:														
Male	878.8	131.6	6.1	17.0	280.9	29.0	11.7	6.1	48.0	29.0	130.1	29.7	40.6	11.6
Female	584.6	101.0	38.8	--	242.3	36.9	--	13.1	13.1	7.5	27.9	--	--	--
Valencia:														
Male	866.2	191.5	28.3	5.5	234.0	32.3	3.3	25.4	36.2	8.2	116.1	35.8	8.1	3.0
Female	684.5	153.4	38.3	--	171.7	55.9	8.6	30.2	31.6	10.5	39.8	3.6	2.1	14.4
DISTRICT IV														
Chaves:														
Male	865.4	177.9	24.0	12.2	251.3	41.1	8.6	22.5	52.8	14.0	76.3	27.1	26.2	8.2
Female	724.3	154.6	23.1	3.1	214.9	67.6	9.8	21.3	34.0	9.6	40.8	4.0	4.5	8.2
Curry:														
Male	784.1	166.0	17.8	6.9	256.6	43.3	8.9	16.2	37.0	16.6	79.7	11.3	9.3	7.9
Female	629.9	118.0	14.0	5.3	194.7	71.1	15.6	28.8	26.2	3.1	28.3	4.8	1.1	4.6
DeBaca:														
Male	1206.6	295.5	9.3	--	264.2	28.4	--	21.5	79.3	41.6	236.4	33.4	--	--
Female	781.2	190.6	33.8	--	233.2	41.8	--	14.3	14.3	14.3	44.8	30.0	--	--
Eddy:														
Male	952.1	186.3	24.4	12.6	295.6	36.6	5.5	29.3	51.5	22.9	108.4	31.4	6.8	8.2
Female	768.6	173.2	30.2	3.4	219.5	64.5	6.7	42.2	34.3	7.0	36.3	2.3	4.0	5.9
Guadalupe:														
Male	816.7	82.7	14.0	14.5	266.0	36.7	--	12.5	10.2	19.1	87.2	12.9	53.2	--
Female	870.3	126.4	46.5	--	267.6	--	--	93.0	38.1	--	73.0	--	--	16.4
Lea:														
Male	899.4	193.6	21.0	2.4	262.1	57.1	10.0	22.8	53.2	13.1	98.2	21.5	17.7	7.9
Female	802.4	154.9	28.0	--	276.4	73.5	13.6	32.8	36.5	10.0	35.9	5.8	6.0	4.2
Lincoln:														
Male	992.7	178.9	36.6	--	307.5	61.6	10.0	10.0	56.8	13.8	93.9	44.9	5.2	15.9
Female	732.9	158.5	12.2	--	196.8	37.4	30.3	13.5	45.6	4.2	51.5	5.3	14.8	--
Quay:														
Male	880.9	174.3	12.2	3.7	336.2	26.1	3.7	19.6	41.4	8.9	100.3	41.1	6.6	8.3
Female	781.6	214.8	24.7	15.1	286.8	33.9	15.6	15.1	19.4	--	49.6	--	3.9	10.2
Roosevelt:														
Male	895.8	153.3	19.7	5.2	322.8	54.2	3.9	15.3	46.7	22.7	93.4	8.9	3.7	11.2
Female	684.8	139.1	10.2	4.6	247.6	70.4	13.1	18.6	12.5	--	47.7	--	8.0	3.2
DISTRICT V														
Catron:														
Male	993.2	184.0	27.9	--	256.4	112.3	13.3	29.4	44.3	--	121.1	60.4	--	27.9
Female	778.0	116.4	43.7	--	217.7	70.3	--	22.8	--	--	87.0	22.8	23.9	--
Dona Ana:														
Male	647.9	143.4	15.7	5.8	173.6	24.8	7.8	13.9	44.5	18.5	63.2	22.8	13.9	4.5
Female	670.8	140.3	27.6	1.4	198.5	70.6	18.1	23.1	28.7	2.2	19.9	3.4	2.8	4.4
Grant:														
Male	783.6	141.8	16.1	2.6	241.1	25.2	--	18.7	49.3	18.3	100.4	38.9	6.6	5.0
Female	669.0	134.0	18.2	--	161.6	86.8	11.5	33.0	49.3	7.7	25.3	5.1	--	2.8
Hidalgo:														
Male	887.4	185.2	12.7	--	261.8	63.3	18.6	39.4	33.6	30.1	84.7	--	--	--
Female	737.1	193.6	62.2	--	191.2	83.5	11.6	22.2	22.7	9.8	--	14.9	9.9	--
Luna:														
Male	856.0	187.8	32.2	15.9	238.5	60.4	2.0	13.7	53.5	5.6	87.6	35.1	20.5	10.3
Female	638.7	120.3	41.9	--	193.1	77.8	--	26.5	6.8	--	22.1	20.5	--	7.9
Otero:														
Male	803.5	162.0	16.1	5.1	230.4	38.2	11.1	15.8	47.1	29.8	81.7	34.0	9.3	8.5
Female	752.8	143.2	24.0	--	248.7	87.4	8.6	19.5	26.4	13.0	31.8	1.3	2.7	5.9
Sierra:														
Male	830.4	148.0	7.7	6.7	216.7	22.9	5.4	15.0	48.5	6.2	73.5	83.4	11.9	12.0
Female	768.5	98.2	9.3	2.5	265.4	49.2	23.4	30.3	17.6	30.4	40.0	45.9	--	5.4
Socorro:														
Male	754.9	136.3	--	20.4	198.2	49.4	--	21.7	44.8	17.8	81.0	31.6	4.3	--
Female	685.6	137.7	12.9	6.6	157.1	65.7	--	29.0	19.7	--	35.1	--	4.6	18.8

Rates per 100,000 population.

TABLE 3.18
AGE-SPECIFIC DEATH RATES FOR SELECTED CAUSES—MALES
New Mexico Residents 1983-85 Average
United States 1984

M A L E (By Age Group)	ALL CAUSES	DISEASES OF HEART 390-398, 402 404-424	MALIGNANT NEOPLASMS 140-208	STROKE 430-438	ALL ACCIDENTS E800-E949	MOTOR VEHICLE E810-E825	ALL OTHER E800-807 E826-E949	INFLUENZA & PNEUMONIA 480-487	DIABETES MELLITUS 250	CIRRHOSIS OF LIVER 571	ATHEROSCLEROSIS 440	SUICIDE E950-E959	C.O.P.D. 490-496	HOMICIDE E960-978	ALCOHOLISM 303
UNITED STATES															
1 - 4	57.0	2.5	4.3	0.4	23.9	8.0	15.9	1.4	0.0	0.1	0.0	--	0.3	2.4	--
5 - 14	32.1	1.0	4.0	0.3	16.5	8.3	8.2	0.5	0.1	0.1	0.0	1.0	0.3	1.3	5.8
15 - 24	141.6	3.3	6.7	0.9	75.7	54.9	20.9	0.7	0.3	0.2	0.0	20.5	0.4	18.1	117.3
25 - 34	173.9	11.3	12.8	2.3	61.9	37.0	24.9	2.0	1.5	3.9	0.0	24.9	0.5	23.5	882.7
35 - 44	270.2	59.7	41.7	7.9	48.6	25.2	23.4	4.0	4.2	14.5	0.2	22.6	1.7	18.2	3251.8
45 - 54	676.6	243.2	177.0	24.5	48.7	22.4	26.3	8.8	10.3	32.5	1.0	23.7	11.0	14.1	6645.3
55 - 64	1705.2	670.7	530.7	63.6	53.1	21.9	31.2	23.7	24.9	49.3	5.2	27.2	60.3	9.4	8993.7
65 - 74	3813.0	1537.2	1088.0	205.4	71.5	24.9	46.6	76.2	59.2	55.5	22.3	33.5	208.4	6.5	7310.7
75 - 84	8445.9	3535.2	1826.6	679.7	147.6	42.0	105.6	306.5	129.4	47.0	103.5	48.5	482.2	6.2	4425.1
85 +	18119.1	8154.8	2427.2	1797.0	357.4	53.4	304.0	1172.1	213.6	29.8	477.4	51.7	691.9	7.3	1683.9
NEW MEXICO (ALL RACES)															
1 - 4	73.8	2.3	2.9	0.6	36.0	15.1	20.9	2.9	0.6	--	--	--	--	5.8	--
5 - 14	41.1	1.1	5.7	0.3	22.1	13.6	8.5	0.6	--	--	--	1.7	--	1.4	--
15 - 24	216.9	4.5	7.4	0.8	125.8	93.2	32.6	0.3	1.1	0.3	--	41.1	--	21.6	--
25 - 34	252.8	11.9	12.8	3.0	102.4	72.0	30.4	1.4	2.2	6.5	--	42.1	0.3	31.2	6.5
35 - 44	265.7	31.2	29.4	3.6	74.0	46.3	27.7	2.9	3.2	23.7	--	27.3	2.5	16.5	16.5
45 - 54	616.0	169.5	135.3	13.7	95.8	46.9	39.0	7.9	15.3	41.6	1.1	32.6	7.4	12.6	22.6
55 - 64	1378.4	455.3	379.9	36.8	81.9	32.7	49.3	12.5	40.4	69.9	3.0	38.6	59.4	10.7	17.2
65 - 74	3095.8	1064.6	838.5	135.6	127.4	48.5	78.9	50.1	74.0	54.3	13.2	41.9	236.8	9.0	20.6
75 - 84	7134.0	2458.7	1443.5	525.1	218.9	52.3	166.6	265.4	164.7	56.2	87.2	54.3	627.8	7.8	19.4
85 +	15117.3	5284.7	2001.8	1233.1	712.6	96.1	616.5	1081.0	264.2	24.0	272.2	80.1	784.8	8.0	16.0
NEW MEXICO NON-HISPANIC WHITE															
1 - 4	60.3	--	5.9	--	29.4	10.3	19.1	1.5	--	--	--	--	--	1.5	--
5 - 14	42.8	0.7	4.0	--	25.4	15.4	10.0	0.7	--	--	--	2.7	--	1.3	--
15 - 24	156.3	5.0	6.7	0.6	91.0	68.1	22.9	--	1.1	0.6	--	30.1	--	12.3	--
25 - 34	188.7	9.9	14.3	2.5	81.3	50.7	30.5	0.5	2.5	1.5	--	36.9	0.5	13.8	1.0
35 - 44	196.3	29.1	32.8	1.9	46.4	30.3	16.1	--	2.5	8.0	--	30.3	3.1	9.9	7.4
45 - 54	574.7	180.7	160.0	11.2	70.5	31.0	39.6	5.2	6.9	24.9	0.9	33.6	7.7	9.5	14.6
55 - 64	1366.0	476.4	407.1	29.8	69.3	25.4	43.9	12.3	30.7	45.6	2.6	46.5	75.5	6.1	11.4
65 - 74	3263.3	1145.1	921.2	143.3	99.9	31.3	68.6	44.6	62.6	43.3	14.4	54.2	297.4	3.6	15.7
75 - 84	7745.8	2652.6	1620.8	576.6	221.3	46.8	174.6	271.2	137.1	53.0	102.9	84.2	826.0	3.1	21.8
85 +	15449.1	5615.4	2049.2	1210.9	758.5	106.5	652.0	1157.7	279.4	13.3	319.4	119.8	851.6	13.3	13.3
NEW MEXICO HISPANIC WHITE															
1 - 4	58.1	1.3	1.3	1.3	26.4	10.6	15.9	2.6	--	--	--	--	--	6.6	--
5 - 14	41.6	1.3	7.2	0.7	22.1	14.3	7.8	0.7	--	--	--	0.7	--	1.3	--
15 - 24	228.6	3.9	7.2	0.7	126.1	95.2	30.9	0.7	0.7	--	--	47.3	--	25.0	--
25 - 34	284.7	12.7	12.7	4.0	102.8	79.9	22.9	2.4	1.6	7.9	--	41.9	--	53.0	4.0
35 - 44	299.3	32.8	27.3	6.6	74.3	40.4	33.9	1.1	5.3	39.3	--	21.8	--	28.4	21.8
45 - 54	614.2	156.9	87.7	16.9	77.6	54.0	23.6	8.4	18.6	60.7	1.7	30.4	5.1	20.2	28.7
55 - 64	1353.4	415.7	320.8	45.2	85.9	49.7	36.2	9.0	63.3	124.3	4.5	18.1	27.1	13.6	31.6
65 - 74	2739.5	957.4	650.1	106.7	165.0	80.9	84.1	64.7	103.5	84.1	9.7	19.4	119.7	19.4	29.1
75 - 84	6339.6	2329.5	1194.0	486.7	155.7	32.4	123.3	227.1	233.6	64.9	71.4	--	343.9	19.5	19.5
85 +	14302.3	4960.6	1877.6	1251.7	602.7	92.7	510.0	973.6	231.8	46.4	185.4	23.2	695.4	--	23.2
NEW MEXICO BLACK															
1 - 4	141.3	23.6	--	--	70.7	47.1	23.6	--	23.6	--	--	--	--	23.6	--
5 - 14	27.7	--	--	--	13.9	13.9	--	--	--	--	--	--	--	--	--
15 - 24	94.0	10.4	10.4	--	41.8	10.4	31.3	--	10.4	--	--	10.4	--	10.4	--
25 - 34	237.7	26.4	--	13.2	66.0	52.8	13.2	--	--	--	--	26.4	--	66.0	--
35 - 44	339.7	67.9	22.6	--	90.6	45.3	45.3	22.6	--	22.6	--	--	22.6	67.9	--
45 - 54	886.4	158.3	189.9	31.7	158.3	63.3	95.0	31.7	63.3	31.7	--	--	63.3	--	31.7
55 - 64	2609.1	854.7	809.7	90.0	90.0	45.0	45.0	45.0	45.0	179.9	--	--	45.0	90.0	--
65 - 74	4336.6	1294.5	1165.0	388.3	194.2	129.4	64.7	129.4	64.7	129.4	--	--	64.7	64.7	--
75 - 84	7758.6	2586.2	2298.9	574.7	287.4	--	287.4	143.7	143.7	--	--	--	287.4	--	--
85 +	11515.2	4242.4	3030.3	606.1	--	--	--	--	--	--	1212.1	--	606.1	--	--
NEW MEXICO INDIAN															
1 - 4	164.2	9.1	--	--	86.7	41.1	45.6	9.1	--	--	9.1	--	--	13.7	--
5 - 14	38.7	2.6	7.7	--	12.9	5.2	7.7	--	--	--	--	2.6	--	2.6	--
15 - 24	526.1	2.9	11.7	2.9	336.1	239.7	96.4	--	--	--	--	76.0	--	61.4	--
25 - 34	640.3	19.3	7.7	--	297.0	219.9	77.1	3.9	3.9	42.4	3.9	92.6	--	54.0	65.6
35 - 44	739.7	33.6	16.8	5.6	325.0	229.8	95.3	33.6	--	89.7	33.6	39.2	5.6	5.6	78.5
45 - 54	1110.8	144.0	113.1	20.6	308.5	195.4	113.1	30.9	82.3	133.7	30.9	51.4	--	10.3	82.3
55 - 64	1451.8	284.7	185.0	85.4	256.2	42.7	213.5	28.5	56.9	56.9	28.5	56.9	14.2	42.7	28.5
65 - 74	2539.2	504.2	630.3	126.1	324.1	108.0	216.1	36.0	90.0	36.0	36.0	--	54.0	18.0	54.0
75 - 84	5228.8	1274.5	784.3	196.1	490.2	228.8	261.4	457.5	130.7	65.4	457.5	--	130.7	--	--
85 +	20408.2	3628.1	2267.6	1814.1	1360.5	--	1360.5	1360.5	453.5	--	1360.5	--	680.3	--	--

New Mexico Rates by Race/Ethnic based on 1985 population estimates

0.0 Equals less than one-tenth of one percent.

Rates Per 100,000 Population

TABLE 3.19
AGE-SPECIFIC DEATH RATES FOR SELECTED CAUSES—FEMALES
New Mexico Residents 1983-85 Average
United States 1984

F E M A L E (By Age Group)	ALL CAUSES	DISEASES OF HEART 390-398, 402 404-424	MALIGNANT NEOPLASMS 140-208	STROKE 430-438	ALL ACCIDENTS E800-E949	MOTOR VEHICLE E810-E825	ALL OTHER E800-807 E826-E949	INFLUENZA & PNEUMONIA 480-487	DIABETES MELLITUS 250	CIRRHOSIS OF LIVER 571	ATHEROSCLEROSIS 440	SUICIDE E950-E959	C.O.P.D. 490-496	HOMICIDE E960-978	ALCOHOLISM 303
UNITED STATES															
1 - 4	46.5	2.3	3.6	0.4	15.5	5.7	9.8	1.5	0.0	0.1	0.0	--	0.4	2.4	--
5 - 14	21.1	1.0	3.0	0.3	8.0	4.9	3.0	0.5	0.1	0.0	0.0	0.3	0.2	1.3	--
15 - 24	51.1	2.1	4.2	0.8	22.5	18.3	4.2	0.5	0.3	0.1	--	4.4	0.4	5.8	30.2
25 - 34	68.5	4.8	13.1	2.2	15.7	10.7	5.0	1.2	1.3	1.8	0.0	6.1	0.6	5.9	320.3
35 - 44	141.5	18.3	51.4	7.2	14.4	9.2	5.2	2.1	2.8	5.8	0.1	7.7	1.5	4.7	1090.6
45 - 54	374.6	75.1	164.4	20.8	16.7	8.8	7.9	4.6	7.6	14.4	0.4	9.2	8.0	3.3	1758.9
55 - 64	918.4	255.4	375.6	48.8	20.8	10.1	10.6	10.7	24.4	22.4	3.1	8.5	33.4	2.6	2178.7
65 - 74	2096.4	764.3	638.1	154.9	33.9	12.7	21.1	36.1	59.5	26.7	13.6	7.3	89.3	2.3	1834.8
75 - 84	5188.2	2283.2	944.2	594.7	83.2	20.0	63.2	162.4	124.2	24.9	79.5	6.3	144.8	3.4	572.9
85 +	14053.9	6885.9	1271.5	1918.9	216.3	13.5	202.8	766.6	218.1	14.2	492.8	4.9	185.2	3.1	262.9
NEW MEXICO (ALL RACES)															
1 - 4	54.2	3.0	2.4	0.6	21.7	13.2	8.4	3.0	--	--	--	--	--	0.6	--
5 - 14	26.1	0.9	4.3	0.3	11.6	9.0	2.6	0.9	0.3	--	--	1.4	--	1.7	--
15 - 24	62.8	2.7	3.8	0.5	28.6	24.2	4.3	0.5	0.8	0.5	--	8.1	0.5	5.1	0.5
25 - 34	82.3	5.4	15.9	1.1	23.7	18.8	4.8	0.8	0.8	1.9	--	8.6	0.8	5.6	2.7
35 - 44	143.2	9.9	43.4	4.6	25.7	20.5	5.3	1.1	2.1	7.8	0.4	11.3	1.4	5.6	1.1
45 - 54	321.6	45.4	131.7	18.2	22.2	13.3	8.9	3.9	8.9	16.8	0.5	8.9	7.4	3.9	4.9
55 - 64	778.0	166.9	298.4	37.0	24.3	11.6	12.7	7.9	43.8	25.9	0.5	12.1	44.9	2.1	7.4
65 - 74	1738.6	467.0	560.9	115.5	49.0	23.1	25.9	37.1	81.9	21.7	14.7	4.9	113.4	1.4	5.6
75 - 84	4221.3	1428.7	777.9	435.6	143.9	25.9	118.0	163.4	181.5	19.4	77.8	11.7	197.1	5.2	1.3
85 +	12962.7	4873.3	1207.2	1621.5	396.5	13.4	383.1	775.1	294.0	8.9	378.6	4.5	392.0	13.4	--
NEW MEXICO NON-HISPANIC WHITE															
1 - 4	46.4	1.5	3.1	--	18.5	15.5	3.1	1.5	--	--	--	--	--	--	--
5 - 14	33.3	0.7	4.2	0.7	16.7	12.5	4.2	1.4	0.7	--	--	1.4	--	2.1	--
15 - 24	52.4	2.3	3.5	--	25.6	22.7	2.9	--	0.6	--	--	8.7	0.6	3.5	0.6
25 - 34	64.7	6.5	13.4	0.5	19.9	15.4	4.5	0.5	1.0	1.0	--	8.0	0.5	3.0	0.5
35 - 44	123.0	6.9	45.2	2.5	20.7	13.2	7.5	0.6	1.3	3.8	0.6	11.7	1.3	5.0	0.6
45 - 54	315.5	42.6	136.1	19.2	18.4	9.2	9.2	3.3	8.3	10.9	0.8	14.2	9.2	2.5	5.8
55 - 64	754.4	154.9	313.0	35.2	21.9	9.4	12.5	7.0	23.5	21.9	0.8	15.7	54.8	3.1	7.8
65 - 74	1736.9	459.1	598.3	112.7	42.7	19.3	23.4	40.6	49.8	19.3	16.3	7.1	138.1	1.0	6.1
75 - 84	4147.2	1459.9	745.5	445.1	152.0	27.5	124.6	155.7	164.9	11.0	37.9	16.5	203.3	3.7	1.8
85 +	13417.9	5254.4	1204.7	1698.1	416.5	12.8	403.7	762.5	269.1	6.4	390.9	6.4	416.5	12.8	--
NEW MEXICO HISPANIC WHITE															
1 - 4	47.5	2.7	2.7	1.4	10.9	4.1	6.8	5.4	--	--	5.4	--	--	1.4	--
5 - 14	23.3	1.3	5.3	--	8.0	6.7	1.3	0.7	--	--	0.7	1.3	--	2.0	--
15 - 24	71.8	3.3	4.0	1.3	29.3	23.3	6.0	0.7	--	1.3	0.7	7.3	0.7	6.0	--
25 - 34	90.1	3.9	18.5	2.3	23.9	17.7	6.2	--	3.9	1.5	--	10.0	1.5	6.9	3.9
35 - 44	149.5	11.5	40.8	6.3	26.1	23.0	3.1	2.1	--	5.2	2.1	12.5	1.0	5.2	--
45 - 54	296.5	46.1	116.7	18.4	21.5	15.4	6.1	3.1	3.1	23.0	3.1	1.5	4.6	6.1	3.1
55 - 64	801.8	197.4	278.0	40.3	22.2	10.1	12.1	8.1	8.1	30.2	8.1	6.0	24.2	--	8.1
65 - 74	1781.3	503.7	517.8	126.6	64.7	31.0	33.8	33.8	2.8	25.3	33.8	--	64.7	2.8	2.8
75 - 84	4426.3	1395.5	889.5	410.1	122.5	21.3	101.2	186.4	--	37.3	186.4	--	207.7	5.3	--
85 +	12868.4	4466.4	1372.9	1665.8	384.4	18.3	366.1	860.3	--	18.3	860.3	--	421.0	--	--
NEW MEXICO BLACK															
1 - 4	24.9	--	--	--	--	--	--	--	--	--	--	--	--	--	--
5 - 14	28.6	--	--	--	14.3	14.3	--	--	--	--	--	--	--	--	--
15 - 24	25.9	--	--	--	12.9	12.9	--	--	--	--	--	--	--	12.9	--
25 - 34	82.1	--	--	--	16.4	16.4	--	--	--	--	--	16.4	--	--	--
35 - 44	147.4	24.6	24.6	24.6	--	--	--	--	--	--	--	--	--	--	--
45 - 54	511.0	210.4	240.5	--	--	--	--	--	--	--	--	--	--	--	--
55 - 64	1256.3	418.8	251.3	41.9	--	--	--	--	209.4	41.9	--	--	83.8	--	--
65 - 74	2040.8	627.9	366.3	366.3	--	--	--	--	314.0	52.3	--	--	52.3	--	--
75 - 84	5797.1	1932.4	966.2	1207.7	120.8	--	120.8	241.5	362.3	120.8	120.8	--	--	--	--
85 +	11245.0	4417.7	803.2	1606.4	--	--	--	--	803.2	--	--	--	--	--	--
NEW MEXICO INDIAN															
1 - 4	110.3	9.2	--	--	73.6	41.4	32.2	--	--	--	--	--	--	--	--
5 - 14	12.6	--	2.5	--	7.6	5.0	2.5	--	--	--	--	2.5	--	--	--
15 - 24	89.1	2.7	5.4	--	45.9	40.5	5.4	2.7	--	--	--	10.8	--	8.1	2.7
25 - 34	185.7	7.0	28.0	--	52.6	49.1	3.5	7.0	3.5	10.5	--	7.0	--	21.0	14.0
35 - 44	288.6	24.1	52.9	9.6	72.2	72.2	--	--	--	52.9	--	4.8	4.8	14.4	9.6
45 - 54	488.5	33.1	149.0	16.6	74.5	49.7	24.8	16.6	33.1	49.7	--	--	8.3	8.3	8.3
55 - 64	934.2	119.8	215.6	47.9	83.8	59.9	24.0	24.0	155.7	59.9	--	--	12.0	--	--
65 - 74	1514.4	338.3	322.2	16.1	80.6	48.3	32.2	16.1	177.2	32.2	16.1	--	32.2	--	16.1
75 - 84	4119.0	915.3	686.5	228.8	152.6	38.1	114.4	152.6	228.8	38.4	76.3	--	76.3	38.1	--
85 +	9308.6	1879.3	593.5	395.6	296.7	--	296.7	791.3	395.6	--	147.8	--	--	98.9	--

New Mexico Rates by Race/Ethnic based on 1985 population estimates

0.0 Equals less than one-tenth of one percent.

Rates Per 100,000 Population

STANDARDIZED MORTALITY RATIOS

Another tool useful in analyzing the mortality of a population is the Standardized Mortality Ratio (SMR).

During the four year period, 1979-1982, there were observed (recorded) 35,589 deaths of New Mexico residents, an average of 8,897 deaths per year.

If New Mexico's population had an age and female/male distribution similar to that of the nation as a whole, and if New Mexicans had been dying at the same rate as was experienced nationally during this time, then we would have expected a total of 37,244 New Mexicans to have died during this period of time. Since only 35,589 deaths were observed, the conclusion then is that New Mexicans generally experienced a lower level of mortality than was experienced nationwide.

The ratio of the observed deaths (35,589) to the expected deaths (37,244) multiplied by 100 is referred to as the standardized mortality ratio, which in this case for New Mexico deaths is "96" (Table 3.20). In other words, the State's overall mortality was 96% of that experienced nationally.

Scanning across the "SMR" row in Table 3.20, reveals excess mortality for the State's population for accidents (53%), motor vehicle accidents in particular (78%), COPD (chronic obstructive pulmonary disease) (64%), diabetes (53%), suicide (68%), influenza/pneumonia (7%), cirrhosis (10%), homicide (29%), and alcoholism mortality in New Mexico is nearly seven times that based on national data.

Mortality from other causes of death are lower in New Mexico than nationally. For example, heart disease in the State is 62% of that nationally, stroke is 72% and cancer is 91%. One caution, however, is that some causes, for example cancer mortality, may overall exhibit an SMR lower than the national mortality experience but further examination by cancer site and sex or ethnicity may and does reveal higher rates for New Mexicans for particular sites.

Another consideration in analyzing the mortality situation of the State is the actual number of deaths and the percent that figure represents of the total. Heart disease mortality in New Mexico, for example, is only 62% of the national level, however, heart disease accounts for one out of every four deaths in New Mexico. Cancer accounts for one out of every five deaths and accidents for one out of every ten. These leading causes together account for 54% of all deaths in the State.

For the State's four largest race/ethnic groups, SMR's indicate an overall "excess" mortality for the Black and Indian populations (Table 3.20).

Deaths from accidents is higher than expected for all four groups, with the level for the Indian being almost four times that nationally, as accidents account for over a fourth of Indian deaths.

COPD deaths is twice the expected number for Non-Hispanic Whites. Diabetes mortality runs twice the expected level for the Hispanic, Black and Indian populations. Excess suicide mortality is noted for the White and Indian populations; excess cirrhosis mortality is noted for Hispanics and Indians.

TABLE 3.20
NEW MEXICO STANDARDIZED MORTALITY RATIOS BY RACE/ETHNIC BASED ON FOUR YEAR RESIDENT MORTALITY 1979-1982 AGE/SEX ADJUSTED

RACE/ETHNIC	All Causes	Heart Disease	Malignant Neoplasms	ALL ACCIDENTS	Motor Vehicle	All Other	Stroke	COPD	Diabetes Mellitus	Suicide	Influenza & Pneumonia	Cirrhosis of Liver	Homicide	Alcoholism	Cert. Cond. in Perinatal Per.	Atherosclerosis	Congenital Anomalies
ALL PERSONS																	
OBSERVED DEATHS	35589	8437	7189	3558	2165	1393	2171	1655	959	998	968	678	691	607	558	408	325
EXPECTED DEATHS	37244	13652	7879	2323	1218	1105	2995	1008	627	593	905	618	535	89	675	494	373
SMR	96	62	91	153	178	126	72	164	153	168	107	110	129	682	83	83	87
% OF TOTAL DEATHS	100.0	23.7	20.2	10.0	6.1	3.9	6.1	4.7	2.7	2.8	2.7	1.9	1.9	1.7	1.6	1.1	0.9
NON-HISPANIC-WHITE																	
OBSERVED DEATHS	22389	6059	5036	1668	960	708	1511	1339	501	570	544	293	236	205	233	311	135
EXPECTED DEATHS	23709	9017	5120	1281	647	633	1991	658	413	340	503	391	291	55	274	331	163
SMR	94	67	98	130	148	112	76	203	122	168	93	75	81	373	85	94	83
% OF TOTAL DEATHS	100.0	27.1	22.5	7.5	4.3	3.2	6.7	6.0	2.3	2.5	2.4	1.3	1.1	0.9	1.0	1.4	0.6
HISPANIC-WHITE																	
OBSERVED DEATHS	10076	2025	1796	1180	747	433	548	283	367	326	321	299	336	243	236	88	137
EXPECTED DEATHS	10786	3740	2210	805	438	367	812	281	172	196	258	180	188	27	298	132	157
SMR	93	54	81	147	171	118	67	101	213	166	124	166	179	900	79	67	87
% OF TOTAL DEATHS	100.0	20.1	17.8	11.7	7.4	4.3	5.4	2.8	3.6	3.2	3.2	3.0	3.3	2.4	2.3	0.9	1.4
BLACK																	
OBSERVED DEATHS	563	114	119	44	19	25	29	12	19	10	10	6	35	19	20	5	6
EXPECTED DEATHS	528	179	109	43	24	19	38	14	8	11	12	9	10	1	15	6	8
SMR	107	64	109	102	79	132	76	86	238	91	83	67	350	1900	133	83	75
% OF TOTAL DEATHS	100.0	20.2	21.1	7.8	3.4	4.4	5.2	2.1	3.4	1.8	1.8	1.1	6.2	3.4	3.6	0.9	1.1
INDIAN																	
OBSERVED DEATHS	2507	224	226	663	437	226	81	19	68	91	92	80	81	140	64	4	47
EXPECTED DEATHS	1968	635	387	170	95	75	137	49	30	39	46	32	39	5	83	22	42
SMR	127	35	58	390	460	301	59	39	227	233	200	250	208	2800	77	18	112
% OF TOTAL DEATHS	100.0	8.9	9.0	26.4	17.4	9.0	3.2	0.8	2.7	3.6	3.7	3.2	3.2	5.6	2.6	0.2	1.9

Expected deaths have been rounded to whole numbers; the SMR's were computed before rounding and may differ somewhat from those that would be arrived at using the rounded figures.

An excess number of homicide deaths is noted for Blacks, and all four groups experience a mortality from alcoholism far beyond what would be expected based on national data.

An excess mortality (beyond what would be expected based on national data) is noted for New Mexicans in age groups between 1 and 44 years of age (Table 3.21).

Excessive mortality is noted for accidental death across all age groups in New Mexico. For age less than one year, accidents account for 3.5% of all deaths and death from motor vehicle accidents in particular occurs at a level three times the expected level. The leading categories of death in this age group: "certain conditions in the perinatal period" and "congenital anomalies" account for 45% and 21% of deaths respectively though the SMRs indicate a lower than expected level of mortality. In examining other causes such as COPD, the reader should cautiously interpret SMRs based on very low frequencies of events.

Accidents account for half of all deaths of New Mexicans 1-4 years of age, with motor vehicle mortality occurring at twice the expected rate.

For ages 5-14, accidents account for 55% of all deaths and though the SMRs are not as high as for the previous age group, motor vehicle accident mortality is 29% higher than expected and "all other" accidents is 14% higher.

With accidents accounting for 57% of deaths, suicide 16%, and homicide 11%, violent causes account for 84% of death in the 15-24 age group. Excessive death of New Mexicans is noted in these mortality categories as motor vehicle accident mortality is 56% higher than expected based on national data; "other accident" mortality is 34% higher, homicide is 20% higher and suicide is twice as high as expected.

The SMR for alcoholism is based on a low frequency of expected deaths, but even the fact that deaths from this cause are observed for this young age group is cause for concern.

Violent death accounts for 56% of all deaths of New Mexicans 25 to 44 years of age. Accident mortality, though accounting for a lower percentage of deaths than for the 15-24 year olds is more excessive in this age group. Motor vehicle accident mortality is twice what would be expected based on national data and "other accident" mortality is 41% higher. A 66% excess mortality is noted for suicide and 34% for homicide. Cancer ranks as third leading cause of death accounting for 11% of deaths in this age group, though at a level 16% below nationally. Alcoholism mortality in the State is 7½ times what would be expected based on national data, as one out of every twenty deaths in this age group in New Mexico is due to alcoholism.

TABLE 3.21
NEW MEXICO STANDARDIZED MORTALITY RATIOS BY RACE/ETHNIC
BASED ON FOUR YEAR RESIDENT MORTALITY
1979-1982 AGE/SEX ADJUSTED

AGE IN YEARS	All Causes	Heart Disease	Malignant Neoplasms	All Accidents	Motor Vehicle	All Other	Stroke	COPD	Diabetes Mellitus	Suicide	Influenza & Pneumonia	Cirrhosis of Liver	Homicide	Alcoholism	Cert. Cond. in Perinatal Per.	Atherosclerosis	Congenital Anomalies
[illegible]																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
1 - 4																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
5 - 14																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
15 - 24																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
25 - 44																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
[illegible]																	
OBSERVED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
EXPECTED DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
SMR	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]
% OF TOTAL DEATHS	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]	[illegible]

Expected deaths have been rounded to whole numbers. The SMR's were computed before rounding and may differ somewhat from those that would be arrived at using the rounded figures.

For New Mexicans age 45-64, cancer accounts for 29% of all deaths and heart disease for 22%. Though these two causes account for over half of all deaths in this age group, SMRs indicate a level of mortality below expected. By this age, accident mortality accounts for only 7% of all deaths but mortality for motor vehicle accidents is twice the expected level, and for "other accidents" it is 13% higher. Other excess mortality is indicated for COPD (42%), diabetes (54%), suicide (44%), homicide (25%), and death from alcoholism is 7½ times the expected level.

By age 65-74, heart disease and cancer account for 57% of deaths. Ranking third, COPD accounts for 8% of deaths as accident mortality has dropped to only 3% of the total. Excess mortality in this age group is noted for motor vehicle accidents (90%), COPD (59%), diabetes (48%), suicide (57%), homicide (59%), while observed deaths from alcoholism overshadow the level expected, based on national data, by a factor of almost six and a half.

One out of every three deaths occurring to New Mexicans 75 years of age and older is due to heart disease; one out of every six is due to cancer, and one out of ten is due to stroke. SMRs for heart disease, stroke, and atherosclerosis are below 100 (less deaths than expected). This age group along with the less than one age group are the only two for which "other accidents" is the major component of accidental death. Only 18% of accidental death for this age group is due to motor vehicle accidents though both components of accidental death are over forty percent higher than expected. Other notable excess mortality is noted for COPD, diabetes, cirrhosis, homicide and, as for younger age groups, alcoholism (Table 3.21).

YEARS OF POTENTIAL LIFE LOST

Years of potential life lost (YPLL) is a measure of premature mortality over the arbitrary span from the first birthday until completion of the 64th year of life.

This measure counts deaths at younger ages more heavily than those at older ages: the younger the person in general, the greater the number of potential years of life yet available to that individual.

The ranking of leading categories of underlying causes of death by contribution to premature mortality (YPLL) is presented in Figure 3.6 for the United States based on age-specific death rates for 1983 and for New Mexico based on age-specific mortality for the period 1979 through 1982.

Though accidents are the fourth leading cause of death in the nation and third leading cause of death in New Mexico, this category accounts for the greatest percentage of years of potential life lost in the State and nationwide (Figure 3.6); 24% nationally and 31% in New Mexico.) That is, the loss of one out of every four years of potential life lost nationally and one out of every three in the State results from accidents.

Cancer which ranks as the second leading cause of death, accounts for the second largest portion of YPLL—19.7% of YPLL nationally and 15.4% in New Mexico. In the State cancer accounts for about half the percentage of years of potential life lost as does accidents.

The leading cause of death in the nation—heart disease—accounts for 17% of YPLL nationally, ranking third; however in New Mexico heart disease ranks fourth accounting for 10.4% of YPLL (Figure 3.6).

Suicides/Homicides ranks third in New Mexico in the percentage of years of potential life lost, exceeding heart disease by 43% and almost equal to the YPLL attributed to cancer in the State.

The four categories—accidents, cancer, suicide/homicide, heart disease—account for 72% of all years of potential life lost in New Mexico and 74% nationally. The two violent categories account for 46% of YPLL in the State and 38% nationwide.

Of the remaining categories presented in Figure 3.6, the percentage of YPLL in New Mexico exceeds that for the nation for: chronic liver disease, chronic obstructive pulmonary disease (COPD), and diabetes.

FIGURE 3.6
YEARS OF POTENTIAL LIFE LOST
FOR PERSONS DYING IN THE 1-64 YEAR AGE GROUP
PERCENT CONTRIBUTION TO TOTAL YEARS BY CAUSE
UNITED STATES 1983
NEW MEXICO 1979-82



Breaking the suicide/homicide category into its two components shows that suicide accounts for 8.4% of YPLL in New Mexico and homicide 6.5%. Examination of an additional category for the state - alcoholism - shows that alcoholism accounts for 3.3% of years of potential life lost in New Mexico.

INFANT MORTALITY

As seen in Figure 3.7, the infant mortality rate in New Mexico has dropped appreciably since 1940, when it was over twice the national rate and ranked highest among the states.

FIGURE 3.7 INFANT MORTALITY RATE, NEW MEXICO RESIDENTS AND UNITED STATES 1940-1985



In 1980, for the first time ever, New Mexico's infant mortality rate dropped below the national rate—11.0 infant deaths per thousand live births in New Mexico compared to 12.5 nationally. A rate of 11.4 was reported for New Mexico in 1982 and 10.0 in 1983—the State's second lowest infant mortality rate.

By 1984, New Mexico's rate had dropped to 9.6 infant deaths per 1,000 live births—the lowest rate ever recorded in New Mexico. In 1985 infant mortality increased to a rate of 10.6.

Nationally, the rate decreased to 10.9 in 1983 and 10.6 infant deaths per thousand live births in 1984 and 1985, the lowest ever reported for the nation as a whole.

Only nineteen states reported a higher infant mortality rate in 1985 than did New Mexico. The States' infant mortality rates ranged from 6.7 (New Hampshire) to 14.0 in (South Carolina). For the District of Columbia, infant mortality was 22.1 infant deaths per thousand live births.

Infant mortality decreased for the Mountain Division in 1985, resulting in a 3% decrease from the previous year. The lowest rate was for the New England Division (Figure 3.8).

Increases in infant mortality rates in 1985 are noted for all of the State's major race/ethnic populations with the exception for the Indian which decreased 2% from 1984. The lowest rates for 1985 are noted for the Hispanic and Navajo Indian (Table 3.22).

FIGURE 3.8 UNITED STATES BY DIVISION INFANT MORTALITY RATE 1985



PROVISIONAL RATE PER 1,000 LIVE BIRTHS

TABLE 3.22 NEW MEXICO RESIDENTS INFANT MORTALITY RATE BY ETHNIC/RACE 1974-1985

NEW MEXICO	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985
ALL RACES	18.3	16.9	15.6	14.4	13.8	14.2	11.0	9.6	11.4	10.0	9.6	10.6
WHITE												
Non-Hispanic	20.7	15.4	16.5	15.5	14.4	13.9	10.8	9.7	11.8	10.9	9.6	11.8
Hispanic	16.9	16.7	14.1	12.1	12.8	12.3	10.5	9.1	11.0	8.8	9.5	9.6
BLACK	33.0	34.5	20.0	24.7	25.1	15.1	26.4	15.7	12.5	12.7	6.3	14.8
INDIAN	20.5	19.1	18.2	17.3	14.1	22.9	11.5	10.1	12.3	11.1	10.4	10.2
Pueblo	13.7	14.6	17.4	27.4	14.2	11.9	2.6	9.1	8.2	9.1	4.9	11.9
Navajo	24.5	21.8	19.4	18.1	14.8	29.2	14.9	10.5	14.6	12.2	12.0	10.2
Apache	16.7	16.1	22.0	10.4	8.2	0.0	13.3	13.4	6.1	10.0	11.4	16.8
ALL OTHER NON-WHITE	16.1	13.9	4.6	9.5	4.7	5.6	3.8	9.1	2.8	2.9	11.9	5.7

Rates Per 1,000 Live Births

Table 3.23 on the following page lists infant death rates by cause for the United States, 1984, and New Mexico by single years, 1981 through 1985.

Notable decreases by cause in the State are noted between 1981 and 1985 for low birth weight, and accidents. Respiratory distress syndrome decreased 28% the prior year.

"Conditions originating in the prenatal period" is the major component of infant mortality in the State; within this category "respiratory distress syndrome" and "short gestation and low birth weight" are the two largest causes of death. The second leading cause of infant mortality is "congenital anomalies" accounting for 27% of all deaths. The mortality rates presented in Table 3.23 are per 100,000 live births.

TABLE 3.23
INFANT MORTALITY RATES BY CAUSE
NEW MEXICO RESIDENTS 1981-1985
UNITED STATES 1984
(Rates Per 100,000 Live Births)

ICDA	CAUSE OF DEATH, 9TH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES	United States 1984	NEW MEXICO Single Year Data				
			1985	1984	1983	1982	1981
	ALL CAUSES	1078.7	1060.1	961.6	996.1	1143.7	963.7
036	MENINGOCOCCAL INFECTION	1.8	---	---	---	---	3.8
038	SEPTICEMIA	8.1	7.3	3.7	---	---	---
Residual of 001-139	OTHER INFECTIOUS & PARASITIC DISEASES	9.8	---	32.9	14.5	25.3	18.8
140-208	MALIGNANT NEOPLASMS	3.1	3.6	11.0	14.5	---	3.8
320-322	MENINGITIS	7.3	18.2	21.9	29.1	21.7	26.4
323-389	OTHER DISEASES OF NERVOUS SYSTEM & SENSE ORGANS	12.3	10.9	14.6	3.6	10.9	11.3
480-487	PNEUMONIA AND INFLUENZA	18.7	21.9	7.3	43.6	14.5	18.8
Residual of 460-519	OTHER DISEASES OF THE RESPIRATORY SYSTEM	17.3	---	3.7	7.3	18.1	22.6
520-579	DISEASES OF THE DIGESTIVE SYSTEM	11.8	18.2	3.7	10.9	7.2	18.8
740-759	CONGENITAL ANOMALIES	233.0	291.4	204.8	243.6	282.3	207.0
740	Anencephalus and similar anomalies	18.5	10.9	11.0	14.5	3.6	11.3
741	Spina bifida	2.6	3.6	---	7.3	10.9	---
742.3	Congenital Hydrocephalus	6.1	14.6	11.0	7.3	7.2	7.5
742.0-742.2, 742.4-742.9, 743	Other congenital anomalies of central nervous system and eye	8.4	18.2	7.3	10.9	18.1	18.8
745-746	Congenital anomalies of heart	69.7	94.7	54.8	87.2	123.1	75.3
747	Other congenital anomalies of circulatory system	22.4	7.3	18.3	7.3	7.2	11.3
748	Congenital anomalies of respiratory system	24.9	21.9	7.3	3.6	10.9	11.3
749-751	Congenital anomalies of digestive system	4.1	14.6	---	21.8	7.2	11.3
752-753	Congenital anomalies of genitourinary system	13.5	10.9	14.6	7.3	7.2	3.8
754-756	Congenital anomalies of musculoskeletal system	18.6	18.2	21.9	14.5	18.1	11.3
758.0	Down's syndrome	2.7	7.3	14.6	10.9	7.2	3.8
758.1-758.9	Other chromosomal anomalies	18.7	21.9	3.7	14.5	18.1	11.3
744, 757, 759	All other and unspecified congenital anomalies	22.9	47.4	40.2	36.4	43.4	30.1
760-779	CERTAIN CONDITIONS ORIGINATING IN THE PERINATAL PERIOD	509.2	419.0	471.7	374.4	416.2	402.8
760	Newborn affected by maternal conditions which may be unrelated to present pregnancy	3.8	7.3	14.6	3.6	---	---
761	Newborn affected by maternal complications of pregnancy	38.3	29.1	18.3	14.5	14.5	11.3
762	Newborn affected by complications of placenta, cord, and membranes	24.2	14.6	25.6	14.5	21.7	18.8
763	Newborn affected by other complications of labor and delivery	1.8	3.6	7.3	---	---	---
764	Slow fetal growth and fetal malnutrition	1.0	---	---	---	---	---
765	Disorders relating to short gestation and unspecified low birthweight	88.9	131.2	138.9	134.5	235.3	233.4
766	Disorders relating to long gestation and high birth weight	0.0	---	---	---	---	---
767	Birth trauma	10.2	3.6	---	14.5	7.2	3.8
768	Intrauterine hypoxia and birth asphyxia	31.9	25.5	25.6	21.8	10.9	22.6
768.2-768.4	Fetal distress in liveborn infant	6.6	10.9	7.3	3.6	3.6	11.3
768.5-768.9	Birth asphyxia	25.3	14.6	18.3	18.2	7.2	11.3
769	Respiratory distress syndrome	96.9	83.8	117.0	65.4	50.7	41.4
770	Other respiratory conditions of newborn	95.3	69.2	36.6	32.7	29.0	26.4
771	Infections specific to the perinatal period	24.2	21.9	25.6	18.2	10.9	15.1
772	Neonatal hemorrhage	10.9	---	7.3	7.3	7.2	3.8
773-774	Hemolytic disease of newborn, due to isoimmunization and other perinatal jaundice	1.5	---	---	3.6	3.6	---
775.0-775.1	Syndrome of "infant of a diabetic mother" and neonatal diabetes mellitus	0.2	---	---	---	---	---
776.0	Hemorrhagic disease of newborn	0.2	---	---	---	---	---
775.2-775.9 & 776.1-779	All other and ill-defined conditions originating in the perinatal period	79.8	29.1	54.8	43.6	25.3	26.4
780-799	SYMPTOMS, SIGNS AND ILL-DEFINED CONDITIONS	160.1	182.2	120.7	174.5	231.6	154.3
798.0	Sudden infant death syndrome	142.9	174.9	113.3	156.3	199.1	139.3
780-797, 798.1-799	Symptoms, signs and all other ill-defined conditions	17.1	7.3	7.3	18.2	32.6	15.1
E800-E949	ACCIDENTS AND ADVERSE EFFECTS	22.8	29.1	21.9	29.1	61.5	37.6
E960-E969	HOMICIDE	6.5	14.6	3.7	3.6	7.2	15.1
E967	Child battering and other maltreatment	2.1	14.6	3.7	3.6	7.2	11.3
E960-E966, E968-E969	Other homicide	4.3	---	---	---	---	3.8
Residual	ALL OTHER CAUSES	57.1	43.7	40.2	47.3	47.1	22.6

As seen in Figure 3.9, between 1973 and 1978, the State's neonatal mortality rate was below the national level, but an increase in the State's rate for 1979 placed New Mexico's rate 1% higher than nationally. In 1980, the State's rate dropped appreciably lower than the U.S. figure. For 1981, a further decrease in the rate left New Mexico's rate 27% lower than nationally (5.7 compared to 7.8). In 1982 the State's rate increased from the previous year to 6.7, but remained 12% lower than nationally. In 1983, the State's neonatal mortality rate decreased to its lowest level yet (5.5), 25% lower than nationally. In 1984, the State's neonatal rate increased 9% from the previous year but remained 12% lower than the national rate. In 1985 the State's rate increased 3% to 6.2, but remains 9% lower than the national rate of 6.8.

FIGURE 3.9
NEONATAL AND POSTNEONATAL MORTALITY RATES NEW MEXICO RESIDENTS AND UNITED STATES
1960-1985



In the postneonatal period, New Mexico's mortality rate has over the years consistently exceeded the national rate. In 1978, the State's rate was 21% higher than nationally; in 1979 the difference increased to 29%. In 1980, the State's rate dropped slightly below the U.S. rate (4.0, compared to 4.1 nationally).

In 1981, New Mexico's postneonatal mortality rate was 3.9 per thousand live births, the same as nationally. New Mexico's rate increased to 4.7 in 1982, dropping to 4.5 in 1983, and dropping further in 1984 to 3.6, the lowest postneonatal mortality rate reported for the State. In 1985 postneonatal mortality increased to a rate of 4.4, 19% higher than the 1985 national rate of 3.7.

Figure 3.10 graphically illustrates the trends and levels of infant mortality in New Mexico in recent years. The lowest neonatal mortality rate in 1985 was the rate recorded for New Mexico's Indian population. The 1985 neonatal mortality rate increased for three of the State's largest race/ethnic groups with the increase being smallest for the Indian, and decreased for Hispanic Whites.

The 1985 postneonatal mortality rate was highest for the State's Black population and lowest for the Hispanic White. The rate for the Indian was a decrease from the prior year's with other groups showing increases as high as 41% (non-Hispanic White).

FIGURE 3.10
NEONATAL AND POSTNEONATAL DEATH RATES BY ETHNIC/RACE, NEW MEXICO RESIDENTS
1978-1985



As was the case in the prior two years, infant mortality rates were higher for males than for females in New Mexico during 1985. The rate decreased over the previous year's rate for both males and females. The State's non-Hispanic White population recorded the highest male infant mortality rate. The highest infant mortality rate for females was noted for Hispanic females (Table 3.24).

TABLE 3.24
NEW MEXICO RESIDENTS, 1985
INFANT MORTALITY RATE BY ETHNIC/RACE AND SEX

(Rates Per 1,000 Live Births)

ETHNIC/RACE	MALE	FEMALE	TOTAL
ALL RACES:			
Infant Deaths	12.4	8.7	10.6
Neonatal Deaths	7.0	5.3	6.2
Postneonatal Deaths	5.4	3.3	4.4
NON-HISPANIC WHITE:			
Infant Deaths	15.2	8.1	11.8
Neonatal Deaths	8.6	5.3	7.0
Postneonatal Deaths	6.6	2.8	4.8
HISPANIC WHITE:			
Infant Deaths	9.7	9.5	9.6
Neonatal Deaths	5.6	5.7	5.7
Postneonatal Deaths	4.1	3.8	4.0
INDIAN:			
Infant Deaths	12.8	7.5	10.2
Neonatal Deaths	5.6	4.9	5.3
Postneonatal Deaths	7.2	2.7	5.0
BLACK:			
Infant Deaths	20.6	8.9	14.8
Neonatal Deaths	14.7	3.0	8.9
Postneonatal Deaths	5.9	6.0	5.9

FIGURE 3.11
NEW MEXICO RESIDENTS
INFANT, NEONATAL, POSTNEONATAL MORTALITY BY LEVEL OF PRENATAL CARE FOR THE 1981 TO 1983 BIRTH COHORT



The marked increase in infant mortality with decreasing levels of prenatal care is presented in Figure 3.11 based on the 1981-1983 birth cohort.

A five year aggregate infant mortality rate by county is presented in Table 3.25. Neonatal mortality rates are higher than postneonatal mortality rates in 26 of the State's counties.

The highest postneonatal mortality rates were recorded in Harding and Sierra counties.

TABLE 3.25
INFANT MORTALITY
NUMBER AND RATE BY COUNTY
NEW MEXICO RESIDENTS 1981-1985 AGGREGATE

HEALTH DISTRICTS	County	Total Infant Deaths		Neonatal		Postneonatal	
		Number	Rate	Number	Rate	Number	Rate
	NEW MEXICO	1,400	10.3	824	6.0	576	4.2
	Cibola *	20	8.7	12	5.2	8	3.5
1	Mckinley	98	11.6	46	5.5	52	6.2
	San Juan	104	8.9	55	4.7	49	4.2
	Colfax	8	6.8	6	5.1	2	1.7
	Harding	2	27.8	1	13.9	1	13.9
	Los Alamos	10	9.9	5	4.9	5	4.9
	Mora	5	12.9	3	7.8	2	5.2
2	Rio Arriba	46	13.2	28	8.1	18	5.2
	San Miguel	17	7.2	12	5.1	5	2.1
	Santa Fe	33	4.9	20	3.0	13	1.9
	Taos	17	7.6	12	5.4	5	2.2
	Union	3	7.1	3	7.1	--	--
	Bernalillo	420	10.9	267	7.0	153	4.0
3	Sandoval	44	10.6	25	6.0	19	4.6
	Torrance	7	10.4	5	7.5	2	3.0
	Valencia *	43	11.3	23	6.1	20	5.3
	Chaves	65	12.1	44	8.2	21	3.9
	Curry	45	9.8	20	4.3	25	5.4
	De Baca	1	5.2	1	5.2	--	--
	Eddy	55	11.1	35	7.1	20	4.0
4	Guadalupe	8	19.0	5	11.8	3	7.1
	Lea	97	13.0	52	7.0	45	6.0
	Lincoln	17	14.2	9	7.5	8	6.7
	Quay	6	6.8	2	2.3	4	4.5
	Roosevelt	17	11.6	11	7.5	6	4.1
	Catron	5	27.5	3	16.5	2	11.0
	Dona Ana	79	7.5	49	4.7	30	2.9
	Grant	25	9.6	14	5.4	11	4.2
5	Hidalgo	8	12.5	6	9.4	2	3.1
	Luna	12	8.0	8	5.3	4	[illegible].7
	Otero	57	10.5	28	5.2	29	5.4
	Sierra	14	26.9	10	19.2	4	7.7
	Socorro	12	9.4	4	3.1	8	6.3

Rates per 1,000 Live Births.
* Prior to 1982, Cibola County data were included with Valencia County.

Analysis of 1985 New Mexico infant mortality by cause is presented in Table 3.26.

In 1985, 42% of all infant deaths occurred in the postneonatal period; 33% at less than one day; 19% at 1-6 days and 7% at 7-27 days.

Disorders relating to short gestation and unspecified low birth weight accounted for 32% of the deaths for infants under one day of age in 1985.

For infants 1-6 days of age respiratory distress syndrome accounted for 20% of deaths in 1985.

Congenital anomalies was the leading cause of death of infants in the 7-27 days age bracket, accounting for 32% of these deaths.

TABLE 3.26
INFANT MORTALITY RATES BY CAUSE/AGE
NEW MEXICO RESIDENTS 1985
(Rates Per 100,000 Live Births)

		NEW MEXICO INFANT DEATH RATES				
ICDA	CAUSE OF DEATH, 9TH REVISION INTERNATIONAL CLASSIFICATION OF DISEASES	< 1 DAY	1-6 DAYS	7-27 DAYS	28 DAYS < 1 YEAR	TOTAL
	ALL CAUSES	353.4	196.7	69.2	440.8	1060.1
036	MENINGOCOCCAL INFECTION	---	---	---	---	---
038	SEPTICEMIA	---	3.6	---	3.6	7.3
Residual of 001-139	OTHER INFECTIOUS & PARASITIC DISEASES	---	---	---	---	---
140-208	MALIGNANT NEOPLASMS	---	---	3.6	---	3.6
320-322	MENINGITIS	---	---	---	18.2	18.2
323-389	OTHER DISEASES OF NERVOUS SYSTEM & SENSE ORGANS	---	3.6	---	7.3	10.9
480-487	PNEUMONIA AND INFLUENZA	---	---	---	21.9	21.9
Residual of 460-519	OTHER DISEASES OF THE RESPIRATORY SYSTEM	---	---	---	---	---
520-579	DISEASES OF THE DIGESTIVE SYSTEM	---	---	---	18.2	18.2
740-759	CONGENITAL ANOMALIES	87.4	83.8	21.9	98.4	291.4
740	Anencephalus and similar anomalies	7.3	3.6	---	---	10.9
741	Spina bifida	3.6	---	---	---	3.6
742.3	Congenital Hydrocephalus	3.6	---	---	10.9	14.6
742.0-742.2, 742.4-742.9, 743	Other congenital anomalies of central nervous system and eye	3.6	7.3	---	7.3	18.2
745-746	Congenital anomalies of heart	10.9	25.5	18.2	40.1	94.7
747	Other congenital anomalies of circulatory system	---	3.6	---	3.6	7.3
748	Congenital anomalies of respiratory system	14.6	---	---	7.3	21.9
749-751	Congenital anomalies of digestive system	---	14.6	---	---	14.6
752-753	Congenital anomalies of genitourinary system	10.9	---	---	---	10.9
754-756	Congenital anomalies of musculoskeletal system	10.9	7.3	---	---	18.2
758.0	Down's syndrome	3.6	---	---	3.6	7.3
758.1-758.9	Other chromosomal anomalies	---	18.2	---	3.6	21.9
744, 757, 759	All other and unspecified congenital anomalies	18.2	3.6	3.6	21.9	47.4
760-779	CERTAIN CONDITIONS ORIGINATING IN THE PERINATAL PERIOD	255.0	98.4	36.4	29.1	419.0
760	Newborn affected by maternal conditions which may be unrelated to present pregnancy	---	7.3	---	---	7.3
761	Newborn affected by maternal complications of pregnancy	29.1	---	---	---	29.1
762	Newborn affected by complications of placenta, cord, and membranes	10.9	3.6	---	---	14.6
763	Newborn affected by other complications of labor and delivery	3.6	---	---	---	3.6
764	Slow fetal growth and fetal malnutrition	---	---	---	---	---
765	Disorders relating to short gestation and unspecified low birthweight	112.9	7.3	3.6	7.3	131.2
766	Disorders relating to long gestation and high birth weight	---	---	---	---	---
767	Birth trauma	---	3.6	---	---	3.6
768	Intrauterine hypoxia and birth asphyxia	14.6	10.9	---	---	25.5
768.2-768.4	Fetal distress in liveborn infant	7.3	3.6	---	---	10.9
768.5-768.9	Birth asphyxia	7.3	7.3	---	---	14.6
769	Respiratory distress syndrome	36.4	40.1	3.6	3.6	83.8
770	Other respiratory conditions of newborn	29.1	18.2	7.3	14.6	69.2
771	Infections specific to the perinatal period	10.9	3.6	7.3	---	21.9
772	Neonatal hemorrhage	---	---	---	---	---
773-774	Hemolytic disease of newborn, due to isoimmunization and other perinatal jaundice	---	---	---	---	---
775.0-775.1	Syndrome of "infant of a diabetic mother" and neonatal diabetes mellitus	---	---	---	---	---
776.0	Hemorrhagic disease of newborn	---	---	---	---	---
775.2-775.9 & 776.1-779	All other and ill-defined conditions originating in the perinatal period	10.9	3.6	14.6	3.6	29.1
780-799	SYMPTOMS, SIGNS AND ILL-DEFINED CONDITIONS	---	---	3.6	178.5	182.2
798.0	Sudden infant death syndrome	---	---	3.6	171.2	174.9
780-797, 798.1-799	Symptoms, signs and all other ill-defined conditions	---	---	---	7.3	7.3
E800-E949	ACCIDENTS AND ADVERSE EFFECTS	---	3.6	---	25.5	29.1
E960-E969	HOMICIDE	---	---	---	14.6	14.6
E967	Child battering and other maltreatment	---	---	---	14.6	14.6
E960-E966, E968-E969	Other homicide	---	---	---	---	---
Residual	ALL OTHER CAUSES	10.9	3.6	3.6	25.5	43.7

Sudden infant death syndrome (SIDS) was the leading cause of death in the postneonatal period in 1985 (Table 3.26).

MATERNAL MORTALITY

Maternal mortality has decreased drastically over the years both in New Mexico and nationally.

FIGURE 3.12
MATERNAL MORTALITY RATE
NEW MEXICO RESIDENTS AND UNITED STATES
1960-1985



In 1985, the deaths of 3 New Mexico resident women were attributed to complications of pregnancy, child birth and puerperium, the number of deaths is two less than the total for the prior year.

In 1980, 3 deaths were reported for a rate of 11.3 maternal deaths per 100,000 live births and in 1981, two maternal deaths for a 7.5 rate, the lowest ever for the State. In 1982 and 1983 there were three maternal deaths reported for each year resulting in a constant rate of 10.9. The 1984 rate was 18.3 decreasing to 10.9 in 1985.

There were 350 maternal deaths reported nationally during 1985 for a rate of 9.3 per 100,000 live births; the rate decreased 19% from the 1984 rate.

The reader is cautioned about the high rates and sometimes drastic changes in rates resulting from the relatively small number of maternal deaths and births in New Mexico in recent years. Also, the reader is reminded that these figures do not represent the deaths of all pregnant women but rather only those deaths attributed to complications of pregnancy, childbirth and puerperium *(ICDA codes 630-676)* according to guidelines set forth by the National Center for Health Statistics.

TABLE 3.27
MATERNAL MORTALITY BY ETHNIC/RACE
NEW MEXICO RESIDENTS 1980-1985
UNITED STATES 1980-1985

	1985		1984		1983		1982		1981		1980	
ETHNIC/RACE	NO.	RATE	NO.	RATE	NO.	RATE	NO.	RATE	NO.	RATE	NO.	RATE
NEW MEXICO:												
ALL RACES	3	10.9	5	18.3	3	10.9	3	10.9	2	7.5	3	11.3
White	1	4.4	4	17.6	3	13.1	2	8.7	2	9.0	2	8.9
Hispanic	1	8.1	3	24.3	3	24.4	2	16.8	2	17.1	1	8.5
Non-Hispanic			1	9.7	-	--	-	--	-	--	1	9.4
Black	1	148.1	-	--	-	--	-	--	-	--	1	175.7
Indian	1	26.3	1	27.3	-	--	1	28.0	-	--	-	--
Other	-	--	-	--	-	--	-	--	-	--	-	--
UNITED STATES:												
ALL RACES	350*	9.3*	285	7.8	290	8.0	292	7.9	309	8.5	334	9.2
White	N/A		159	5.4	170	5.9	171	5.8	184	6.3	193	6.7
Other	N/A		126	16.9	120	16.3	121	16.4	125	17.3	141	19.8

Rate per 100,000 Live Births
* Provisional

The State's maternal rate continues to exceed the National rate—in 1985, a rate 17% higher than the rate of 9.3 nationally (Table 3.27).

The three resident maternal deaths in the State in 1985 occurred, one each, in Bernalillo, Rio Arriba, and Cibola counties. All three deaths resulted from post-partum hemorrhage.

The three maternal deaths in 1985 were of one White Hispanic woman, one Black woman, and one Indian woman.

Table 3.26 presents latest available U.S. (1984) maternal mortality by age group as well as New Mexico data for two time periods, 1981-83 and 1983-85.

TABLE 3.28
MATERNAL MORTALITY RATES BY AGE OF MOTHER
NEW MEXICO RESIDENTS 1981-1983, 1983-1985 AVERAGES
UNITED STATES 1984
(Rates per 100,000 Live Births)

AGE GROUP	UNITED STATES		NEW MEXICO			
	1984		1981-1983		1983-1985	
	NUMBER	RATE	NUMBER	RATE	NUMBER	RATE
< 15	1	10.0	0	--	0	--
15-19	29	6.2	1	7.1	1	7.6
20-24	49	4.3	2	7.1	2	7.1
25-29	81	6.9	2	8.7	3	12.8
30-34	76	11.5	2	16.6	3	23.7
35-39	39	19.9	0	-	2	48.0
40 +	10	35.8	1	147.3	0	-
TOTAL	285	7.8	8	9.8	11	13.4

FETAL MORTALITY

A fetal death is defined as a death in utero. New Mexico statute had, up through 1979, required the reporting of all fetal deaths of 20 weeks or more gestation, however, in 1980, the regulations were revised to base the requirement on weight rather than gestation. The law now requires the reporting of fetal deaths weighing 500 or more grams.

FIGURE 3.13
FETAL MORTALITY RATIO
NEW MEXICO AND UNITED STATES
1945-1985



*(1985 U.S. DATA NOT AVAILABLE AT PRINTING)

New Mexico's 1985 total of 157 fetal deaths resulted in a fetal death ratio of 5.7 fetal deaths per thousand live births. This represents a decrease from the 1984 figure of 186 fetal deaths and a ratio of 6.8.

New Mexico's fetal death ratios for 1984 and 1985 were lower than the 1984 national ratio of 8.5 fetal deaths per thousand live births. The State's fetal death ratio has consistently been lower than the national ratio (Figure 3.13). The lowest ratio nationwide is for the New England Division (Figure 3.14).

New Mexico's fetal death ratios by age of mother (Figure 3.15) are lower than nationally. The low ratio for New Mexico's under 15 age group is most likely a random fluctuation of the small numbers involved; the general trend is high rates for younger and older women.

Fetal death data for New Mexico counties are presented in Table 3.29 for a seven year period. Notable fluctuations in ratios are noted for some counties because of the small number of events involved.

FIGURE 3.14
UNITED STATES BY DIVISION
FETAL DEATH RATIOS*
1984



*RATIO PER 1,000 LIVE BIRTHS

FIGURE 3.15
FETAL DEATH RATIO BY AGE OF MOTHER
NEW MEXICO RESIDENTS 1983-1985 AGGREGATE
UNITED STATES 1984



The fetal death ratio for New Mexico decreased from 1984 to 1985 for Hispanic White and Indian, but increased for the non-Hispanic White (11%) as well as for Blacks (43%).

In 1985 thirty-five percent of the fetal deaths reported an unknown or unspecified cause of death. For specified causes, "Fetus affected by complications of placenta, cord and membranes" was the category most often cited as cause of death. (Table 3.30) on Page 60.

TABLE 3.29
FETAL DEATHS
NUMBER AND RATIO BY COUNTY
NEW MEXICO RESIDENTS 1979-1985

HEALTH DISTRICTS

		1985		1984		1983		1982		1981		1980		1979	
	COUNTY	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO
	NEW MEXICO	157	5.7	186	6.8	192	7.0	202	7.3	180	6.8	204	7.7	206	8.4
	CIBOLA	--	--	4	7.5	5	8.7	4	6.1	*	*	*	*	*	*
1	MCKINLEY	8	4.7	10	6.0	11	6.6	16	9.3	15	9.0	18	10.8	26	16.0
	SAN JUAN	14	6.3	7	3.1	16	6.9	15	6.0	10	4.1	16	6.9	17	9.1
	COLFAX	4	15.6	--	--	4	16.0	1	4.0	3	13.8	4	16.7	3	13.6
	HARDING	--	--	--	--	--	--	--	--	--	--	--	--	--	--
	LOS ALAMOS	3	14.7	1	5.0	4	19.4	1	5.1	--	--	2	10.0	2	10.4
	MORA	2	27.8	--	--	--	--	1	12.0	--	--	3	33.3	--	--
2	RIO ARRIBA	2	2.6	6	8.2	4	5.9	2	3.1	5	7.6	5	7.6	7	11.0
	SAN MIGUEL	4	8.6	2	4.0	4	8.3	3	6.7	4	8.5	3	6.0	5	10.9
	SANTA FE	6	4.1	8	5.6	19	13.9	16	13.1	9	7.0	9	7.1	7	5.7
	TAOS	1	2.2	2	4.5	1	2.2	1	2.2	3	7.2	4	9.6	8	20.1
	UNION	2	26.3	--	--	1	12.5	--	--	--	--	--	--	1	13.2
	BERNALILLO	36	4.5	61	7.8	52	6.8	60	7.8	50	6.8	62	8.1	43	6.0
	SANDOVAL	6	6.1	9	10.0	8	10.1	14	18.9	6	8.3	5	7.3	5	9.2
3	TORRANCE	1	7.4	1	7.4	1	6.8	2	15.9	1	7.9	2	15.4	1	9.2
	VALENCIA	4	6.4	3	4.8	2	3.4	7	11.3	10	7.6	13	9.0	15	11.2
	CHAVES	4	4.0	8	7.5	11	9.4	5	4.4	6	6.1	10	9.9	7	7.6
	CURRY	10	11.2	3	3.1	6	6.6	5	5.4	5	5.5	7	7.2	2	2.1
	DE BACA	--	--	--	--	1	23.8	3	75.0	--	--	--	--	1	23.8
	EDDY	10	11.3	8	8.6	10	9.[illegible]	8	6.9	10	10.4	11	11.2	11	12.8
4	GUADALUPE	--	--	--	--	--	--	1	11.2	1	11.8	--	--	1	13.9
	LEA	15	11.0	14	10.2	2	1.3	4	2.5	9	6.0	4	3.0	8	7.0
	LINCOLN	4	16.3	5	20.3	2	7.9	2	8.4	1	4.6	1	5.9	--	--
	QUAY	--	--	3	16.4	2	12.1	2	10.0	--	--	1	5.6	3	16.7
	ROOSEVELT	2	7.0	3	11.2	5	17.0	1	3.0	1	3.4	1	3.4	2	7.3
	CATRON	--	--	--	--	--	--	--	--	2	48.8	--	--	--	--
	DONA ANA	10	4.7	11	4.8	9	4.3	9	4.3	10	5.1	5	2.7	19	11.2
	GRANT	2	3.9	8	17.0	4	8.6	5	9.2	9	15.1	4	7.3	3	5.5
5	HIDALGO	--	--	--	--	--	--	--	--	--	--	--	--	1	7.7
	LUNA	1	3.3	4	13.7	2	6.8	--	--	3	10.2	2	6.8	3	10.1
	OTERO	5	4.5	4	3.7	6	5.2	8	7.5	4	4.1	9	9.7	4	4.2
	SIERRA	--	--	--	--	--	--	4	41.7	2	19.2	1	11.0	--	--
	SOCORRO	1	4.0	1	4.2	--	--	2	8.7	1	3.6	2	8.1	1	4.2

---

*Prior to 1982, Cibola data were included in Valencia County
Ratio per 1,000 Live Births.

TABLE 3.30
FETAL DEATHS
NUMBER AND RATIO BY CAUSE
AND ETHNIC/RACE
NEW MEXICO RESIDENTS 1985

CAUSE		TOTAL		NON-HISPANIC WHITE		HISPANIC WHITE		BLACK		INDIAN		OTHER UNSPECIFIED	
9th Revision ICDA #		NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO	NO.	RATIO
	ALL FETAL DEATHS	157	5.7	74	7.2	66	5.3	2	3.0	15	3.9	--	--
740-759	CONGENITAL ANOMALIES	21	0.8	7	0.7	12	1.0	--	--	2	0.5	--	--
760.0-760.9	FETUS AFFECTED BY MATERNAL CONDITIONS WHICH MAY BE UNRELATED TO PRESENT PREGNANCY	6	0.2	1	0.1	3	0.2	1	1.5	1	0.3	--	--
760.0	Maternal hypertensive disorders	3	0.1	1	0.1	1	0.1	1	1.5	--	--	--	--
760.1	Maternal renal and urinary tract diseases	1	0.0	--	--	--	--	--	--	1	0.3	--	--
760.2	MATERNAL INFECTION	1	0.0	--	--	1	0.1	--	--	--	--	--	--
760.3	Other chronic maternal circulatery and respiratory diseases	--	--	--	--	--	--	--	--	--	--	--	--
760.5	Maternal injury	1	0.0	--	--	1	0.1	--	--	--	--	--	--
761.0-761.9	FETUS AFFECTED BY MATERNAL COMPLICATIONS OF PREGNANCY	4	0.1	4	0.4	--	--	--	--	--	--	--	--
761.0	Incompetent cervix	2	0.1	2	0.2	--	--	--	--	--	--	--	--
761.1	Premature rupture of membranes	2	0.1	2	0.2	--	--	--	--	--	--	--	--
761.3	Polyhydramnics	--	--	--	--	--	--	--	--	--	--	--	--
761.5	Multiple pregnancy	--	--	--	--	--	--	--	--	--	--	--	--
761.6	Maternal deaths	--	--	--	--	--	--	--	--	--	--	--	--
761.8	Other	--	--	--	--	--	--	--	--	--	--	--	--
762	FETUS AFFECTED BY COMPLICATIONS OF PLACENTA, CORD AND MEMBRANES	56	2.0	33	3.2	21	1.7	--	--	2	0.5	--	--
763	FETUS AFFECTED BY OTHER COMPLICATIONS OF LABOR AND DELIVERY	--	--	--	--	--	--	--	--	--	--	--	--
764	SLOW FETAL GROWTH AND FETAL MALNUTRITION	--	--	--	--	--	--	--	--	--	--	--	--
765	DISORDERS RELATING TO SHORT GESTATION AND UNSPECIFIED LOW BIRTH WEIGHT	8	0.3	3	0.3	5	0.4	--	--	--	--	--	--
768	INTRAUTERINE HYPOXIA AND BIRTH ASPHYXIA	--	--	--	--	--	--	--	--	--	--	--	--
769	RESPIRATORY DISTRESS	--	--	--	--	--	--	--	--	--	--	--	--
770	OTHER RESPIRATORY CONDITIONS OF FETUS	--	--	--	--	--	--	--	--	--	--	--	--
771	INFECTIONS SPECIFIC TO THE PERINATAL PERIOD	1	0.0	--	--	--	--	--	--	1	0.3	--	--
772	FETAL HEMORRHAGE	1	0.0	1	0.1	--	--	--	--	--	--	--	--
773	HAEMOLYTIC DISEASE OF FETUS DUE TO ISOIMMUNIZATION	2	0.1	--	--	2	0.2	--	--	--	--	--	--
775	SYNDROME OF"INFANT OF A DIABETIC MOTHER"	1	0.0	--	--	1	0.1	--	--	--	--	--	--
776	HAEMATOLOGICAL DISORDERS OF FETUS AND NEWBORN	1	0.0	1	0.1	--	--	--	--	--	--	--	--
777	PERINATAL DISORDERS OF DIGESTIVE SYSTEM	--	--	--	--	--	--	--	--	--	--	--	--
778.0	HYDROPS FETALIS NOT DUE TO ISOIMMUNIZATION	1	0.0	--	--	1	0.1	--	--	--	--	--	--
779.9	CAUSE UNKNOWN/UNSNSPECIFIED	55	2.0	24	2.3	21	1.7	1	1.5	9	2.4	--	--

---

Ratio Per 1,000 Live Births
0.0 Equals less Than a Tenth of One Percent

TABLE 3.31
NEW MEXICO RESIDENTS
MORTALITY BY COUNTY
1985

HEALTH DISTRICTS		DEATHS		DEATH RATE PER 1,000 POP		INFANT MORTALITY						MATERNAL	FETAL	DEATHS
						NEONATAL		POSTNEONATAL		TOTAL INFANT DEATH				
	COUNTY	MALE	FEMALE	NO.	RATE	NO.	RATE	NO.	RATE	NO.	RATE	DEATHS	NO.	RATIO
	NEW MEXICO	5477	4160	9637	6.6	170	6.2	121	4.4	291	10.6	3	157	5.7
	CIBOLA	78	53	131	5.2	3	5.7	3	5.7	6	11.4	1	0	0.0
1	MCKINLEY	233	122	355	5.6	8	4.7	11	6.4	19	11.0	0	8	4.7
	SAN JUAN	251	176	427	4.7	11	4.9	8	3.6	19	8.5	0	14	6.3
	COLFAX	74	51	125	8.7	2	7.8	0	0.0	2	7.8	0	4	15.6
	HARDING	7	8	15	15.0	0	0.0	1	62.5	1	62.5	0	0	0.0
	LOS ALAMOS	48	24	72	3.9	1	4.9	3	14.7	4	19.6	0	3	14.7
	MORA	18	13	31	6.7	3	41.7	1	13.9	4	55.6	0	2	27.8
2	RIO ARRIBA	162	106	268	8.3	4	5.3	4	5.3	8	10.6	1	2	2.6
	SAN MIGUEL	111	76	187	7.5	3	6.4	1	2.1	4	8.6	0	4	8.6
	SANTA FE	257	242	499	5.9	4	2.8	2	1.4	6	4.1	0	6	4.1
	TAOS	94	57	151	6.9	4	8.7	1	2.2	5	10.9	0	1	2.2
	UNION	32	27	59	11.6	2	26.3	0	0.0	2	26.3	0	2	26.3
	BERNALILLO	1719	1319	3038	6.5	60	7.6	39	4.9	99	12.5	1	36	4.5
3	SANDOVAL	186	126	312	7.1	6	6.1	4	4.1	10	10.2	0	6	6.1
	TORRANCE	44	24	68	7.9	3	22.1	0	0.0	3	22.1	0	1	7.4
	VALENCIA	162	104	266	7.4	8	12.7	1	1.6	9	14.3	0	4	6.4
	CHAVES	290	244	534	9.5	6	5.9	4	4.0	10	9.9	0	4	4.0
	CURRY	134	146	280	6.6	3	3.4	9	10.1	12	13.4	0	10	11.2
	DE BACA	26	18	44	18.3	0	0.0	0	0.0	0	0.0	0	0	0.0
	EDDY	268	195	463	8.8	6	6.8	1	1.1	7	7.9	0	10	11.3
	GUADALUPE	24	12	36	8.2	1	12.8	1	12.8	2	25.6	0	0	0.0
4	LEA	210	188	398	6.1	7	5.1	6	4.4	13	9.5	0	15	11.0
	LINCOLN	69	35	104	7.5	0	0.0	0	0.0	0	0.0	0	4	16.3
	QUAY	64	51	115	9.8	0	0.0	0	0.0	0	0.0	0	0	0.0
	ROOSEVELT	65	50	115	7.0	2	7.0	2	7.0	4	14.0	0	2	7.0
	CATRON	16	10	26	9.3	2	58.8	0	0.0	2	58.8	0	0	0.0
	DONA ANA	303	282	585	4.9	8	3.7	7	3.3	15	7.0	0	10	4.7
5	GRANT	102	85	187	6.8	5	9.7	3	5.8	8	15.5	0	2	3.9
	HIDALGO	19	23	42	6.8	0	0.0	0	0.0	0	0.0	0	0	0.0
	LUNA	129	86	215	12.3	2	6.6	0	0.0	2	6.6	0	1	3.3
	OTERO	156	111	267	5.4	5	4.5	8	7.2	13	11.7	0	5	4.5
	SIERRA	90	69	159	16.7	1	10.2	1	10.2	2	20.4	0	0	0.0
	SOCORRO	36	27	63	4.6	0	0.0	0	0.0	0	0.0	0	1	4.0

Infant Mortality Rates and Fetal Death Ratios per 1,000 Live Births

COMMUNICABLE DISEASES

INTRODUCTION

The distribution of communicable disease in New Mexico is influenced by a blend of three cultures in a unique environment. Disease more commonly associated with certain racial groups. . . for example, diphtheria among the Navajo; shigella on Indian Reservations; disease, like brucellosis, related to ingestion of Mexican dairy products; and diseases of geography, like plague, endemic to New Mexico. . . these all color the overall disease picture.

Data presented are limited to those diseases declared to be notifiable according to the regulations governing the reporting in New Mexico.

VENEREAL DISEASE

Gonorrhea is the most frequently reported communicable disease in the United States. Between 1960 and 1975, the number of gonorrhea cases reported in the United States increased substantially, plateauing at about one million cases a year. Since 1975 the nation has experienced a downward trend in the number of cases.

In 1985, the number of provisional gonorrhea cases reported increased to 911,419, a 4% increase from the 878,556 cases reported in 1984. Gonorrhea rates also increased to 384.5 per 100,000 population, up 3% from 1984, nationally.

New Mexico has experienced a decreasing trend in recent years in the number of reported cases. Since peaking in 1976 at 7,137 cases reported, the number per year has decreased to about half as many cases.

FIGURE 4.1
GONORRHEA CASES REPORTED
NEW MEXICO OCCURRENCES
1966-1985



The number of reported cases in New Mexico dropped by 5% in 1980, 0.9% in 1981, 6% in 1982, and then dropped 19% in 1983, 5% in 1984 and another 6% in 1985.

The total cases of gonorrhea reported in the State were 4,402 cases in 1982, 3,553 cases in 1983, 3,372 in 1984 and 3,181 cases in 1985.

Comparison of case rates (number of cases per 100,000 population) for gonorrhea in New Mexico reflects the decreasing trend in the State which began in the mid-Seventies. In 1978, for the first time New Mexico's gonorrhea case rate was lower than nationally. The further decreases up through 1985 has left the State's rate 43% lower than the provisional 1985 national rate. This is a notable change from earlier years when New Mexico's rate was as much as 30% higher than the national rate.

FIGURE 4.2
GONORRHEA
CASES PER 100,000 POPULATION
NEW MEXICO RESIDENTS
AND UNITED STATES
1975-1985



Based on 1984 data by States, New Mexico's gonorrhea case rate ranked thirty-third highest among the states, down from twenty-ninth in 1983. Georgia reported the highest incidence rate among the states in 1984.

Thirty-four of the states reported decreases in incidence rate resulting in the overall decrease in the rate of incidence of gonorrhea nationally in 1984.

The decrease in the State's incidence rate in 1985 resulted from a decrease in incidence rate for age groups 25-29, and older (Table 4.1 on the following page).

The State's incidence rates are lower than national rates for all age groups based on comparison of 1984 U.S. data which is the latest available national data by age with 1985 New Mexico data.

The incidence of gonorrhea for New Mexicans age 0-14 is 46% lower than for the same age group nationally; for teens 15-19 it is 49% lower than for teens nationally; for persons in their early twenties it is 45% lower, late twenties, 45% lower; early thirties, 40% lower.

The only age group showing an increased incidence over 1976 rates in New Mexico is age 0-14. The greatest decrease has been for ages 25-29 which decreased 73% since 1976.

The number of reported cases of gonorrhea and syphilis case rates per 100,000 population for New Mexico's counties for 1985 are presented in Table 4.2. The case rates for 1985 ranged from a high of 579.8 for McKinley County to a low of 16.3 in Los Alamos County, with one county reporting no cases. The rate decreased in fifteen counties in 1985 resulting in an overall State decrease.

TABLE 4.1
NEW MEXICO & UNITED STATES
GONORRHEA
AGE SPECIFIC CASE RATES
NEW MEXICO 1976-1985, UNITED STATES 1976-1984

CASE RATES PER 100,000 POPULATION	ALL AGES	AGE GROUP 0-14	15-19	20-24	25-29	30-34	35+
NEW MEXICO:							
1985	215.0	10.5	582.8	882.1	470.8	255.7	54.6
1984	230.0	8.4	510.3	859.6	495.1	292.2	65.4
1983	251.2	10.7	516.0	993.5	563.6	319.0	64.0
1982	320.3	8.7	722.5	1237.8	751.1	401.7	68.5
1981	335.9	12.7	662.2	1422.7	720.5	488.9	66.3
1980	350.9	13.3	706.8	1337.6	899.1	505.9	75.1
1979	396.3	13.8	815.3	1611.1	1054.1	598.8	70.5
1978	462.0	13.3	939.2	1922.8	1251.1	602.5	91.9
1977	495.4	14.7	873.8	2112.6	1461.1	734.7	102.0
1976	611.0	7.6	982.6	2892.5	1775.3	860.9	141.1
UNITED STATES:							
1984	374.7	19.3	1133.7	1594.8	859.0	423.3	64.4
1983	387.6	19.0	1161.4	1616.6	881.0	406.6	69.6
1982	417.5	20.3	1198.9	1720.6	957.7	442.3	75.7
1981	435.2	21.1	1145.0	1749.2	1023.9	453.3	79.7
1980	443.3	55.4	1168.3	1780.5	1040.7	484.0	59.7
1979	459.4	24.3	1211.4	1910.0	1115.5	524.6	76.9
1978	468.3	11.1	977.6	2409.9	1648.0	833.8	127.0
1977	465.9	23.4	1212.4	2009.4	1153.5	523.7	76.5
1976	470.5	22.8	1253.6	2070.6	1136.7	535.5	74.1

United States Data Source:
VD Statistical Letter 1976-1984 Center for Disease Control
Military cases not included.

TABLE 4.2
NEW MEXICO
OCCURRENCE BY COUNTY
GONORRHEA AND SYPHILIS CASES REPORTED
1985

HEALTH DISTRICTS	COUNTY	GONORRHEA NUMBER	GONORRHEA RATE	SYPHILIS INFECTIOUS NUMBER	SYPHILIS INFECTIOUS RATE	SYPHILIS NON-INFECTIOUS NUMBER	SYPHILIS NON-INFECTIOUS RATE
	NEW MEXICO	3,181	219.4	237	16.3	89	6.1
	CIBOLA	14	55.3	4	15.8	-	--
1	MCKINLEY	367	579.8	71	112.2	6	9.5
	SAN JUAN	349	383.1	15	16.5	10	11.0
	COLFAX	10	69.9	-	--	3	21.0
	HARDING	1	100.0	-	--	-	--
	LOS ALAMOS	3	16.3	-	--	-	--
	MORA	13	282.6	2	43.5	1	21.7
2	RIO ARRIBA	81	250.0	3	9.3	4	12.3
	SAN MIGUEL	91	366.9	14	56.5	1	4.0
	SANTA FE	156	184.4	3	3.5	6	7.1
	TAOS	54	246.6	-	--	2	9.1
	UNION	16	313.7	-	--	-	--
	BERNALILLO	1,080	232.6	67	14.4	25	5.4
	SANDOVAL	33	74.8	8	18.1	4	9.1
3	TORRANCE	9	104.7	-	--	-	--
	VALENCIA	24	66.9	2	5.6	4	11.1
	CHAVES	127	224.8	4	7.1	-	--
	CURRY	144	338.0	6	14.1	-	--
	DE BACA	1	41.7	-	--	1	41.7
	EDDY	62	118.1	-	--	-	--
4	GUADALUPE	9	204.5	2	45.5	-	--
	LEA	189	288.5	3	4.6	5	7.6
	LINCOLN	9	64.7	-	--	1	7.2
	QUAY	19	162.4	-	--	1	8.5
	ROOSEVELT	46	278.8	-	--	1	6.1
	CATRON	-	--	-	--	3	107.1
	DONA ANA	180	152.2	29	24.5	8	6.8
	GRANT	16	58.6	-	--	1	3.7
5	HIDALGO	6	96.8	-	--	-	--
	LUNA	16	91.4	-	--	2	11.4
	OTERO	46	93.1	3	6.1	-	--
	SIERRA	3	31.6	1	10.5	-	--
	SOCORRO	7	51.5	-	--	-	--

Rate Per 100,000 Population
Includes both Civilian and Military Population.

The incidence rate for primary and secondary syphilis decreased in 1984 both nationally and in New Mexico. Nationally the rate decreased by 15% while the State's incidence rate decreased 49% in 1984 (Table 4.3), with national rates exceeding State rates for all ages. 1985 data shows the State rate increasing by 31% to 8.8.

TABLE 4.3
NEW MEXICO & UNITED STATES
PRIMARY AND SECONDARY SYPHILIS
AGE SPECIFIC CASE RATES
NEW MEXICO 1976-1985, UNITED STATES 1976-1984

CASE RATES PER 100,000 POPULATION	ALL AGES	AGE GROUP 0-14	15-19	20-24	25-29	30-34	35+
NEW MEXICO:							
1985	8.8	.3	15.3	14.3	18.1	20.1	7.2
1984	6.7	--	6.9	24.0	19.0	12.6	2.8
1983	13.2	.3	9.2	40.8	37.1	21.1	8.4
1982	14.6	.3	12.2	41.0	37.2	29.2	10.1
1981	10.4	.9	12.6	28.1	22.1	17.4	7.8
1980	8.6	--	8.3	20.7	27.8	18.8	4.9
1979	7.8	.0	11.5	25.6	12.2	19.0	4.7
1978	7.3	.3	10.8	23.7	19.2	17.7	2.9
1977	8.5	.3	5.4	33.3	24.4	10.7	5.7
1976	12.4	.6	11.6	47.2	38.2	21.7	6.0
UNITED STATES:							
1984	12.0	.3	17.3	39.1	33.1	23.0	5.6
1983	14.1	.4	23.2	43.7	36.6	25.8	6.3
1982	14.6	.5	23.0	44.8	38.4	27.0	6.6
1981	13.7	.4	19.6	41.1	37.1	25.1	6.3
1980	12.0	.9	16.9	35.8	32.6	23.2	4.1
1979	11.4	.3	16.2	35.3	31.6	22.4	3.4
1978	10.0	.3	14.6	30.8	29.0	19.5	4.6
1977	9.5	.3	13.8	30.3	26.6	18.4	4.5
1976	11.1	.4	17.3	35.1	31.2	22.9	5.2

Source: VD Statistical Letter 1976-1984 Center for Disease Control. Military cases not included.

A breakdown of cases of infectious and non-infectious syphilis also illustrates the increase in reported cases in New Mexico (Figure 4.3) over the prior year.

The number of reported non-infectious cases increased 31% in 1985 and the number of infectious increased 25%. The number of reported syphilis cases and case rates by county are presented in Table 4.2.

FIGURE 4.3
SYPHILIS CASES REPORTED
NEW MEXICO OCCURRENCES
1968-1985



PLAGUE

A variety of factors—including climatic conditions, rodent and flea population dynamics, and steady human encroachment into previously unsettled areas—are probably responsible for the increased number of plague cases during recent years.

FIGURE 4.4
PLAGUE
TOTAL CASES REPORTED BY YEAR
NEW MEXICO OCCURRENCES
1970-1985



FIGURE 4.5
PLAGUE
TOTAL CASES REPORTED BY COUNTY
NEW MEXICO OCCURRENCES
1985



Of the seventeen cases of plague reported in the nation for 1985, fourteen cases (82%) occurred in New Mexico.

The fourteen cases for New Mexico in 1985 follow 16 cases in 1984 and 26 in 1983. Though decreasing, as seen in Figure 4.4 these have been three consecutive years of higher than average plague case incidence.

The map of Figure 4.5 depicts cases reported for 1985 by county. Three cases occurred in Bernalillo County, two each in McKinley and Los Alamos counties, and one each in Taos, Colfax, Sandoval, Santa Fe, San Miguel, Mora and Grant counties.

Of plague cases reported in 1985, three cases were in the 0-9 age group, three were in the 10-15 age group, three in the 20-29 age group and 5 in the >30 age group.

The case ratio in 1985 was almost four males to every female, with a total of two fatal cases for the year.

TABLE 4.4
PLAGUE
NEW MEXICO OCCURRENCE
1977-1985

YEAR	1977	1978	1979	1980	1981	1982	1983	1984	1985
Total Cases	8	5	4	13	6	9	26	16	14
Month Acquired:									
January	-	-	-	-	-	-	-	-	-
February	-	-	-	-	1	-	-	-	-
March	-	-	-	-	-	-	-	1	-
April	-	-	1	-	1	-	2	1	1
May	2	-	-	1	1	2	2	2	3
June	-	1	2	3	-	-	8	2	3
July	2	1	1	2	1	4	8	2	3
August	1	2	-	5	2	2	4	3	1
September	2	-	-	2	-	-	1	2	-
October	1	1	-	-	-	1	1	2	3
November	-	-	-	-	-	-	-	1	-
All Ages:	8	5	4	13	6	9	26	16	14
Male	6	2	1	5	3	6	13	12	11
Female	2	3	3	8	3	3	13	4	3
0 - 9	3	1	-	6	-	1	5	-	3
Male	3	-	-	3	-	1	2	-	2
Female	-	1	-	3	-	-	3	-	1
10-15	-	2	1	1	-	2	6	5	3
Male	-	1	-	1	-	2	2	3	3
Female	-	1	1	-	-	-	4	2	-
16-19	-	-	1	1	-	-	1	1	-
Male	-	-	1	-	-	-	1	1	-
Female	-	-	-	1	-	-	-	-	-
20-29	-	-	-	-	4	3	3	1	3
Male	-	-	-	-	3	1	2	1	2
Female	-	-	-	-	1	2	1	-	1
> 30	5	2	2	5	2	3	11	9	5
Male	3	1	-	1	-	2	6	7	4
Female	2	1	2	4	2	1	5	2	1
FATAL CASES	-	-	2	3	3	2	2	2	2

RABIES

The number of rabies cases reported in New Mexico in 1979 was the second highest since the rabies epizootic of 1974. Since 1979 the number of rabies cases has decreased, with a total of 12 cases reported in 1984 and 1985.

Of the 12 cases reported in New Mexico in 1985, there were 2 skunk, 9 bat and 1 cat. There were no dog rabies reported in 1985 or the prior two years.

FIGURE 4.6
RABIES IN ANIMALS
CASES REPORTED
NEW MEXICO OCCURRENCES
1969-1985



There were 5,606 reported cases of rabies in the United States and Puerto Rico in 1985, an increase of 39 cases compared with reported cases in 1984.

Of the cases of rabies reported in 1985, in New Mexico, the counties in the southern part of the state accounted for 58%: 3 in Dona Ana, and 1 each in Eddy and Luna, and 2 in Otero County.

Dona Ana and Bernalillo counties reported the highest number of rabies in the State during 1985 (Figure 4.7). Santa Fe and Los Alamos counties were the other counties reporting one case each.

FIGURE 4.7
RABIES
NUMBER OF CASES REPORTED
NEW MEXICO OCCURRENCES
1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
MORA
HARDING
LOS ALAMOS
SANTA FE
SANDOVAL
MCKINLEY
SAN MIGUEL
BERNALILLO
VALENCIA
TORRANCE
GUADALUPE
QUAY
CURRY
DE BACA
ROOSEVELT
CATRON
SOCORRO
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
3
1
2
0

DIPHTHERIA

In the last eight years not a single case of diphtheria has been reported in New Mexico (Figure 4.8). The last major outbreak of diphtheria in the State occurred in 1973.

FIGURE 4.8
DIPHTHERIA
CASES REPORTED
NEW MEXICO OCCURRENCES
1969-1985



TUBERCULOSIS

In New Mexico the number of tuberculosis cases decreased by 18 from 112 cases in 1984 to 94 cases in 1985. Tuberculosis cases have been on a downward trend since 1981 when the number of reported cases was 152; subsequently dropping 38% to the 1985 level (Figure 4.9).

FIGURE 4.9
TUBERCULOSIS
CASES REPORTED
NEW MEXICO OCCURRENCES
1970-1985



The State's incidence rate for tuberculosis decreased slightly from 7.9 cases per 100,000 population in 1984 to 6.5 in 1985 (Table 4.5). New Mexico's case rate of 6.5 in 1985 was 31% lower than the 1984 national incidence rate of 9.4, (Table 4.7).

Table 4.5 illustrates the marked decreasing trend in New Mexico over the years for both tuberculosis incidence and mortality.

TABLE 4.5
NEW MEXICO
TUBERCULOSIS MORBIDITY & MORTALITY
1930-1985

YEAR	INCIDENCE	MORTALITY
1930	*	169.6
1940	*	71.8
1950	*	37.0
1960	59.3	7.8
1970	19.9	2.7
1975	12.7	2.1
1976	15.5	2.7
1977	12.8	2.5
1978	12.3	1.8
1979	12.3	1.3
1980	11.2	1.1
1981	11.4	1.1
1982	8.9	0.8
1983	8.3	1.1
1984	7.9	0.8
1985	6.5	1.0

RATE PER 100,000 POPULATION

*Due to changes in reporting over the years, these figures would not be directly comparable to figures for more recent years.

In New Mexico, the incidence of tuberculosis is notably higher for the State's Indian population than for the rest of the State's population. Though the incidence rate for the Indian has decreased in recent years, it remains over four times higher than for the non-Indian population of the State. In 1985 the State experienced a 29% decrease in the tuberculosis incidence rate for the Indian and 12% for the non-Indian population (Figure 4.10).

FIGURE 4.10
TUBERCULOSIS INCIDENCE RATES
NEW MEXICO OCCURRENCES
1976-1985



Mortality from tuberculosis in New Mexico has averaged over the last five years about one death per hundred thousand population. The rate for the Indian is three times higher than the overall State rate. Because of the relatively few number of tuberculosis deaths in the State in recent years the yearly fluctuation in the rates by race/ethnicity is not unexpected.

TABLE 4.6
NEW MEXICO RESIDENTS
TUBERCULOSIS DEATHS
RATES BY RACE/ETHNIC
1978-1985

ETHNIC/RACE	Tuberculosis Mortality Rate							
	1978	1979	1980	1981	1982	1983	1984	1985
Total	1.8	1.3	1.1	1.1	0.8	1.1	0.8	1.0
White								
Non-Hispanic	1.1	0.9	1.0	0.6	0.1	0.7	0.8	0.7
Hispanic	1.7	1.3	0.4	1.0	1.2	1.0	0.4	0.8
Black	0	0	0	0.4	0	0	0	3.7
Indian	7.4	4.2	4.7	3.7	3.6	4.4	3.4	3.4

Rate Per 100,000 Population

Roosevelt County showed the highest incidence rate (cases per hundred thousand population) in the State for 1985 (30.3) followed by McKinley County (22.1). Tuberculosis cases were reported for 18 of the State's 33 counties (Figure 4.11).

FIGURE 4.11
TUBERCULOSIS
NUMBER OF CASES REPORTED
Per 100,000 Population
New Mexico Occurrence
1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
MORA
HARDING
McKINLEY
LOS ALAMOS
SANDOVAL
SANTA FE
SAN MIGUEL
QUAY
BERNALILLO
CIBOLA
VALENCIA
GUADALUPE
TORRANCE
CURRY
DE BACA
ROOSEVELT
CATRON
SOCORRO
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
LUNA
DONA ANA
EDDY
HIDALGO
6.5 STATE AVERAGE
22.1-30.3
1.5-4.7
11.9-16.5
0
5.6-9.3

TABLE 4.7
TUBERCULOSIS
CASE RATE PER 100,000 POPULATION
SPECIFIC AGE GROUPS
NEW MEXICO OCCURRENCES 1983-1985
UNITED STATES 1982-1984

AGE	UNITED STATES			NEW MEXICO		
	1982	1983	1984	1983	1984	1985
TOTAL	11.0	10.2	9.4	8.3	7.9	6.5
0 - 4	4.6	4.6	4.3	2.4	6.0	5.0
5 - 14	1.7	1.6	1.4	0.4	1.2	1.7
15 - 24	4.7	4.7	4.2	1.8	1.8	4.4
25 - 44	10.2	9.6	9.0	9.8	5.8	3.0
45 - 64	16.8	15.4	14.4	10.6	8.8	9.2
65 +	27.9	25.6	23.2	34.5	37.8	25.2

1984 Tuberculosis Statistics, CDC
Rates Per 100,000 Population

The 18% decrease in the statewide incidence rate for tuberculosis in 1985 was the net result of increases for the 5-14, 15-24, and 45-64 age groups and decreases in rate for the 0-4, 25-44 and 65+ age groups. New Mexico's age specific rates are lower than nationally except for the 25-64 age group; this is based on comparison of 1985 New Mexico data to latest available (1984) national data.

RUBELLA

As seen in Figure 4.12, the level of rubella reported in New Mexico has remained drastically reduced from those levels prior to 1975, as in 1983 for the first time there were no cases of rubella reported in the State, only one case reported in 1984, and [illegible] 2 cases reported in 1985.

Rubella, like measles, has responded to intensive vaccination efforts and enforcement of school immunization laws.

FIGURE 4.12
RUBELLA
CASES REPORTED
NEW MEXICO OCCURRENCES
1969-1985



AIDS

By the end of 1985, physicians and health departments in the United States and Puerto Rico had reported a total of 16,533 cases of acquired immunodeficiency syndrome (AIDS). Of these cases, 56 cases were diagnosed before 1981, 247 in 1981, 965 in 1982, 2,473 in 1983, 4,445 in 1984, and 8347 in 1985.

There were three cases of AIDS reported in New Mexico in 1984, and 14 cases in 1985, bringing the State total since 1979 to eighteen cases.

MEASLES

The last major outbreak of measles in the State occurred in 1977 and centered in Valencia and Torrance counties. The incidence rate for measles in New Mexico then dropped from 3.1 cases per 100,000 population in 1979 to zero cases in 1982 and 1983 with some fluctuation in between including a "minor" outbreak in 1979 in Albuquerque and Silver City.

In 1984 the number of measles cases reached the second highest total in the last eleven years and then dropped to only six cases in 1985.

FIGURE 4.13
MEASLES
CASES REPORTED
NEW MEXICO OCCURRENCES
1970-1985



Of the 6 cases reported in 1985 statewide, 2 were reported in Valencia County, 2 in Chaves County and one each in Sandoval and Dona Ana counties.

The Lea County outbreak in 1984 occurred in Hobbs beginning in one junior high school and spreading through a final total of seven schools. 62% of the persons attended grades 7-9 while 19% attended grades 10-12; in all, persons affected were from the first, second and fifth through twelfth grades.

The incidence rate for measles (cases per hundred thousand population) in 1984 was almost six times higher for New Mexico than nationally. The rate for 1985 was significantly lower than the previous year, 0.4 as compared to 6.2 in 1984.

FIGURE 4.14
MEASLES
NUMBER OF CASES REPORTED
PER 100,000 POPULATION
NEW MEXICO OCCURRENCES
1985


SAN JUAN
RIO ARRIBA
TAOS
COLFAX
UNION
LOS ALAMOS
SANDOVAL
SANTA FE
MORA
HARDING
McKINLEY
SAN MIGUEL
BERNALILLO
CIBOLA
VALENCIA
GUADALUPE
QUAY
TORRANCE
CURRY
DE BACA
CATRON
SOCORRO
ROOSEVELT
LINCOLN
CHAVES
SIERRA
GRANT
OTERO
LEA
DONA ANA
LUNA
EDDY
HIDALGO
NEW MEXICO 0.4
5.6
2.3
3.5
0.8
0

OTHER COMMUNICABLE DISEASES

The number of cases of notifiable diseases and the corresponding cases rates are presented for New Mexico and the United States in Table 4.8 on the following page. The data are based on cases of notifiable diseases reported for New Mexico for 1985, and for the United States for 1984. New Mexico's case rates exceed national rates for psittacosis, plague, salmonellosis, shigellosis, viral hepatitis (A, B, and unspecified), legionellosis, malaria, tularemia, and typhoid.

Nationally, in 1983, the number of salmonella isolates reported reached the highest level ever and then decreased in 1984. New Mexico also reported a decrease in cases in 1984 (20%) but rose in 1985 to near the 1983 figure (406 cases and a rate of 28.0, in 1985).

In 1985, 400 cases of shigellosis were reported, 12% higher than the previous year, but still remaining three times higher than the 1984 national rate.

New Mexico's case rate for hepatitis B is 92% higher than the national rate, and the number of cases in the State has been increasing. Hepatitis A has decreased markedly in the State since 1979; the number of cases did increase in 1985 over 1984 pushing the State case rate to over three times the 1984 national level.

New Mexico has the highest plague case rate in the nation; 100 times the national rate.

In Table 4.9, the number of cases reported for selected communicable disease in New Mexico are presented for years 1968 to 1985.

Over the years presented, major annual rabies activity has occurred in 1968, 1974, 1975, 1979 and 1980, with the 1984 and 1985 total being the lowest case total since 1973. The last major diphtheria outbreak was in 1973, with no cases reported the last eight years. Measles and rubella have shown dramatic decreases in total cases compared to the early Seventies though experiencing outbreaks on occasion; for example, the increase in measles cases in 1984. A major hepatitis outbreak occurred in 1979, followed by general decreases, reaching in 1983 the lowest level since 1970, then increasing to almost double the 1985 figure.

Major pertussis activity occurred in 1975 and 1977, decreasing to eight cases in 1982 then increasing to 13 cases in 1983 and 15 for 1985. Reported plague cases reached an all time high total of 26 cases in 1983, then dropped to its second highest total (16) equal to the 1975 total and then dropped to 14 cases in 1985.

Salmonella cases, which reached the highest level ever reported in 1983, dropped in 1984 to its 1981 level and then increased in 1985 back to near the 1983 level. Shigellosis cases peaked in 1980 and has fluctuated between 351 and 432 cases since then.

Tuberculosis activity has decreased over the last four years. Three typhoid cases have been reported in 1983 and 1984 and 4 in 1985. Gonorrhea cases have been decreasing since 1976. Infectious syphilis cases which had increased notably in 1982 and 1983 dropped sharply in 1984, and increased for 1985, while non-infectious syphilis has fluctuated between 52 and 100 cases in the Eighties.

TABLE 4.8
NOTIFIABLE DISEASES REPORTED CASES AND CASE RATES NEW MEXICO 1985 UNITED STATES 1984

NOTIFIABLE DISEASES	NEW MEXICO		UNITED STATES	
	Cases	Rate	Cases	Rate
Aids	14	1.0	4,445	1.9
Amebiasis	9	0.6	5,252	2.2
Anthrax	-	-	1	0.00
Aseptic Meningitis	40	2.8	8,326	3.5
Botulism	-	-	123	0.1
Brucellosis	1	0.1	131	0.1
Chancroid	-	-	665	0.3
Cholera	-	-	1	0.00
Diphtheria	-	-	1	0.00
Encephalitis:				
Other Primary Infection	5	0.3	1,117	0.5
Arthropod-borne	-	-	129	0.1
Post-Infectious	-	-	119	0.1
Legionellosis (Legionnaires Disease)	7	0.5	750	0.3
Leprosy	-	-	290	0.1
Leptospirosis	-	-	40	0.01
Malaria	13	0.9	1,007	0.4
Measles	6	0.4	2,587	1.1
Meningococcal Infections	-	-	2,746	1.2
Pertussis	15	1.0	2,276	1.0
Plague	14	1.0	31	0.01
Poliomyelitis:				
Total	-	-	8	0.03
Paralytic	-	-	8	0.03
Psittacosis	6	0.4	172	0.1
Rubella	2	0.1	752	0.3
Congenital Syndrome	-	-	5	0.02
Salmonellosis (Excluding Typhoid Fever)	406	28.0	40,861	17.3
Shigellosis	400	27.6	17,371	7.4
Tetanus	-	-	74	0.03
Trichinosis	-	-	68	0.02
Tularemia	6	0.4	291	0.1
Typhoid Fever	4	0.3	390	0.2
Typhus Fever				
Flea-Borne, Endemic, Murine	-	-	53	0.02
Typhus Fever	-	-		
Tick Borne, Rky. Mt. Spotted)	-	-	838	0.4
Viral Hepatitis				
A	458	31.6	22,040	9.3
B	309	21.3	26,115	11.1
Unspecified	61	4.2	9,402	4.0

SOURCE: MMWR, Vol. 34, No. 38
Rates Per 100,000 Population

TABLE 4.9
SELECTED COMMUNICABLE DISEASES NUMBER OF REPORTED CASES NEW MEXICO OCCURRENCES 1968-1985

	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985
Animal Rabies:	40	20	16	9	10	7	78	43	22	21	27	49	45	28	23	15	12	12
Dog	6	1	0	0	3	0	31	5	0	0	0	8	2	0	2	0	0	0
Cat	4	3	0	0	0	1	6	3	0	0	2	1	1	0	0	1	1	1
Bat	7	5	6	1	4	5	30	19	7	10	10	12	17	18	13	7	2	9
Skunk	18	11	8	7	2	1	11	13	10	9	14	27	25	10	8	7	8	2
Other Species	5	0	2	1	1	0	0	3	5	2	1	1	0	0	0	0	1	0
Diphtheria	3	1	1	8	12	42	17	9	3	5	0	0	0	0	0	0	0	0
Hepatitis	290	320	369	605	522	519	755	554	449	771	663	1149	440	419	581	405	486	828
Measles	176	284	306	364	129	115	63	17	17	257	0	38	12	9	0	0	88	6
Pertussis	27	12	13	3	4	14	21	109	18	76	42	30	29	16	8	13	13	15
Plague	0	5	9	1	0	1	7	16	9	8	5	4	13	6	9	26	16	14
Rubella	136	321	237	251	128	174	130	19	4	11	4	12	5	5	6	0	1	2
Salmonella, Non-Typhoidal	96	87	102	88	94	154	104	96	83	116	265	184	212	327	259	409	327	406
Shigellosis	236	109	208	259	261	224	250	361	323	345	551	622	627	419	351	432	353	400
Tuberculosis	193	199	202	141	208	222	195	146	181	152	149	153	146	152	122	116	112	94
Typhoid	9	11	6	3	2	9	5	1	2	0	2	4	3	0	0	3	3	4
Gonorrhea	1876	2306	2882	2969	4377	4805	5433	7023	7137	5941	5600	4918	4676	4636	4402	3553	3372	3181
Syphilis																		
Infectious	171	263	128	162	223	183	184	284	286	180	145	159	218	248	321	350	189	237
Non-infectious	566	712	430	236	253	344	328	223	207	182	133	74	52	100	62	79	68	89

SPECIAL FEATURE

NEW MEXICO COUNTY HEALTH PROFILES

BERNALILLO COUNTY

BERNALILLO COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	315774	419700	464300	588500
% 19 AND YOUNGER	42.1%	33.2%	30.1%	28.3%
% 65 AND OLDER	6.1%	8.1%	9.2%	11.8%
% WHITE NON-HISPANIC	50.9%	56.9%	XXXXX	XXXXX
% WHITE HISPANIC	39.2%	36.4%	XXXXX	XXXXX
% BLACK	1.9%	2.3%	XXXXX	XXXXX
% INDIAN	7.2%	2.7%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	13.0%	10.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	31.1%	32.2%	32.0%	31.8%

•Almost one-third of the state's population resides in Bernalillo County making it New Mexico's most populous county with a population density of 397.2 persons per square mile.

•The county population is projected to approach 600,000 by the year 2000 as the percent of population 19 and younger decreases and the percent 65 and older increases.

•The birth rate has remained slightly over 17 births per 1000 population the past three years - a rate 10% below the overall state rate, but accounting for 29% of New Mexico births.

•The percentages of low birth weight births and births to teen women have decreased, while the percentage of births to single woman has increased.

•A larger percentage of births are to women 25 to 39 than occur for the state as a whole.

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

BERNALILLO	YEAR 1983	YEAR 1984	YEAR 1985
LIVE BIRTHS	7650	7785	7937
WHITE NON-HISPANIC	3139	3333	3348
WHITE HISPANIC	3711	3726	3784
BLACK	282	238	261
INDIAN	329	320	379
OTHER	189	168	165
COUNTY % OF TOTAL STATE BIRTHS	27.8%	28.5%	28.9%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	BERNALILLO NUMBER	BERNALILLO %	NEW MEXICO %
ALL AGES	7937	100.0%	100.0%
<15	13	0.2%	0.2%
15-17	351	4.4%	5.4%
18-19	714	9.0%	9.9%
(15-19)	1065	13.4%	15.3%
20-24	2524	31.8%	33.6%
25-29	2393	30.1%	28.8%
30-34	1425	18.0%	15.7%
35-39	469	5.9%	5.5%
=> 40	48	0.6%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
BERNALILLO:										
1985	66.5	0.8	62.3	34.7	102.6	113.1	98.1	63.9	24.7	--
1984	67.5	1.3	48.3	26.6	78.3	100.8	106.3	73.4	27.9	1.8
1983	67.1	1.2	49.8	30.0	77.2	105.2	104.1	68.3	24.7	2.2
1982	68.8	1.1	51.4	30.6	80.1	107.6	111.2	67.1	23.3	1.9
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



BERNALILLO COUNTY

BERNALILLO COUNTY DEATHS

			TOTAL	
YEAR	MALE	FEMALE	NUMBER	CRUDE DEATH RATE
1985	1719	1319	3038	6.5
1984	1551	1299	2850	6.3
1983	1535	1206	2741	6.2
1982	1519	1254	2773	6.4
COUNTY % OF TOTAL (1985) STATE DEATHS	31.4	31.7	31.5	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

BERNALILLO CO.	
MALE	851.8
FEMALE	688.7
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate increased to 6.5 deaths per 1000 population in 1985, as the county accounts for 32% of all New Mexico resident deaths. Adjusted rates show levels of mortality higher than state levels for males but slightly lower for females.
- Heart disease, cancer and accidents account for over half of all deaths in the county, however levels of mortality are lower than expected for heart disease and cancer, but 8% higher than expected for accidents. The level of deaths from chronic obstructive pulmonary disease (COPD) is twice the expected level. The observed deaths for diabetes, suicide and alcoholism are also notably higher than expected.
- Both the neonatal and postneonatal components of infant mortality are higher than state levels.
- Over 98% of resident births occur within the county. Overall, the county experiences 2.6 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

BERNALILLO CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	10525	11280	93	100.0	100.0
HEART DISEASE	2484	4078	61	23.6	23.7
MALIGNANT NEOPLASMS	2303	2411	76	21.9	20.2
ALL ACCIDENT	807	749	108	7.7	10.0
MOTOR VEHICLE	448	400	112	4.3	6.1
ALL OTHER	359	349	103	3.4	3.9
STROKE	632	890	71	6.0	6.1
C.O.P.D.	593	296	200	5.6	4.7
DIABETES MELLITUS	290	191	152	2.8	2.7
SUICIDE	340	196	173	3.2	2.8
INFLUENZA/PNEUMONIA	270	266	102	2.6	2.7
CIRRHOSIS OF LIVER	192	198	97	1.8	1.9
HOMICIDE	218	179	122	2.1	1.9
ALCOHOLISM	164	29	566	1.6	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	147	191	77	1.4	1.6
ATHEROSCLEROSIS	130	144	90	1.2	1.1
CONGENITAL ANOMALIES	95	109	87	.9	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

	% OF TOTAL DEATHS	
AGE	BERNALILLO	NEW MEXICO
ALL	100.0	100.0
< 1	3.1	2.9
1 - 4	.5	.8
5- 14	.6	.8
15-24	2.8	3.7
25-34	4.3	4.4
35-44	4.1	4.0
45-54	6.7	6.4
55-64	14.4	13.4
65-74	22.1	22.1
75-84	24.2	24.5
85+	17.2	16.9

INFANT MORTALITY

BERNALILLO COUNTY	TOTAL INFANT		NEONATAL		POSTNEONATAL	
	Number	Rate	Number	Rate	Number	Rate
1980	81	10.5	58	7.5	23	3.0
1981	66	9.0	41	5.6	25	3.4
1982	89	11.6	57	7.4	32	4.2
1983	90	11.8	52	6.8	38	5.0
1984	76	9.8	57	7.3	19	2.4
1985	99	12.5	60	7.6	39	4.9
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

BERNALILLO COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Albuquerque-Urban	7248	2852	2.5
Albuquerque-Rural	518	141	3.7
Sandia/Kirtland AFB (Military)	9	1	9.0
Other & Unspecified	162	44	3.7
COUNTY TOTAL	7937	3038	2.6
% OF COUNTY RESIDENT BIRTHS:			
Born in County	98.85%	XXXXX	XXXXX
Born in Other NM County	0.98%	XXXXX	XXXXX
Born Out of State	0.16%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

CATRON COUNTY

CATRON COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	2198	2720	2800	3300
% 19 AND YOUNGER	38.1%	34.9%	31.6%	28.9%
% 65 AND OLDER	12.3%	13.5%	14.2%	13.7%
% WHITE NON-HISPANIC	56.0%	69.4%	XXXXX	XXXXX
% WHITE HISPANIC	42.2%	28.3%	XXXXX	XXXXX
% BLACK	0.3%	0.4%	XXXXX	XXXXX
% INDIAN	0.5%	1.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	14.3%	17.4%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.2%	0.2%	0.2%	0.2%

- Catron County ranks as the state's 31st most populous county and has a population density of .4 persons per square mile. Only 0.2% of the state's population resides in Catron County.
- The county population is projected to reach 3300 persons by the year 2000 with a 17% decrease in the percentage of persons 19 and younger as compared to 1980.
- The birth rate which had been decreasing increased to a level of 12.1 births per 1000 population in 1985 which is 36% lower than the state rate, accounting for .1% of all the state's births.
- The percent low birth weight births and births to teen mothers dropped notably. Births to single mothers increased but remained well below the state percentage.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

CATRON	YEAR 1983	1984	1985
LIVE BIRTHS	37	30	34
WHITE NON-HISPANIC	21	26	20
WHITE HISPANIC	15	4	13
BLACK	0	0	0
INDIAN	1	0	1
OTHER	0	0	0
COUNTY % OF TOTAL STATE BIRTHS	.1%	.1%	.1%



BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	CATRON NUMBER	CATRON %	NEW MEXICO %
ALL AGES	34	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	0	0.0%	5.4%
18-19	3	8.8%	9.9%
(15-19)	3	8.8%	15.3%
20-24	11	32.4%	33.6%
25-29	10	29.4%	28.8%
30-34	5	14.7%	15.7%
35-39	3	8.8%	5.5%
=> 40	2	5.9%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
CATRON:										
1985	54.8	--	25.9	--	56.6	85.9	144.9	47.6	31.3	6.5
1984	53.0	--	37.0	--	102.0	132.4	37.0	62.5	65.2	--
1983	64.7	--	51.5	--	142.9	130.4	73.4	113.4	21.5	--
1982	72.5	--	38.2	36.1	41.7	149.3	142.9	53.2	44.9	8.3
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







CATRON COUNTY

CATRON COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	16	10	26	9.3
1984	16	8	24	8.7
1983	16	12	28	10.0
1982	19	8	27	10.0
COUNTY % OF TOTAL (1985) STATE DEATHS	0.3	0.2	0.3	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

	1983 - 85 AVERAGE
CATRON CO.	
MALE	993.2
FEMALE	778.0
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate increased from 8.7 deaths per 1000 population to 9.3, accounting for 0.3% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 68% of all deaths. Observed deaths are less than expected for heart disease and cancer, but over three times the expected level for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), and influenza/pneumonia.
- Due to relatively small number of births each year, infant mortality figures fluctuate and must be cautiously interpreted.
- Over half of resident births occur in another county and 41% in another state. Overall, the county experiences slightly more than one birth for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

CATRON CO. — FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	97	105	92	100.0	100.0
HEART DISEASE	30	41	73	30.9	23.7
MALIGNANT NEOPLASMS	20	22	91	20.6	20.2
ALL ACCIDENT	16	5	320	16.5	10.0
MOTOR VEHICLE	13	2	650	13.4	6.1
ALL OTHER	3	3	100	3.1	3.9
STROKE	3	9	33	3.1	6.1
C.O.P.D.	6	3	200	6.2	4.7
DIABETES MELLITUS	--	2	--	--	2.7
SUICIDE	1	1	100	1.0	2.8
INFLUENZA/PNEUMONIA	6	3	200	6.2	2.7
CIRRHOSIS OF LIVER	--	2	--	--	1.9
HOMICIDE	1	1	100	1.0	1.9
ALCOHOLISM	1	.18	555	1.0	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	1	1	100	1.0	1.6
ATHEROSCLEROSIS	1	1	100	1.0	1.1
CONGENITAL ANOMALIES	1	1	100	1.0	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS CATRON	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.8	2.9
1 - 4	--	.8
5- 14	3.8	.8
15-24	2.6	3.7
25-34	5.1	4.4
35-44	2.6	4.0
45-54	2.6	6.4
55-64	7.7	13.4
65-74	28.2	22.1
75-84	26.9	24.5
85+	16.7	16.9

INFANT MORTALITY

CATRON COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	0	0.0	0	0.0	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	2	50.0	0	0.0	2	50.0
1983	0	0.0	0	0.0	0	0.0
1984	1	33.3	1	33.3	0	0.0
1985	2	58.8	2	58.8	0	0.0
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

CATRON COUNTY PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Reserve-All	14	11	1.3
Other & Unspecified	20	15	1.3
COUNTY TOTAL	34	26	1.3
% OF COUNTY RESIDENT BIRTHS:			
Born in County	5.88%	XXXXX	XXXXX
Born in Other NM County	52.94%	XXXXX	XXXXX
Born Out of State	41.18%	XXXXX	XXXXX

***The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

CHAVES COUNTY

CHAVES COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	43335	51103	56500	63200
% 19 AND YOUNGER	42.9%	34.3%	32.8%	32.1%
% 65 AND OLDER	9.0%	14.1%	15.2%	16.0%
% WHITE NON-HISPANIC	66.9%	66.1%	XXXXX	XXXXX
% WHITE HISPANIC	27.9%	30.4%	XXXXX	XXXXX
% BLACK	3.3%	2.2%	XXXXX	XXXXX
% INDIAN	1.4%	0.6%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	20.4%	14.3%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	4.3%	3.9%	3.9%	3.4%

- Chaves County ranks as the state's 7th most populous county and has a population density of 9.3 persons per square mile. About 4% of the state's population resides in Chaves County.
- The county population is projected to reach over 63,000 persons by the year 2000 with nearly one-third being 19 years of age or younger.
- The birth rate dropped in 1985 to 17.9 births per 1000 population - a rate 5% below the state rate, accounting for nearly 4% of New Mexico births. The fertility rates shows a higher than statewide rate of childbearing for Chaves County women.
- The percentages of low birth weight births and births to single mothers are below state levels, but the percent of births to teen mothers though decreasing is 18% higher than statewide.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

CHAVES	1983	1984	1985
LIVE BIRTHS	1173	1060	1012
WHITE NON-HISPANIC	563	461	469
HISPANIC	568	566	510
BLACK	29	19	23
INDIAN	7	4	4
OTHER	6	10	6
COUNTY % OF TOTAL STATE BIRTHS	4.3%	3.9%	3.7%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	CHAVES NUMBER	CHAVES %	NEW MEXICO %
ALL AGES	1012	100.0%	100.0%
<15	6	0.6%	0.2%
15-17	77	7.6%	5.4%
18-19	102	10.1%	9.9%
(15-19)	179	17.7%	15.3%
20-24	390	38.5%	33.6%
25-29	269	26.6%	28.8%
30-34	125	12.4%	15.7%
35-39	36	3.6%	5.5%
=> 40	7	0.7%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
CHAVES:										
1985	85.8	3.0	81.2	60.1	110.6	169.3	141.4	59.4	20.0	--
1984	87.3	1.7	71.6	48.8	107.1	174.3	126.9	73.3	23.5	2.7
1983	101.1	1.3	84.4	44.8	145.9	201.8	160.3	65.1	34.9	1.1
1982	100.6	1.4	88.6	51.2	146.8	204.5	133.8	86.6	26.7	3.3
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







CHAVES COUNTY

CHAVES COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	290	244	534	9.5
1984	275	235	510	8.7
1983	270	234	504	9.0
1982	279	211	490	9.2
COUNTY % OF TOTAL (1985) STATE DEATHS	5.3	5.9	5.5	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

CHAVES CO.	
MALE	865.4
FEMALE	724.3
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate has increased to 9.5 deaths per 1000 population, accounting for 5.5% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and accidents rank as the leading causes of death, accounting for 52% of all deaths. Observed deaths from heart disease are only 59% of expected, cancer is 7% higher than expected and accidents 39% higher than expected. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, homicide and alcoholism.
- Infant mortality is lower in the county than statewide.
- Over 95% of resident births occur within the county. Overall, the county experiences 1.9 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

CHAVES CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1999	2000	100	100.0	100.0
HEART DISEASE	456	775	59	22.8	23.7
MALIGNANT NEOPLASMS	458	427	107	22.9	20.2
ALL ACCIDENT	133	96	139	6.7	10.0
MOTOR VEHICLE	82	47	174	4.1	6.1
ALL OTHER	51	49	104	2.6	3.9
STROKE	141	174	81	7.1	6.1
C.O.P.D.	88	58	152	4.4	4.7
DIABETES MELLITUS	65	35	186	3.3	2.7
SUICIDE	39	24	163	2.0	2.8
INFLUENZA/PNEUMONIA	41	51	80	2.1	2.7
CIRRHOSIS OF LIVER	32	29	110	1.6	1.9
HOMICIDE	35	20	175	1.8	1.9
ALCOHOLISM	28	4	700	1.4	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	31	26	119	1.6	1.6
ATHEROSCLEROSIS	37	29	128	1.9	1.1
CONGENITAL ANOMALIES	12	15	80	.6	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS CHAVES	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.4	2.9
1 - 4	.6	.8
5- 14	.8	.8
15-24	2.8	3.7
25-34	2.6	4.4
35-44	2.7	4.0
45-54	5.6	6.4
55-64	13.9	13.4
65-74	23.2	22.1
75-84	28.5	24.5
85+	17.0	16.9

INFANT MORTALITY

CHAVES COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	12	11.9	7	7.0	5	5.0
1981	15	15.2	10	10.1	5	5.1
1982	13	11.4	9	7.9	4	3.5
1983	15	12.8	10	8.5	5	4.3
1984	12	11.3	9	8.5	3	2.8
1985	10	9.9	6	5.9	4	4.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

CHAVES COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Dexter-All	55	24	2.3
Hagerman-Urban	14	11	1.3
Hagerman-Rural	9	3	3.0
Roswell-Urban	842	451	1.9
Roswell-Rural	66	39	1.7
Other & Unspecified	26	6	4.3
COUNTY TOTAL	1012	534	1.9
% OF COUNTY RESIDENT BIRTHS:			
Born in County	95.65%	XXXXX	XXXXX
Born in Other NM County	4.15%	XXXXX	XXXXX
Born Out of State	0.20%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

CIBOLA COUNTY

CIBOLA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	#####	30347	25300	31100
% 19 AND YOUNGER	###	42.8%	40.9%	35.4%
% 65 AND OLDER	###	4.6%	7.2%	10.9%
% WHITE NON-HISPANIC	###	###	XXXXX	XXXXX
% WHITE HISPANIC	###	###	XXXXX	XXXXX
% BLACK	###	###	XXXXX	XXXXX
% INDIAN	###	###	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	###	###	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	###	2.3%	1.7%	1.7%

- Cibola County ranks as the state's 15th most populous county with a population density of 6.1 persons per square mile, accounting for 1.7% of the total state population.

- The county population is projected to be at 31,100 persons by the year 2000 - a 2.5% increase from the 1980 figure with doubling of the percentage of persons 65 and older.

- Remaining the same as for the previous year the birth rate is 10% higher than the state rate. The county rate of 20.8 births per 1000 population accounts for 1.9% of New Mexico resident births.

- Percentages of births to teens and births to single women exceed state levels, but low birth weight births are below state levels.

***DATA AVAILABLE ONLY IN A FORM INCLUDING BOTH VALENCIA AND CIBOLA



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

CIBOLA	1983	1984	1985
LIVE BIRTHS	574	533	525
WHITE			
NON-HISPANIC	117	110	88
HISPANIC	207	183	196
BLACK	1	2	8
INDIAN	246	235	229
OTHER	3	3	4
COUNTY % OF TOTAL STATE BIRTHS	2.1%	1.9%	1.9%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	CIBOLA NUMBER	CIBOLA %	NEW MEXICO %
ALL AGES	525	100.0%	100.0%
<15	3	0.6%	0.2%
15-17	36	6.9%	5.4%
18-19	60	11.4%	9.9%
(15-19)	96	18.3%	15.3%
20-24	170	32.4%	33.6%
25-29	140	26.7%	28.8%
30-34	80	15.2%	15.7%
35-39	32	6.1%	5.5%
=> 40	4	0.8%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
CIBOLA:										
1985	99.0	2.5	99.1	61.9	155.0	263.2	162.6	67.6	33.0	--
1984	83.2	2.4	82.4	56.8	129.2	125.3	120.9	66.1	28.9	3.6
1983	83.8	2.9	81.8	38.3	161.8	121.8	128.0	62.8	34.4	3.4
1982	96.1	3.7	94.0	50.9	173.2	159.0	132.4	72.7	25.8	3.4
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







CIBOLA COUNTY

CIBOLA COUNTY
DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	78	53	131	5.2
1984	75	53	128	5.0
1983	64	52	116	4.2
1982	85	54	139	4.7
COUNTY % OF TOTAL (1985) STATE DEATHS	1.4	1.3	1.4	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
CIBOLA CO.	
MALE	709.3
FEMALE	764.1
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 5.2 deaths per 1000 population accounts for 1.4% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for females but lower for males.
- Mortality by cause is based on combined Cibola/Valencia county data and shows heart disease, cancer and accidents to be the leading causes of death, accounting for 52% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but higher than expected for accidents, chronic obstructive pulmonary disease (COPD), diabetes, suicide, influenza/pneumonia and alcoholism.
- Postneonatal mortality exceeded state levels in 1985.
- Nearly 51% of resident births occur in another county. Overall, the county experiences 4 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

VALENCIA CO. (CIBOLA INCLUDED)　　FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1506	1545	97	100.0	100.0
HEART DISEASE	316	547	58	21.0	23.7
MALIGNANT NEOPLASMS	276	324	85	18.3	20.2
ALL ACCIDENT	194	105	185	12.9	10.0
MOTOR VEHICLE	126	56	225	8.4	6.1
ALL OTHER	68	49	139	4.5	3.9
STROKE	81	117	69	5.4	6.1
C.O.P.D.	54	41	132	3.6	4.7
DIABETES MELLITUS	48	25	192	3.2	2.7
SUICIDE	52	26	200	3.5	2.8
INFLUENZA/PNEUMONIA	45	36	125	3.0	2.7
CIRRHOSIS OF LIVER	26	27	96	1.7	1.9
HOMICIDE	24	25	96	1.6	1.9
ALCOHOLISM	22	4	550	1.5	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	33	39	85	2.2	1.6
ATHEROSCLEROSIS	19	19	100	1.3	1.1
CONGENITAL ANOMALIES	24	20	120	1.6	.9

PERCENTAGE DISTRIBUTION
OF DEATHS BY AGE
1983-1985

AGE	% OF TOTAL DEATHS CIBOLA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.5	2.9
1 - 4	1.9	.8
5- 14	.8	.8
15-24	8.3	3.7
25-34	6.1	4.4
35-34	5.1	4.0
45-54	8.5	6.4
55-64	13.6	13.4
65-74	20.3	22.1
75-84	20.0	24.5
85+	12.0	16.9

INFANT MORTALITY

CIBOLA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	**	**	**	**	**	**
1981	**	**	**	**	**	**
1982	7	10.6	4	6.1	3	4.5
1983	2	3.5	1	1.7	1	1.7
1984	5	9.4	4	7.5	1	1.9
1985	6	11.4	3	5.7	3	5.7
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

CIBOLA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Grants-Urban	196	50	3.9
Grants-Rural	38	0	---
Milan-Urban	32	11	2.9
Milan-Rural	9	0	---
Other & Unspecified	250	70	3.6
COUNTY TOTAL	525	131	4.0
% OF COUNTY RESIDENT BIRTHS:			
Born in County	49.14%	XXXXX	XXXXX
Born in Other NM County	50.86%	XXXXX	XXXXX
Born Out of State	----	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL <28 days, POSTNEONATAL = 28 days to <1 year.

COLFAX COUNTY

COLFAX COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	12170	13667	14300	16600
% 19 AND YOUNGER	40.8%	35.4%	33.8%	31.3%
% 65 AND OLDER	12.3%	13.0%	13.1%	13.3%
% WHITE NON-HISPANIC	45.6%	51.3%	XXXXX	XXXXX
% WHITE HISPANIC	53.1%	47.3%	XXXXX	XXXXX
% BLACK	0.5%	0.4%	XXXXX	XXXXX
% INDIAN	0.4%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	20.1%	13.5%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.2%	1.0%	1.0%	0.9%

- Colfax County, ranks as the state's 21st most populous county and has a population density of 3.8 persons per square mile. 1% of the state's population resides in Colfax County.

- The county population is projected to reach almost 17,000 persons by the year 2000.

- The present birth rate of 17.9 births per 1000 population is an increase from previous years leaving the county rate 5% below the state rate. The county accounts for .9% of all New Mexico resident births.

- A slightly higher percentage of births to teen mothers is noted for the county as compared to state levels, and while the percent of births to single mothers is lower than statewide, the percent has increased. The level of low birth weight births dropped for 1985.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

COLFAX	YEAR 1983	1984	1985
LIVE BIRTHS	250	212	256
WHITE NON-HISPANIC	73	76	91
HISPANIC	173	131	163
BLACK	2	1	0
INDIAN	2	3	0
OTHER	0	1	2
COUNTY % OF TOTAL STATE BIRTHS	.9%	.8%	.9%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	COLFAX NUMBER	COLFAX %	NEW MEXICO %
ALL AGES	256	100.0%	100.0%
<15	1	0.4%	0.2%
15-17	21	8.2%	5.4%
18-19	26	10.2%	9.9%
(15-19)	47	18.4%	15.3%
20-24	86	33.6%	33.6%
25-29	74	28.9%	28.8%
30-34	39	15.2%	15.7%
35-39	7	2.7%	5.5%
=> 40	2	0.8%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
COLFAX										
1985	83.4	1.7	79.4	60.2	107.0	145.3	150.1	79.1	14.2	--
1984	77.5	1.8	59.1	29.6	112.7	136.7	129.4	66.8	26.2	1.7
1983	90.5	1.8	68.8	34.8	130.4	171.8	132.0	77.3	44.2	--
1982	96.2	3.9	81.8	51.9	136.1	173.3	142.9	92.9	22.4	7.3
NEW MEXICO: 1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







COLFAX COUNTY

COLFAX COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	74	51	125	8.7
1984	64	38	102	7.6
1983	59	52	111	8.2
1982	58	44	102	7.5
COUNTY % OF TOTAL (1985) STATE DEATHS	1.4	1.2	1.3	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

COLFAX CO.	
MALE	859.0
FEMALE	619.2
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 8 7 deaths per 1000 population accounts for 1 3% of all New Mexico resident deaths Adjusted rates show lower than state levels of mortality for famales and higher for males

- Heart disease, cancer and accidents are the leading causes of death, accounting for 57% of all deaths Observed deaths are lower than expected levels for heart disease and cancer, but 27% higher than expected for accidents Observed deaths are higher than expected for chronic obstrurtive pulmonary disease (COPD), diabetes, suicide and alcoholism

- Neonatal mortality exceeded the state rate in 1985

- Nearly 80% of resident births occur within the county Overall, the county experiences 2 births for every death

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

COLFAX CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	442	523	85	100.0	100.0
HEART DISEASE	125	204	61	28.3	23.7
MALIGNANT NEOPLASMS	92	109	84	20.8	20.2
ALL ACCIDENT	33	26	127	7.5	10.0
MOTOR VEHICLE	22	13	169	5.0	6.1
ALL OTHER	11	13	85	2.5	3.9
STROKE	24	46	52	5.4	6.1
C.O.P.D.	19	15	127	4.3	4.7
DIABETES MELLITUS	14	9	156	3.2	2.7
SUICIDE	10	7	143	2.3	2.8
INFLUENZA/PNEUMONIA	13	14	93	2.9	2.8
CIRRHOSIS OF LIVER	8	7	114	1.8	2.7
HOMICIDE	5	5	100	1.1	1.9
ALCOHOLISM	3	1	300	.7	1.9
CERTAIN CONDITIONS					1.7
IN PERINATAL PERIOD	6	6	100	1.4	1.6
ATHEROSCLEROSIS	4	8	50	.9	1.1
CONGENITAL ANOMALIES	--	3	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS COLFAX	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	1.8	2.9
1 - 4	.3	.8
5- 14	.6	.8
15-24	3.2	3.7
25-34	2.9	4.4
35-44	3.2	4.0
45-54	4.1	6.4
55-64	12.7	13.4
65-74	22.1	22.1
75-84	27.1	24.5
85+	21.8	16.9

INFANT MORTALITY

COLFAX COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	5	20.9	3	12.6	2	8.4
1981	1	4.6	1	4.6	0	0.0
1982	1	4.0	0	0.0	1	4.0
1983	2	8.0	1	4.0	1	4.0
1984	2	9.4	2	9.4	0	0.0
1985	2	7.8	2	7.8	0	0.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

COLFAX COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Cimarron-All	28	14	2.0
Raton-Urban	144	81	1.8
Raton-Rural	12	5	2.4
Springer-Urban	22	14	1.6
Springer-Rural	3	2	1.5
Eagle Nest-All	15	1	15.0
Other & Unspecified	32	8	4.0
COUNTY TOTAL	256	125	2.0
% OF COUNTY RESIDENT BIRTHS:			
Born in County	79.30%	XXXXX	XXXXX
Born in Other NM County	13.28%	XXXXX	XXXXX
Born Out of State	7.42%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

CURRY COUNTY

CURRY COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	39517	42019	42600	41700
% 19 AND YOUNGER	41.1%	36.9%	35.5%	31.7%
% 65 AND OLDER	7.4%	9.6%	11.5%	14.4%
% WHITE NON-HISPANIC	74.5%	71.6%	XXXXX	XXXXX
% WHITE HISPANIC	17.4%	19.2%	XXXXX	XXXXX
% BLACK	6.4%	6.8%	XXXXX	XXXXX
% INDIAN	0.3%	0.4%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	15.1%	11.7%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	3.9%	3.2%	2.9%	2.3%

- Curry County ranks as the state's 11th most populous county and has a population density of 30.4 persons per square mile. About 3% of the state population resides in Curry County.
- The county population is projected to drop to less than 42,000 persons by the year 2000, accounting for a decreasing percentage of the state's population.
- Accounting for 3.3% of the state's births the county birth rate decreased to 21.0 births per 1000 population - a level 11% higher than state-wide.
- The percents of low birth weight births and births to teen mothers are down from the prior year, but the percent of births to teens exceeds state levels. The percent of births to single mothers increased but remains below state level.


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

CURRY	YEAR 1983	1984	1985
LIVE BIRTHS	904	980	893
WHITE NON-HISPANIC	510	571	505
HISPANIC	274	280	270
BLACK	77	97	77
INDIAN	7	5	4
OTHER	36	27	37
COUNTY % OF TOTAL STATE BIRTHS	3.3%	3.6%	3.3%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	CURRY NUMBER	CURRY %	NEW MEXICO %
ALL AGES	893	100.0%	100.0%
<15	3	0.3%	0.2%
15-17	51	5.7%	5.4%
18-19	110	12.3%	9.9%
(15-19)	161	18.0%	15.3%
20-24	381	42.7%	33.6%
25-29	233	26.1%	28.8%
30-34	91	10.2%	15.7%
35-39	20	2.2%	5.5%
=> 40	4	0.4%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
CURRY:										
1985	96.8	1.7	98.5	51.3	171.9	248.7	142.5	55.7	13.1	--
1984	97.7	1.1	87.5	50.5	145.4	202.5	134.7	60.2	14.9	1.0
1983	89.7	1.6	92.6	53.6	153.8	181.4	114.7	50.4	17.1	3.3
1982	91.3	1.1	84.5	50.7	137.5	178.9	125.3	56.3	24.0	3.8
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

CURRY COUNTY

CURRY COUNTY DEATHS

			TOTAL	
YEAR	MALE	FEMALE	NUMBER	CRUDE DEATH RATE
1985	134	146	280	6.6
1984	176	132	308	7.0
1983	168	121	289	6.6
1982	178	134	312	7.3
COUNTY % OF TOTAL (1985) STATE DEATHS	2.4	3.5	2.9	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

CURRY CO.	
MALE	784.1
FEMALE	629.9
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 6.6 deaths per 1000 population accounts for 2.9% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for both males and females.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 59% of all deaths. Observed deaths for heart disease and cancer are below expected levels, while deaths from accidents, chronic obstructive pulmonary disease (COPD), diabetes, homicide and alcoholism are notably above expected levels.
- Postneonatal mortality is higher than statewide.
- Over 73% of resident births occur within the county, while nearly 3% occur in another state. Overall, the county experiences 3.2 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

CURRY CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1235	1268	97	100.0	100.0
HEART DISEASE	373	471	79	30.2	23.7
MALIGNANT NEOPLASMS	247	257	96	20.0	20.2
ALL ACCIDENT	103	77	134	8.3	10.0
MOTOR VEHICLE	53	40	133	4.3	6.1
ALL OTHER	50	37	135	4.0	3.9
STROKE	115	108	106	9.3	6.1
C.O.P.D.	47	33	142	3.8	4.7
DIABETES MELLITUS	29	21	138	2.3	2.7
SUICIDE	21	19	111	1.7	2.8
INFLUENZA/PNEUMONIA	36	32	113	2.9	2.7
CIRRHOSIS OF LIVER	12	19	63	1.0	1.9
HOMICIDE	30	17	176	2.4	1.9
ALCOHOLISM	13	3	433	1.1	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	16	25	64	1.3	1.6
ATHEROSCLEROSIS	20	19	105	1.6	1.1
CONGENITAL ANOMALIES	18	13	138	1.5	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS	
	CURRY	NEW MEXICO
ALL	100.0	100.0
< 1	2.7	2.9
1 - 4	.8	.8
5- 14	.7	.8
15-24	2.9	3.7
25-34	2.5	4.4
35-44	3.1	4.0
45-54	7.2	6.4
55-64	12.1	13.4
65-74	21.0	22.1
75-84	27.8	24.5
85+	19.3	16.9

INFANT MORTALITY

CURRY COUNTY	TOTAL INFANT		NEONATAL		POSTNEONATAL	
	Number	Rate	Number	Rate	Number	Rate
1980	10	10.4	6	6.2	4	4.1
1981	10	11.0	8	8.8	2	2.2
1982	11	11.9	5	5.4	6	6.5
1983	4	4.4	2	2.2	2	2.2
1984	8	8.2	2	2.0	6	6.1
1985	12	13.4	3	3.4	9	10.1
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

CURRY COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Clovis-Urban	694	249	2.8
Clovis-Rural	44	8	5.5
Texico-All	28	5	5.6
Melrose-All	4	10	0.4
Cannon AFB (Military)	112	2	56.0
Other & Unspecified	11	6	1.8
COUNTY TOTAL	893	280	3.2
% OF COUNTY RESIDENT BIRTHS:			
Born in County	[illegible]7%	XXXXX	XXXXX
Born in Other NM County	23.85%	XXXXX	XXXXX
Born Out of State	2.58%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

DE BACA COUNTY

DE BACA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	2547	2454	2400	2300
% 19 AND YOUNGER	35.1%	28.6%	27.8%	32.1%
% 65 AND OLDER	18.1%	21.0%	22.1%	20.5%
% WHITE NON-HISPANIC	58.8%	68.5%	XXXXX	XXXXX
% WHITE HISPANIC	40.8%	30.9%	XXXXX	XXXXX
% BLACK	0.2%	0.0%	XXXXX	XXXXX
% INDIAN	0.1%	0.1%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	24.5%	14.5%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.3%	0.2%	0.2%	0.1%

- De Baca County ranks as the state's second least populated county and has a population density of one person per square mile, accounting for 0.2% of the state's population.
- The county population in the year 2000 is not projected to differ greatly from the 1980 census count resulting in the county's accounting for a decreasing percent of the total state population.
- Birth rates have run below state levels. The 1985 rate of 15 births per 1000 population is 21% below the state birth rate as the county accounts for only .1% of all New Mexico resident births.
- The percent of births to teen mothers is well above the state level, while the percentages of births to single mothers and of low birth weight births have dropped below state levels.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

DEBACA	1983	1984	1985
LIVE BIRTHS	42	38	36
WHITE NON-HISPANIC	25	16	19
HISPANIC	17	21	17
BLACK	0	0	0
INDIAN	0	0	0
OTHER	0	1	0
COUNTY % OF TOTAL STATE BIRTHS	.2%	.1%	.1%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	DE BACA NUMBER	DE BACA %	NEW MEXICO %
ALL AGES	36	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	3	8.3%	5.4%
18-19	8	22.2%	9.9%
(15-19)	11	30.6%	15.3%
20-24	11	30.6%	33.6%
25-29	11	30.6%	28.8%
30-34	1	2.8%	15.7%
35-39	1	2.8%	5.5%
=> 40	1	2.8%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
DE BACA:										
1985	80.7	--	123.6	56.6	222.2	118.3	180.3	14.1	16.4	--
1984	91.1	--	82.5	64.5	114.3	206.3	138.9	96.8	14.5	--
1983	97.2	--	108.9	46.2	222.2	197.0	148.6	76.9	--	16.3
1982	92.8	--	89.1	92.3	83.3	276.9	121.6	30.8	28.2	--
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







DE BACA COUNTY

DE BACA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	26	18	44	18.3
1984	22	15	37	16.0
1983	20	18	38	15.8
1982	21	12	33	13.6
COUNTY % OF TOTAL (1985) STATE DEATHS	0.5	0.4	0.5	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

DEBACA CO.	
MALE	1206.6
FEMALE	781.2
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate increased to 18.3 deaths per 1000 population, accounting for 0.5% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both males and females.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 61% of all deaths. Observed deaths are less than expected for heart disease and cancer, but 80% higher than expected for accidents.
- Due to the relatively small number of births each year, infant mortality figures fluctuate and must be cautiously interpreted.
- 36% of resident births occur within the county. Overall, the county experiences less than one birth for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

DEBACA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	127	144	88	100.0	100.0
HEART DISEASE	46	59	78	36.2	23.7
MALIGNANT NEOPLASMS	23	29	79	18.1	20.2
ALL ACCIDENT	9	5	180	7.1	10.0
MOTOR VEHICLE	5	2	250	3.9	6.1
ALL OTHER	4	3	133	3.1	3.9
STROKE	5	14	36	3.9	6.1
C.O.P.D.	4	4	100	3.1	4.7
DIABETES MELLITUS	--	3	--	--	2.7
SUICIDE	1	1	100	.8	2.8
INFLUENZA/PNEUMONIA	5	4	125	3.9	2.7
CIRRHOSIS OF LIVER	1	2	50	.8	1.9
HOMICIDE	2	1	200	1.6	1.9
ALCOHOLISM	1	.23	435	.8	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	--	1	--	--	1.6
ATHEROSCLEROSIS	1	3	33	.8	1.1
CONGENITAL ANOMALIES	--	1	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS DEBACA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	.8	2.9
1 - 4	--	.8
5- 14	--	.8
15-24	1.7	3.7
25-34	4.2	4.4
35-44	2.5	4.0
45-54	.8	6.4
55-64	5.9	13.4
65-74	26.9	22.1
75-84	29.4	24.5
85+	27.7	16.9

INFANT MORTALITY

DE BACA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	0	0.0	0	0.0	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	0	0.0	0	0.0	0	0.0
1983	1	23.8	1	23.8	0	0.0
1984	0	0.0	0	0.0	0	0.0
1985	0	0.0	0	0.0	0	0.0
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

DE BACA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Ft. Sumner-Urban	19	32	0.6
Ft. Sumner-Rural	14	9	1.6
Other & Unspecified	3	3	1.0
COUNTY TOTAL	36	44	0.8
% OF COUNTY RESIDENT BIRTHS:			
Born in County	36.11%	XXXXX	XXXXX
Born in Other NM County	61.11%	XXXXX	XXXXX
Born Out of State	2.78%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

DONA ANA COUNTY

DONA ANA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	69773	96340	118300	179900
% 19 AND YOUNGER	45.6%	38.0%	34.8%	33.4%
% 65 AND OLDER	5.3%	7.2%	8.3%	10.5%
% WHITE NON-HISPANIC	45.9%	44.5%	XXXXX	XXXXX
% WHITE HISPANIC	50.8%	51.8%	XXXXX	XXXXX
% BLACK	1.9%	1.7%	XXXXX	XXXXX
% INDIAN	0.3%	0.8%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	20.5%	18.3%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	6.9%	7.4%	8.2%	9.7%

- Dona Ana County ranks as the state's second most populous county and has a population density of 31.1 persons per square mile, accounting for 8% of the state's total population.
- The county population is projected to reach nearly 180,000 persons by the year 2000 accounting for an increasing percent of the state population.
- The county accounts for almost 8% of all resident births in the state with a birth rate slightly below the overall state rate. The 1985 birth rate of 18.1 births per 1000 live births represents an 11% decrease from the prior year.
- Percentages of low birth weight births, and births to single mothers are below state levels, while that for births to teens is just about at the state level.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

DONA ANA	1983	1984	1985
LIVE BIRTHS	2112	2283	2140
WHITE			
NON-HISPANIC	633	648	596
HISPANIC	1398	1553	1469
BLACK	52	37	36
INDIAN	9	17	22
OTHER	20	28	17
COUNTY % OF TOTAL STATE BIRTHS	7.7%	8.3%	7.8%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	DONA ANA NUMBER	DONA ANA %	NEW MEXI %
ALL AGES	2140	100.0%	100.0%
<15	4	0.2%	0.2%
15-17	109	5.1%	5.4%
18-19	222	10.4%	9.9%
(15-19)	331	15.5%	15.3%
20-24	756	35.3%	33.6%
25-29	620	29.0%	28.8%
30-34	295	13.8%	15.7%
35-39	118	5.5%	5.5%
=> 40	16	0.7%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
DONA ANA:										
1985	74.3	0.8	59.6	32.9	99.1	109.3	102.6	77.6	32.7	0.2
1984	80.3	1.2	47.9	32.6	63.4	104.8	153.3	90.2	36.5	3.2
1983	77.2	2.1	54.3	40.8	67.9	101.1	136.5	84.3	35.0	4.1
1982	78.6	0.9	54.2	42.0	66.4	109.9	133.7	87.2	28.7	5.0
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







DONA ANA COUNTY

DONA ANA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	303	282	585	4.9
1984	323	281	604	5.4
1983	316	241	557	5.2
1982	305	218	523	5.1
COUNTY % OF TOTAL (1985) STATE DEATHS	5.5	6.8	6.1	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
DONA ANA CO.	
MALE	647.9
FEMALE	670.8
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased slightly to 4.9 deaths per 1000 population, accounting for 6.1% of New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for males and females.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 56% of all deaths. Observed deaths are less than expected for heart disease and cancer and 10% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide and alcoholism.
- Infant mortality is below state levels.
- About 16% of resident births occur in another state. Overall, the county experiences 3.7 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

DONA ANA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	2072	2403	86	100.0	100.0
HEART DISEASE	524	856	61	25.3	23.7
MALIGNANT NEOPLASMS	448	505	89	21.6	20.2
ALL ACCIDENT	190	172	110	9.2	10.0
MOTOR VEHICLE	121	94	129	5.8	6.1
ALL OTHER	69	78	88	3.3	3.9
STROKE	139	186	75	6.7	6.1
C.O.P.D.	85	63	135	4.1	4.7
DIABETES MELLITUS	79	40	198	3.8	2.7
SUICIDE	46	43	107	2.2	2.8
INFLUENZA/PNEUMONIA	34	57	60	1.6	2.7
CIRRHOSIS OF LIVER	33	41	81	1.6	1.9
HOMICIDE	30	40	75	1.4	1.9
ALCOHOLISM	17	6	283	.8	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	33	48	69	1.6	1.6
ATHEROSCLEROSIS	27	30	90	1.3	1.1
CONGENITAL ANOMALIES	24	27	89	1.2	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS DONA ANA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.6	2.9
1 - 4	.7	.6
5- 14	.7	.8
15-24	3.3	3.7
25-34	3.7	4.4
35-44	3.0	4.0
45-54	7.8	6.4
55-64	13.6	13.4
65-74	23.8	22.1
75-84	23.4	24.5
85+	17.4	16.9

INFANT MORTALITY

DONA ANA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	19	10.3	12	6.5	7	3.8
1981	13	6.7	7	3.6	6	3.1
1982	21	10.5	14	7.0	7	3.5
1983	15	7.1	10	4.7	5	2.4
1984	15	6.6	10	4.4	5	2.2
1985	15	7.0	8	3.7	7	3.3
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

DONA ANA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Hatch-All	56	25	2.2
Las Cruces-Urban	1232	360	3.4
Las Cruces-Rural	49	46	1.1
Mesilla-Urban	2	8	0.3
Mesilla-Rural	54	1	54.0
University Park-All	0	0	---
White Sands Missle Range (Mil.)	69	6	11.5
La Mesa-All	56	16	3.5
Other & Unspecified	622	123	5.1
COUNTY TOTAL	2140	585	3.7
% OF COUNTY RESIDENT BIRTHS:			
Born in County	82.52%	XXXXX	XXXXX
Born in Other NM County	1.40%	XXXXX	XXXXX
Born Out of State	16.07%	XXXXX	XXXXX

***The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

EDDY COUNTY

EDDY COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	41119	47855	52500	61500
% 19 AND YOUNGER	41.5%	35.1%	34.9%	34.5%
% 65 AND OLDER	8.9%	12.5%	13.4%	14.0%
% WHITE NON-HISPANIC	66.6%	66.9%	XXXXX	XXXXX
% WHITE HISPANIC	30.5%	30.6%	XXXXX	XXXXX
% BLACK	2.2%	1.8%	XXXXX	XXXXX
% INDIAN	0.2%	0.3%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	17.8%	10.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	4.0%	3.7%	3.6%	3.3%

- Eddy County ranks as the state's 8th most populous county and has a population density of 12.6 persons per square mile, accounting for 3.6% of the state's total population.

- The county population is projected to exceed 61,000 persons by the year 2000.

- The county birth rate has decreased notably the past four years dropping below the state level in 1983 and further decreasing to 16.9 births per 1000 population by 1985, accounting for 3.2% of New Mexico's births. The fertility rate does indicate however, a higher than statewide rate of childbearing for Eddy County women.

- The percentage of low birth weight births remains above state levels; births to teens has decreased, but remains above state levels; births to single mothers has increased but remains below state levels.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

EDDY	YEAR 1983	1984	1985
LIVE BIRTHS	1019	930	885
WHITE NON-HISPANIC	540	473	442
HISPANIC	450	433	428
BLACK	18	15	11
INDIAN	5	4	1
OTHER	6	5	3
COUNTY % OF TOTAL STATE BIRTHS	3.7%	3.4%	3.2%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	EDDY NUMBER	EDDY %	NEW MEXICO %
ALL AGES	885	100.0%	100.0%
<15	1	0.1%	0.2%
15-17	60	6.8%	5.4%
18-19	111	12.5%	9.9%
(15-19)	171	19.3%	15.3%
20-24	311	35.1%	33.6%
25-29	268	30.3%	28.8%
30-34	90	10.2%	15.7%
35-39	37	4.2%	5.5%
=> 40	7	0.8%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
EDDY:										
1985	80.4	0.5	82.2	49.0	129.8	157.0	142.4	47.8	20.8	--
1984	81.3	1.4	75.5	46.3	121.8	152.8	113.8	65.7	19.4	1.9
1983	91.3	1.9	88.2	57.0	137.8	164.8	129.7	71.5	24.9	2.7
1982	108.7	1.5	106.4	59.5	180.9	195.2	159.3	85.7	23.7	1.6
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







EDDY COUNTY

EDDY COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	268	195	463	8.8
1984	246	219	465	8.6
1983	251	202	453	8.6
1982	237	179	416	8.1
COUNTY % OF TOTAL (1985) STATE DEATHS	4.9	4.7	4.8	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
EDDY CO.	
MALE	952.1
FEMALE	768.6
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 8.8 deaths per 1000 population accounts for 4.8% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 57% of all deaths. Observed deaths are less than expected for heart disease and cancer, but 47% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, homicide, alcoholism and perinatal conditions.
- Neonatal mortality is 10% higher than statewide.
- Over 92% of resident births occur within the county. Overall, the county experiences 1.9 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

EDDY CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1629	1730	94	100.0	100.0
HEART DISEASE	457	664	69	28.1	23.7
MALIGNANT NEOPLASMS	342	367	93	21.0	20.2
ALL ACCIDENT	128	87	147	7.9	10.0
MOTOR VEHIC	65	43	151	4.0	6.1
ALL OTHER	63	44	143	3.9	3.9
STROKE	114	149	77	7.0	6.1
C.O.P.D.	80	49	163	4.9	4.7
DIABETES MELLITUS	44	30	147	2.7	2.7
SUICIDE	27	22	123	1.7	2.8
INFLUENZA/PNEUMONIA	46	44	105	2.8	2.7
CIRRHOSIS OF LIVER	18	24	75	1.1	1.9
HOMICIDE	22	18	122	1.4	1.9
ALCOHOLISM	8	4	200	.5	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	28	24	117	1.7	1.6
ATHEROSCLEROSIS	23	25	92	1.4	1.1
CONGENITAL ANOMALIES	15	14	107	.9	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS EDDY	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.2	2.9
1 - 4	.7	.8
5- 14	.6	.8
15-24	3.0	3.7
25-34	3.1	4.4
35-44	2.7	4.0
45-54	4.4	6.4
55-64	12.4	13.4
65-74	23.0	22.1
75-84	27.4	24.5
85+	20.6	16.9

INFANT MORTALITY

EDDY COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	13	13.2	10	10.2	3	3.1
1981	8	8.3	5	5.2	3	3.1
1982	17	14.6	11	9.5	6	5.2
1983	12	11.8	5	4.9	7	6.9
1984	11	11.8	8	8.6	3	3.2
1985	7	7.9	6	6.8	1	1.1
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

EDDY COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Artesia-Urban	253	109	2.3
Artesia-Rural	52	22	2.4
Carlsbad-Urban	422	290	1.5
Carlsbad-Rural	98	23	4.3
Loving-Urban	27	11	2.5
Loving-Rural	5	2	2.5
Other & Unspecified	28	6	4.7
COUNTY TOTAL	885	463	1.9
% OF COUNTY RESIDENT BIRTHS:			
Born in County	92.66%	XXXXX	XXXXX
Born in Other NM County	7.01%	XXXXX	XXXXX
Born Out of State	0.34%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's are computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

GRANT COUNTY

GRANT COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	22030	26204	27300	32200
% 19 AND YOUNGER	42.5%	37.4%	36.0%	33.9%
% 65 AND OLDER	7.9%	10.5%	11.0%	13.4%
% WHITE NON-HISPANIC	42.2%	47.4%	XXXXX	XXXXX
% WHITE HISPANIC	56.1%	51.2%	XXXXX	XXXXX
% BLACK	0.5%	0.5%	XXXXX	XXXXX
% INDIAN	0.4%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	11.9%	12.4%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	2.2%	2.0%	1.9%	1.7%

- Grant County ranks as the state's 14th most populous county and has a population density of 6.9 persons per square mile, accounting for almost 2% of the state population.
- The county population is projected to reach 32,200 persons by the year 2000, with a 28% increase in the percentage of persons 65 and older.
- The county birth rate increased to 18.9 births per 1000 population thus equaling the state rate, and accounting for 1.9% of the state's resident births.
- The percentage of low birth weight births has decreased below state levels; the percent of births to teens rose in 1985, remaining higher than for the state; and the percent of births to single mothers increased but remained below state levels.

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

GRANT	1983	1984	1985
LIVE BIRTHS	464	470	515
WHITE			
NON-HISPANIC	150	142	172
HISPANIC	300	320	328
BLACK	7	4	6
INDIAN	6	2	5
OTHER	1	2	4
COUNTY % OF TOTAL STATE BIRTHS	1.7%	1.7%	1.9%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	GRANT NUMBER	GRANT %	NEW MEXICO %
ALL AGES	515	100.0%	100.0%
<15	1	0.2%	0.2%
15-17	40	7.8%	5.4%
18-19	65	12.6%	9.9%
(15-19)	105	20.4%	15.3%
20-24	165	32.0%	33.6%
25-29	141	27.4%	28.8%
30-34	69	13.4%	15.7%
35-39	29	5.6%	5.5%
=> 40	5	1.0%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
GRANT:										
1985	81.0	0.8	87.2	54.0	140.4	137.0	140.4	68.7	28.9	1.3
1984	77.0	0.8	59.4	36.9	95.6	156.3	111.7	61.7	28.7	3.8
1983	74.9	1.6	67.3	35.2	118.9	125.1	111.2	68.4	34.3	3.7
1982	85.8	--	64.5	33.0	115.2	164.1	130.1	82.6	24.1	3.6
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



GRANT COUNTY

GRANT COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	102	85	187	6.8
1984	115	97	212	7.7
1983	90	65	155	5.5
1982	108	77	185	6.5
COUNTY % OF TOTAL (1985) STATE DEATHS	1.9	2.0	1.9	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

GRANT CO.	
MALE	783.6
FEMALE	669.0
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate has decreased to 6.8 deaths per 1000 population, accounting for 1.9% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 52% of all deaths. Observed deaths are less than expected for heart disease and cancer, but 28% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, influenza/pneumonia, cirrhosis and alcoholism.
- Both components of the county infant mortality rate exceed state levels.
- About 95% of resident births occur within the county. Overall, the county experiences 2.8 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

GRANT CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	736	851	86	100.0	100.0
HEART DISEASE	177	321	55	24.0	23.7
MALIGNANT NEOPLASMS	149	179	83	20.2	20.2
ALL ACCIDENT	59	46	128	8.0	10.0
MOTOR VEHICLE	30	23	130	4.1	6.1
ALL OTHER	29	24	121	3.9	3.9
STROKE	45	71	63	6.1	6.1
C.O.P.D.	47	24	134	6.4	4.7
DIABETES MELLITUS	24	15	160	3.3	2.7
SUICIDE	14	12	117	1.9	2.8
INFLUENZA/PNEUMONIA	29	21	138	3.9	2.7
CIRRHOSIS OF LIVER	30	12	250	4.1	1.9
HOMICIDE	7	10	70	1.0	1.9
ALCOHOLISM	13	2	650	1.8	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	11	15	73	1.5	1.6
ATHEROSCLEROSIS	5	12	42	.7	1.1
CONGENITAL ANOMALIES	6	8	75	.8	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS GRANT	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.3	2.9
1 - 4	.2	.8
5- 14	.9	.8
15-24	4.0	3.7
25-34	4.0	4.4
35-44	3.1	4.0
45-54	5.0	6.4
55-64	13.2	13.4
65-74	22.5	22.1
75-84	26.3	24.5
85+	18.6	16.9

INFANT MORTALITY

GRANT COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	7.3	3	5.5	1	1.8
1981	4	6.7	3	5.0	1	1.7
1982	8	14.7	4	7.3	4	7.3
1983	2	4.3	1	2.2	1	2.2
1984	3	6.4	1	2.1	2	4.3
1985	8	15.5	5	9.7	3	5.8
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

GRANT COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Bayard-Urban	55	19	2.9
Bayard-Rural	2	1	2.0
Santa Rita-Urban	0	0	---
Santa Rita-Rural	0	0	---
Central-Urban	42	17	2.5
Central-Rural	2	2	1.0
Hurley-Urban	21	14	1.5
Hurley-Rural	4	1	4.0
Silver City-Urban	238	90	2.6
Silver City-Rural	52	17	3.1
Fort Bayard-All	6	1	6.0
Other & Unspecified	93	25	3.7
COUNTY TOTAL	515	187	2.8
% OF COUNTY RESIDENT BIRTHS:			
Born in County	94.76%	XXXXX	XXXXX
Born in Other NM County	3.69%	XXXXX	XXXXX
Born Out of State	1.55%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT <1 year, NEONATAL <28 days, POSTNEONATAL = 28 days to 1 year

GUADALUPE COUNTY

GUADALUPE COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	4969	4496	4400	4800
% 19 AND YOUNGER	47.2%	38.6%	36.0%	33.8%
% 65 AND OLDER	9.4%	12.2%	10.9%	11.5%
% WHITE NON-HISPANIC	15.0%	16.5%	XXXXX	XXXXX
% WHITE HISPANIC	84.5%	82.4%	XXXXX	XXXXX
% BLACK	0.1%	0.1%	XXXXX	XXXXX
% INDIAN	---	0.2%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	37.9%	27.0%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.5%	0.3%	0.3%	0.3%

- Guadalupe County ranks as the state's 30th most populous county and has a population density of 1.5 persons per square mile, accounting for 0.3% of the state's total population.
- County population projections to the year 2000, project the population to increase to 4,800 persons, continuing to account for 3% of the state population.
- The county birth rate dropped 6% below the state level to 17.7 births per 1000 population in 1985, accounting for 3% of the state's resident births.
- The percentage of low birth weight births decreased by 29% from the prior year and the percentages of births to teen and to single mothers are above state levels - 57% and 17% above, respectively.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

GUADALUPE	YEAR 1983	1984	1985
LIVE BIRTHS	92	78	78
WHITE			
NON-HISPANIC	8	7	9
HISPANIC	80	71	65
BLACK	2	0	0
INDIAN	0	C	1
OTHER	2	0	3
COUNTY % OF TOTAL STATE BIRTHS	.3%	.3%	.3%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	GUADALUPE NUMBER	%	NEW MEXICO %
ALL AGES	78	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	7	9.0%	5.4%
18-19	12	15.4%	9.9%
(15-19)	19	24.4%	15.3%
20-24	28	35.9%	33.6%
25-29	16	20.5%	28.8%
30-34	9	11.5%	15.7%
35-39	5	6.4%	5.5%
=> 40	1	1.3%	0.7%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
GUADALUPE:										
1985	82.1	--	96.4	61.4	144.6	133.3	108.1	60.0	38.2	--
1984	79.3	--	73.0	59.5	94.3	146.1	120.3	79.1	18.0	--
1983	92.6	--	68.8	65.1	74.8	222.2	119.5	71.4	35.7	--
1982	89.5	--	87.0	47.3	149.5	155.6	118.8	71.4	62.5	4.1
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







GUADALUPE COUNTY

GUADALUPE COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	24	12	36	8.2
1984	21	13	34	7.6
1983	14	18	32	7.1
1982	22	18	40	9.1
COUNTY % OF TOTAL (1985) STATE DEATHS	0.4	0.3	0.4	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
GUADALUPE CO.	
MALE	816.7
FEMALE	870.3
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 8.2 deaths per 1000 population accounts for 0.4% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for females and lower for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 51% of all deaths. Observed deaths are less than expected for heart disease and cancer, but twice the expected level for accidents. Observed deaths are higher than expected for diabetes, suicide, cirrhosis, homicide, alcoholism and atherosclerosis.
- Both components of infant mortality are above state levels.
- Over 61% of resident births are delivered in another county. Overall, the county experiences 2.2 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

GUADALUPE CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	162	160	101	100.0	100.0
HEART DISEASE	37	61	61	22.8	23.7
MALIGNANT NEOPLASMS	28	34	82	17.3	20.2
ALL ACCIDENT	17	8	212	10.5	10.0
MOTOR VEHICLE	11	4	275	6.8	6.1
ALL OTHER	6	4	150	3.7	3.9
STROKE	6	14	43	3.7	6.1
C.O.P.D.	4	5	80	2.5	4.7
DIABETES MELLITUS	5	3	167	3.1	2.7
SUICIDE	7	2	350	4.3	2.8
INFLUENZA/PNEUMONIA	3	4	75	1.9	2.7
CIRRHOSIS OF LIVER	3	2	150	1.9	1.9
HOMICIDE	4	2	200	2.5	1.9
ALCOHOLISM	6	.34	1765	3.7	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	2	2	100	1.2	1.6
ATHEROSCLEROSIS	4	2	200	2.5	1.1
CONGENITAL ANOMALIES	--	1	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS GUADALUPE	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	5.9	2.9
1 - 4	1.0	.8
5- 14	1.0	.8
15-24	2.9	3.7
25-34	5.9	4.4
35-44	5.9	4.0
45-54	5.9	6.4
55-64	6.9	13.4
65-74	15.7	22.1
75-84	31.4	24.5
85+	17.6	16.9

INFANT MORTALITY

GUADALUPE COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	1	10.5	1	10.5	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	2	22.5	1	11.2	1	11.2
1983	1	10.9	1	10.9	0	0.0
1984	3	38.5	2	25.6	1	12.8
1985	2	25.6	1	12.8	1	12.8
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

GUADALUPE COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Santa Rosa-Urban	51	19	2.7
Santa Rosa-Rural	12	1	12.0
Vaughn-Urban	5	9	0.6
Vaughn-Rural	4	1	4.0
Other & Unspecified	6	6	1.0
COUNTY TOTAL	78	36	2.2
% OF COUNTY RESIDENT BIRTHS:			
Born in County	38.46%	XXXXX	XXXXX
Born in Other NM County	61.54%	XXXXX	XXXXX
Born Out of State	----	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

HARDING COUNTY

HARDING COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	1348	1090	1000	900
% 19 AND YOUNGER	38.4%	29.5%	29.4%	29.6%
% 65 AND OLDER	14.6%	17.5%	17.8%	21.8%
% WHITE NON-HISPANIC	50.2%	55.1%	XXXXX	XXXXX
% WHITE HISPANIC	49.3%	44.3%	XXXXX	XXXXX
% BLACK	---	---	XXXXX	XXXXX
% INDIAN	0.5%	0.1%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	28.3%	16.9%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.1%	0.1%	0.1%	.0%

- Harding County ranks as the state's least populous county and has a population density of .5 persons per square mile, accounting for 0.1% of New Mexico's population
- The county population is projected to decrease to 900 persons by the year 2000; about a 10% drop in population
- The county birth rate is 15% below the state rate. The county rate of 16 births per 1000 population accounts for 0.1% of state resident births
- Because of the relatively few births, birth characteristics fluctuate, however, the percent low birth weight dropped below state levels, while the percentage of births to single women is generally less than half the state figure



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

HARDING	1983	1984	1985
LIVE BIRTHS	16	11	16
WHITE			
NON-HISPANIC	7	5	7
HISPANIC	9	6	9
BLACK	0	0	0
INDIAN	0	0	0
OTHER	0	0	0
COUNTY % OF TOTAL STATE BIRTHS	.1%	.04%	.1%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	HARDING NUMBER	HARDING %	NEW MEXICO %
ALL AGES	16	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	1	6.3%	5.4%
18-19	1	6.3%	9.9%
(15-19)	2	12.5%	15.3%
20-24	5	31.3%	33.6%
25-29	7	43.8%	28.8%
30-34	2	12.5%	15.7%
35-39	0	0.0%	5.5%
=> 40	0	0.0%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
HARDING										
1985	98.8	--	66.7	58.8	76.9	147.1	411.8	71.4	--	--
1984	65.1	--	--	--	--	80.0	193.5	54.1	66.7	--
1983	92.5	--	71.4	--	176.5	230.8	62.5	105.3	62.5	--
1982	69.4	--	23.8	--	58.8	230.8	93.8	52.6	--	--
NEW MEXICO: 1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







HARDING COUNTY

HARDING COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	7	8	15	15.0
1984	6	3	9	9.2
1983	3	8	11	11.0
1982	7	6	13	13.0
COUNTY % OF TOTAL (1985) STATE DEATHS	0.1	0.2	0.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

HARDING CO.	
MALE	681.9
FEMALE	921.0
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate has increased to 15 deaths per 1000 population, accounting for 0.2% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for females and lower for males.
- Heart disease, cancer and chronic obstructive pulmonary disease (COPD) are the leading causes of death, accounting for 65% of all deaths, as compared to 49% statewide. Observed deaths are almost 20% less than expected for heart disease and cancer.

Due to the relatively small number of births each year, infant mortality rates can fluctuate radically.

- No resident births occur within the county; 19% occur out of state. Overall, the county experiences 1.1 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

HARDING CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	46	53	87	100.0	100.0
HEART DISEASE	17	21	81	37.0	23.7
MALIGNANT NEOPLASMS	10	12	83	21.7	20.2
ALL ACCIDENT	2	2	100	4.3	10.0
MOTOR VEHICLE	1	1	100	2.2	6.1
ALL OTHER	1	1	100	2.2	3.9
STROKE	2	5	40	4.3	6.1
C.O.P.D.	3	2	150	6.5	4.7
DIABETES MELLITUS	2	1	200	4.3	2.7
SUICIDE	1	1	100	2.2	2.8
INFLUENZA/PNEUMONIA	1	1	100	2.2	2.7
CIRRHOSIS OF LIVER	1	1	100	2.2	1.9
HOMICIDE	--	--	--	--	1.9
ALCOHOLISM	--	--	--	--	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	1	--	--	2.2	1.6
ATHEROSCLEROSIS	1	1	100	2.2	1.1
CONGENITAL ANOMALIES	1	--	--	2.2	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS HARDING	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.9	2.9
1 - 4	--	.8
5- 14	--	.8
15-24	2.9	3.7
25-34	--	4.4
35-44	--	4.0
45-54	--	6.4
55-64	5.7	13.4
65-74	11.4	22.1
75-84	34.3	24.5
85+	42.9	16.9

INFANT MORTALITY

HARDING COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	0	0.0	0	0.0	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	1	83.3	1	83.3	0	0.0
1983	0	0.0	0	0.0	0	0.0
1984	0	0.0	0	0.0	0	0.0
1985	1	62.5	0	0.0	1	62.5
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

HARDING COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Roy-All	9	9	1.0
Mosquero-All	2	3	0.7
Other & Unspecified	5	3	1.7
COUNTY TOTAL	16	15	1.1
% OF COUNTY RESIDENT BIRTHS:			
Born in County	----	XXXXX	XXXXX
Born in Other NM County	81.25%	XXXXX	XXXXX
Born Out of State	18.75%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

HIDALGO COUNTY

HIDALGO COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	4734	6049	6200	7600
% 19 AND YOUNGER	44.8%	40.7%	39.2%	39.8%
% 65 AND OLDER	8.8%	9.3%	9.9%	9.7%
% WHITE NON-HISPANIC	39.8%	52.0%	XXXXX	XXXXX
% WHITE HISPANIC	58.8%	47.0%	XXXXX	XXXXX
% BLACK	0.4%	0.2%	XXXXX	XXXXX
% INDIAN	0.4%	0.1%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	21.7%	14.7%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.5%	0.5%	0.4%	0.4%

- Hidalgo County ranks as the state's 27th most populous county and has a population density of 1.8 persons accounting for 0.4% of New Mexico's population.
- The county population is projected to reach 7,600 persons by the year 2000.
- The county birth rate has dropped to 17.3 births per 1000 population, a rate 8% lower than the statewide rate as the county accounts for 0.4% of New Mexico resident births.
- The percentage of births to teens has increased above state levels while the percentage of births to single women though below state levels is increasing. The percent of low birth weight births averages about the state level.


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

HIDALGO	1983	1984	1985
LIVE BIRTHS	144	136	107
WHITE NON-HISPANIC	65	52	43
HISPANIC	77	84	64
BLACK	0	0	0
INDIAN	1	0	0
OTHER	1	0	0
COUNTY % OF TOTAL STATE BIRTHS	.5%	.5%	.4%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	HIDALGO NUMBER	HIDALGO %	NEW MEXICO %
ALL AGES	107	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	11	10.3%	5.4%
18-19	10	9.3%	9.9%
(15-19)	21	19.6%	15.3%
20-24	41	38.3%	33.6%
25-29	31	29.0%	28.8%
30-34	9	8.4%	15.7%
35-39	3	2.8%	5.5%
=> 40	2	1.9%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
HIDALGO:										
1985	76.3	--	72.9	62.9	98.5	156.5	147.6	44.1	12.6	--
1984	96.2	3.1	64.8	36.2	126.2	196.2	140.5	55.8	81.4	3.3
1983	102.3	--	101.5	81.4	144.2	218.0	140.5	64.1	17.4	3.3
1982	99.4	--	100.0	44.4	219.0	174.7	146.3	80.2	34.3	3.2
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

HIDALGO COUNTY

HIDALGO COUNTY DEATHS

			TOTAL	
YEAR	MALE	FEMALE	NUMBER	CRUDE DEATH RATE
1985	19	23	42	6.8
1984	27	19	46	7.2
1983	23	18	41	6.4
1982	16	24	40	6.3
COUNTY % OF TOTAL (1985) STATE DEATHS	0.3	0.6	0.4	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
HIDALGO CO.	
MALE	887.4
FEMALE	737.1
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county death rate decreased to 6.8 deaths per 1000 population, accounting for 0.4% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and chronic obstructive pulmonary disease (COPD), are the leading causes of death, accounting for 52% of all deaths. Observed deaths are lower than expected for heart disease, but 14% higher than expected for cancer and 50% higher for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), influenza/pneumonia, homicide and alcoholism.
- No infant deaths were reported for the county for 1985.
- Over 89% of resident births occur in another county. Overall, the county experiences 2.5 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

HIDALGO CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	192	179	107	100.0	100.0
HEART DISEASE	41	67	61	21.4	23.7
MALIGNANT NEOPLASMS	42	37	114	21.9	20.2
ALL ACCIDENT	15	10	150	7.8	10.0
MOTOR VEHICLE	11	5	220	5.7	6.1
ALL OTHER	4	5	80	2.1	3.9
STROKE	11	15	73	5.7	6.1
C.O.P.D.	16	5	320	8.3	4.7
DIABETES MELLITUS	2	3	67	1.0	2.7
SUICIDE	1	3	33	.5	2.8
INFLUENZA/PNEUMONIA	6	4	150	3.1	2.7
CIRRHOSIS OF LIVER	2	2	100	1.0	1.9
HOMICIDE	5	2	250	2.6	1.9
ALCOHOLISM	3	.34	882	1.6	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	2	4	50	1.0	1.6
ATHEROSCLEROSIS	3	3	100	1.6	1.1
CONGENITAL ANOMALIES	2	2	100	1.0	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS	
	HIDALGO	NEW MEXICO
ALL	100.0	100.0
< 1	3.9	2.9
1 - 4	--	.8
5- 14	.8	.8
15-24	1.6	3.7
25-34	3.1	4.4
35-44	2.3	4.0
45-54	4.7	6.4
55-64	8.5	13.4
65-74	29.5	22.1
75-84	25.6	24.5
85+	20.2	16.9

INFANT MORTALITY

HIDALGO COUNTY	TOTAL INFANT		NEONATAL		POSTNEONATAL	
	Number	Rate	Number	Rate	Number	Rate
1980	2	14.4	1	7.2	1	7.2
1981	2	17.9	1	8.9	1	8.9
1982	1	7.0	1	7.0	0	0.0
1983	2	13.9	1	6.9	1	6.9
1984	3	22.1	3	22.1	0	0.0
1985	0	0.0	0	0.0	0	0.0
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

HIDALGO COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Lordsburg-Urban	63	35	1.8
Lordsburg-Rural	8	0	---
Other & Unspecified	36	7	5.1
COUNTY TOTAL	107	42	2.5
% OF COUNTY RESIDENT BIRTHS:			
Born in County	----	XXXXX	XXXXX
Born in Other NM County	89.72%	XXXXX	XXXXX
Born Out of State	10.28%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

LEA COUNTY

LEA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	49554	55993	65500	77100
% 19 AND YOUNGER	42.6%	37.8%	38.9%	37.9%
% 65 AND OLDER	5.4%	8.0%	8.5%	10.0%
% WHITE NON-HISPANIC	81.2%	73.1%	XXXXX	XXXXX
% WHITE HISPANIC	12.5%	21.2%	XXXXX	XXXXX
% BLACK	5.3%	4.8%	XXXXX	XXXXX
% INDIAN	0.4%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	12.5%	9.0%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	4.9%	4.3%	4.5%	4.2%

- Lea County ranks as the state's 5th most populous county and has a population density of 14.9 persons per square mile accounting for 4.5% of the state's population.
- The county population is projected to reach 77,100 persons by the year 2000, accounting for a decreasing percent of the state population and maintaining a high percentage of population 19 and younger.
- The county birth rate increased slightly for 1985 to 20.9 births per 1000 population remaining 11% higher then the state level, and accounting for 5% of resident births in the state.
- The percentage of low birth weight births decreased but stayed above state levels; the percentage of births to teens increased, remaining 16% higher than for the state. The percent of births to single women is well below the state figure.

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

LEA	1983	1984	1985
LIVE BIRTHS	1596	1368	1367
WHITE			
NON-HISPANIC	875	756	764
HISPANIC	618	529	514
BLACK	87	57	74
INDIAN	9	14	6
OTHER	7	12	9
COUNTY % OF TOTAL STATE BIRTHS	5.8%	5.0%	5.0%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	LEA NUMBER	LEA %	NEW MEXICO %
ALL AGES	1367	100.0%	100.0%
<15	6	0.4%	0.2%
15-17	96	7.0%	5.4%
18-19	144	10.5%	9.9%
(15-19)	240	17.6%	15.3%
20-24	494	36.1%	33.6%
25-29	430	31.5%	28.8%
30-34	152	11.1%	15.7%
35-39	36	2.6%	5.5%
=> 40	9	0.7%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
LEA:										
1985	99.5	2.1	105.3	69.8	159.3	199.4	160.7	59.0	16.5	--
1984	97.9	1.5	82.5	52.4	131.0	193.5	140.4	71.2	17.5	3.0
1983	111.4	2.9	119.2	69.9	198.8	212.2	150.6	66.5	26.2	2.1
1982	122.5	1.2	129.5	73.0	220.0	237.9	160.9	78.5	20.9	4.5
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



LEA COUNTY

LEA COUNTY
DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	210	188	398	6.1
1984	240	173	413	6.3
1983	260	208	468	7.3
1982	246	178	424	6.7
COUNTY % OF TOTAL (1985) STATE DEATHS	3.8	4.5	4.1	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

LEA CO.	
MALE	899.4
FEMALE	802.4
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 6.1 deaths per 1000 population accounts for 4.1% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 57% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 85% higher than expected for accidents. Observed deaths are notably higher than expected for chronic obstructive pulmonary disease (COPD), suicide and alcoholism.
- Infant mortality is slightly below the state level.
- Almost 10% of resident births occur in another state. Overall, the county experiences 3.4 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

LEA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1636	1511	108	100.0	100.0
HEART DISEASE	446	541	82	27.3	23.7
MALIGNANT NEOPLASMS	310	324	96	18.9	20.2
ALL ACCIDENT	179	97	185	10.9	10.0
MOTOR VEHICLE	122	51	239	7.5	6.1
ALL OTHER	57	46	124	3.5	3.9
STROKE	104	115	90	6.4	6.1
C.O.P.D.	68	40	170	4.2	4.7
DIABETES MELLITUS	31	25	124	1.9	2.7
SUICIDE	51	25	204	3.1	2.8
INFLUENZA/PNEUMONIA	25	35	71	1.5	2.7
CIRRHOSIS OF LIVER	23	27	85	1.4	1.9
HOMICIDE	28	22	127	1.7	1.9
ALCOHOLISM	24	4	600	1.5	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	38	35	109	2.3	1.6
ATHEROSCLEROSIS	19	18	106	1.2	1.1
CONGENITAL ANOMALIES	23	18	128	1.4	.9

PERCENTAGE DISTRIBUTION
OF DEATHS BY AGE
1983-1985

AGE	% OF TOTAL DEATHS LEA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.8	2.9
1 - 4	.4	.8
5- 14	.9	.8
15-24	3.3	3.7
25-34	4.6	4.4
35-44	4.0	4.0
45-54	7.3	6.4
55-64	17.0	13.4
65-74	22.4	22.1
75-84	24.2	24.5
85+	12.2	16.9

INFANT MORTALITY

LEA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	26	19.8	17	13.0	9	6.9
1981	29	19.2	18	11.9	11	7.3
1982	19	11.7	9	5.5	10	6.2
1983	22	13.8	12	7.5	10	6.3
1984	14	10.2	6	4.4	8	5.8
1985	13	9.5	7	5.1	6	4.4
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

LEA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Eunice-Urban	62	16	3.9
Eunice-Rural	4	2	2.0
Hobbs-Urban	794	245	3.2
Hobbs-Rural	142	19	7.5
Jal-Urban	47	15	3.1
Jal-Rural	7	2	3.5
Lovington-Urban	229	79	2.9
Lovington-Rural	35	9	3.9
Tatum-Urban	21	5	4.2
Tatum-Rural	4	2	2.0
Other & Unspecified	22	4	5.5
COUNTY TOTAL	1367	398	3.4
% OF COUNTY RESIDENT BIRTHS:			
Born in County	87.78%	XXXXX	XXXXX
Born in Other NM County	2.34%	XXXXX	XXXXX
Born Out of State	9.88%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

LINCOLN COUNTY

LINCOLN COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	7560	10997	13900	18000
% 19 AND YOUNGER	36.5%	31.2%	29.3%	28.3%
% 65 AND OLDER	11.6%	12.2%	12.2%	15.0%
% WHITE NON-HISPANIC	63.5%	72.0%	XXXXX	XXXXX
% WHITE HISPANIC	34.0%	26.5%	XXXXX	XXXXX
% BLACK	0.5%	0.5%	XXXXX	XXXXX
% INDIAN	1.1%	0.6%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	20.7%	13.2%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.7%	0.8%	1.0%	1.0%

- Lincoln County ranks as the state's 22nd most populous county and has a population density of 2 9 persons per square mile One percent of the state's population resides in Lincoln County
- The county population is projected to reach 18,000 persons by the year 2000, continuing to account for 1% of the state population
- The county birth rate, usually at about the state figure, dropped below the state level in 1984 and 1985 to 17 7 births per 1000 population - 6% below the state level and accounting for 0 9% of New Mexico births
- A notable drop in the percent of low birth weight births in 1984 left the county percentage at about the state level in 1985 The percentage of births to single women remains well below state levels, while the percent of births to teen women decreased well below the state levels

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

LINCOLN	YEAR 1983	1984	1985
LIVE BIRTHS	254	246	246
WHITE NON-HISPANIC	162	155	151
HISPANIC	86	85	85
BLACK	0	2	2
INDIAN	6	4	6
OTHER	0	0	2
COUNTY % OF TOTAL STATE BIRTHS	.9%	.9%	.9%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	LINCOLN NUMBER	%	NEW MEXICO %
ALL AGES	246	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	9	3.7%	5.4%
18-19	19	7.7%	9.9%
(15-19)	28	11.4%	15.3%
20-24	96	39.0%	33.6%
25-29	72	29.3%	28.8%
30-34	36	14.6%	15.7%
35-39	9	3.7%	5.5%
=> 40	5	2.0%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
LINCOLN:										
1985	81.0	--	59.2	32.8	95.5	203.0	152.2	63.5	15.9	1.2
1984	85.7	--	79.8	16.5	220.9	170.8	129.4	80.1	20.3	--
1983	94.1	2.0	69.0	29.3	157.9	216.5	136.2	71.4	28.8	9.4
1982	92.5	--	75.9	42.8	149.7	182.3	153.0	64.0	42.6	1.6
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



LINCOLN COUNTY

LINCOLN COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	69	35	104	7.5
1984	64	50	114	8.4
1983	71	44	115	9.1
1982	59	43	102	8.6
COUNTY % OF TOTAL (1985) STATE DEATHS	1.3	0.8	1.1	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
LINCOLN CO.	
MALE	992.7
FEMALE	732.9
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 7.5 deaths per 1000 population accounts for 1.1% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both males and females.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 54% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 65% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), suicide and homicide to name a few.
- No infant deaths were reported for the county for 1985.
- 11% of resident births occur in another county. Overall, the county experiences 2.4 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

LINCOLN CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	334	383	90	100.0	100.0
HEART DISEASE	87	145	60	25.3	23.7
MALIGNANT NEOPLASMS	65	86	76	18.9	20.2
ALL ACCIDENT	33	20	165	9.6	10.0
MOTOR VEHICLE	21	10	210	6.1	6.1
ALL OTHER	12	10	120	3.5	3.9
STROKE	23	31	74	6.7	6.1
C.O.P.D.	21	11	191	6.1	4.7
DIABETES MELLITUS	9	7	129	2.6	2.7
SUICIDE	8	5	160	2.3	2.8
INFLUENZA/PNEUMONIA	11	9	122	3.2	2.7
CIRRHOSIS OF LIVER	8	7	114	2.3	1.9
HOMICIDE	7	5	140	2.0	1.9
ALCOHOLISM	4	.92	435	1.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	4	4	100	1.2	1.6
ATHEROSCLEROSIS	3	5	60	.9	1.1
CONGENITAL ANOMALIES	3	3	100	.9	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS LINCOLN	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.1	2.9
1 - 4	.9	.8
5- 14	1.8	.8
15-24	1.2	3.7
25-34	6.6	4.4
35-44	3.6	4.0
45-54	6.9	6.4
55-64	9.9	13.4
65-74	26.7	22.1
75-84	23.7	24.5
85+	16.5	16.9

INFANT MORTALITY

LINCOLN COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	0	0.0	0	0.0	0	0.0
1981	5	23.1	3	13.9	2	9.3
1982	5	20.9	3	12.6	2	8.4
1983	5	19.7	3	11.8	2	7.9
1984	2	8.1	0	0.0	2	8.1
1985	0	0.0	0	0.0	0	0.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

LINCOLN COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Capitan-All	23	9	2.6
Carrizozo-Urban	17	16	1.1
Carrizozo-Rural	2	2	1.0
Ruiduso-Urban	117	45	2.6
Ruidoso-Rural	12	1	12.0
Ruidoso Downs-All	39	13	3.0
Other & Unspecified	36	18	2.0
COUNTY TOTAL	246	104	2.4
% OF COUNTY RESIDENT BIRTHS:			
Born in County	88.62%	XXXXX	XXXXX
Born in Other NM County	10.98%	XXXXX	XXXXX
Born Out of State	0.41%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

LOS ALAMOS COUNTY

LOS ALAMOS COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	15198	17599	18400	20800
% 19 AND YOUNGER	44.2%	34.0%	29.2%	25.0%
% 65 AND OLDER	2.1%	4.1%	6.5%	16.3%
% WHITE NON-HISPANIC	80.9%	86.1%	XXXXX	XXXXX
% WHITE HISPANIC	17.8%	11.4%	XXXXX	XXXXX
% BLACK	0.4%	0.4%	XXXXX	XXXXX
% INDIAN	0.5%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	2.1%	3.5%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.5%	1.4%	1.3%	1.1%

- Los Alamos County ranks as the state's 18th most populous county and has a population density of 170.4 persons per square mile, accounting for 1.3% of the state's total population.
- The county population is projected to reach 20,800 persons by the year 2000 with the percentage of population 65 and older more than tripling.
- The county birth rate has over the last four years been about half the state rate. The county rate of 11.1 births per 1000 population accounts for 0.7% of all New Mexico resident births.
- Levels of births to teen mothers, and births to single women are well below state levels, while the percentage low birth weight increased approaching the state level.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

LOS ALAMOS	YEAR 1983	1984	1985
LIVE BIRTHS	206	200	204
WHITE			
NON-HISPANIC	160	142	159
HISPANIC	37	53	39
BLACK	3	1	0
INDIAN	1	2	3
OTHER	5	2	3
COUNTY % OF TOTAL STATE BIRTHS	.7%	.7%	.7%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	LOS ALAMOS NUMBER	%	NEW MEXICO %
ALL AGES	204	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	1	0.5%	5.4%
18-19	6	2.9%	9.9%
(15-19)	7	3.4%	15.3%
20-24	46	22.5%	33.6%
25-29	75	36.8%	28.3%
30-34	50	24.5%	15.7%
35-39	23	11.3%	5.5%
=> 40	3	1.5%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
LOS ALAMOS:										
1985	50.0	--	8.8	2.3	16.7	66.1	188.4	83.8	33.0	--
1984	45.9	--	6.1	1.6	19.1	73.7	125.0	80.1	22.5	0.7
1983	48.4	--	15.1	8.4	34.3	98.2	119.8	75.5	15.4	5.1
1982	44.0	--	13.1	3.2	41.9	106.3	94.6	60.5	27.1	1.4
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







LOS ALAMOS COUNTY

LOS ALAMOS COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	48	24	72	3.9
1984	32	27	59	3.1
1983	41	30	71	3.8
1982	37	40	77	4.4
COUNTY % OF TOTAL (1985) STATE DEATHS	0.9	0.6	0.7	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

LOS ALAMOS CO.	
MALE	665.3
FEMALE	546.9
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate increased to 3 9 deaths per 1000 population, accounting for 0 7% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for both females and males.
- Cancer, heart disease and accidents are the leading causes of death, accounting for 63% of all deaths. Observed deaths are lower than expected levels for cancer and heart disease, while the observed number of accident deaths are right at the expected level. Suicide is 50% higher than expected.

Infant mortality jumped notably above the state level.

- Over 88% of resident births occur within the county. Overall, the county experiences 2 8 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

LOS ALAMOS CO. — FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	252	373	68	100.0	100.0
HEART DISEASE	50	126	40	19.8	23.7
MALIGNANT NEOPLASMS	81	88	92	32.1	20.2
ALL ACCIDENT	29	29	100	11.5	10.0
MOTOR VEHICLE	21	16	131	8.3	6.1
ALL OTHER	8	13	62	3.2	3.9
STROKE	8	24	33	3.2	6.1
C.O.P.D.	9	9	100	3.6	4.7
DIABETES MELLITUS	3	6	50	1.2	2.7
SUICIDE	12	8	150	4.8	2.8
INFLUENZA/PNEUMONIA	4	7	57	1.6	2.7
CIRRHOSIS OF LIVER	4	9	44	1.6	1.9
HOMICIDE	5	8	63	2.0	1.9
ALCOHOLISM	3	1	300	1.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	5	6	83	2.0	1.6
ATHEROSCLEROSIS	4	3	133	1.6	1.1
CONGENITAL ANOMALIES	1	4	25	.4	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS LOS ALAMOS	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.4	2.9
1 - 4	--	.8
5- 14	1.5	.8
15-24	2.5	3.7
25-34	1.0	4.4
35-44	5.4	4.0
45-54	8.4	6.4
55-64	23.2	13.4
65-74	21.2	22.1
75-84	18.2	24.5
85+	15.3	16.9

INFANT MORTALITY

LOS ALAMOS COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	1	5.0	1	5.0	0	0.0
1981	1	4.9	1	4.9	0	0.0
1982	2	10.2	1	5.1	1	5.1
1983	2	9.7	2	9.7	0	0.0
1984	1	5.0	0	0.0	1	5.0
1985	4	19.6	1	4.9	3	14.7
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

LOS ALAMOS COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Los Alamos-Urban	190	66	2.9
Los Alamos-Rural	12	2	6.0
White Rock-All	2	4	0.5
Other & Unspecified	0	0	---
COUNTY TOTAL	204	72	2.8
% OF COUNTY RESIDENT BIRTHS:			
Born in County	88.73%	XXXXX	XXXXX
Born in Other NM County	10.29%	XXXXX	XXXXX
Born Out of State	0.98%	XXXXX	XXXXX

***The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

LUNA COUNTY

LUNA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	11706	15585	17500	22500
% 19 AND YOUNGER	43.0%	34.0%	30.0%	28.0%
% 65 AND OLDER	10.2%	16.8%	21.1%	24.4%
% WHITE NON-HISPANIC	50.9%	58.3%	XXXXX	XXXXX
% WHITE HISPANIC	46.5%	39.2%	XXXXX	XXXXX
% BLACK	2.0%	1.6%	XXXXX	XXXXX
% INDIAN	0.1%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	20.5%	19.2%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.2%	1.2%	1.2%	1.2%

- Luna County ranks as the state's 19th most populous county and has a population density of 5.9 persons per square mile, accounting for 1.2% of the state's population.
- The county population is projected to reach 22,500 persons by the year 2000 continuing to account for 1.2 percent of New Mexico's population with a relatively high percentage of persons 65 and older.
- The county birth rate has decreased to 17.3 births per 1000 population, a rate 8% below the state rate and accounting for 1.1% of the state's resident births.
- The percent of births to teen mothers remains 50% higher in the county than statewide while the percents of low birth weight births and births to single women exceed state levels.


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

LUNA	YEAR 1983	1984	1985
LIVE BIRTHS	296	291	302
WHITE NON-HISPANIC	104	93	78
HISPANIC	187	192	216
BLACK	4	5	7
INDIAN	0	0	0
OTHER	1	1	1
COUNTY % OF TOTAL STATE BIRTHS	1.1%	1.1%	1.1%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	LUNA NUMBER	LUNA %	NEW MEXICO %
ALL AGES	302	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	29	9.6%	5.4%
18-19	41	13.6%	9.9%
(15-19)	70	23.2%	15.3%
20-24	99	32.8%	33.6%
25-29	73	24.2%	28.8%
30-34	38	12.6%	15.7%
35-39	18	6.0%	5.5%
=> 40	4	1.3%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
LUNA:										
1985	100.2	--	120.1	84.5	170.8	169.8	150.2	78.2	37.0	--
1984	94.5	1.4	86.2	40.0	180.3	182.0	180.3	50.3	26.2	9.5
1983	97.4	--	112.2	69.9	198.3	175.7	146.6	70.9	31.2	4.8
1982	107.5	4.4	110.0	51.8	228.1	214.0	164.8	76.1	39.0	2.4
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

LUNA COUNTY

LUNA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	129	86	215	12.3
1984	118	61	179	10.8
1983	113	66	179	10.8
1982	111	87	198	12.1
COUNTY % OF TOTAL (1985) STATE DEATHS	2.4	2.1	2.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
LUNA CO.	
MALE	856.0
FEMALE	638.7
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 12.3 deaths per 1000 population accounts for 2.2% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for males, but lower for females.
- Heart disease, cancer and stroke are the leading causes of death, accounting for 61% of all deaths. Observed deaths are lower than expected for heart disease, but 15% higher than expected for cancer and 3% higher for stroke. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, cirrhosis, alcoholism and accidents.
- Infant mortality remains below state levels for 1985.
- Over 20% of resident births occur in another county. Overall, the county experiences 1.4 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

LUNA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	732	692	106	100.0	100.0
HEART DISEASE	211	272	78	28.8	23.7
MALIGNANT NEOPLASMS	173	151	115	23.6	20.2
ALL ACCIDENT	33	29	114	4.5	10.0
MOTOR VEHICLE	28	14	200	3.8	6.1
ALL OTHER	5	16	31	.7	3.9
STROKE	62	60	103	8.5	6.1
C.O.P.D.	47	21	224	6.4	4.7
DIABETES MELLITUS	22	12	183	3.0	2.7
SUICIDE	18	7	257	2.5	2.8
INFLUENZA/PNEUMONIA	23	18	128	3.1	2.7
CIRRHOSIS OF LIVER	13	10	130	1.8	1.9
HOMICIDE	11	6	183	1.5	1.9
ALCOHOLISM	12	1	1200	1.6	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	10	8	125	1.4	1.6
ATHEROSCLEROSIS	7	10	70	1.0	1.1
CONGENITAL ANOMALIES	2	4	50	.3	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS LUNA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	.9	2.9
1 - 4	.2	.8
5- 14	1.0	.8
15-24	2.4	3.7
25-34	1.0	4.4
35-44	1.7	4.0
45-54	4.5	6.4
55-64	13.8	13.4
65-74	30.0	22.1
75-84	29.6	24.5
85+	14.8	16.9

INFANT MORTALITY

LUNA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	2	6.8	2	6.8	0	0.0
1981	3	10.1	2	6.8	1	3.4
1982	4	12.3	3	9.2	1	3.1
1983	3	10.1	1	3.4	2	6.8
1984	0	0.0	0	0.0	0	0.0
1985	2	6.6	2	6.6	0	0.0
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

LUNA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Deming-Urban	190	157	1.2
Deming-Rural	91	43	2.1
Other & Unspecified	21	15	1.4
COUNTY TOTAL	302	215	1.4
% OF COUNTY RESIDENT BIRTHS:			
Born in County	78.81%	XXXXX	XXXXX
Born in Other NM County	20.53%	XXXXX	XXXXX
Born Out of State	0.66%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

MCKINLEY COUNTY

MCKINLEY COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	43208	56449	63300	89200
% 19 AND YOUNGER	51.8%	46.1%	43.9%	42.2%
% 65 AND OLDER	4.4%	4.7%	5.1%	6.3%
% WHITE NON-HISPANIC	17.0%	20.3%	XXXXX	XXXXX
% WHITE HISPANIC	20.0%	12.5%	XXXXX	XXXXX
% BLACK	0.9%	0.6%	XXXXX	XXXXX
% INDIAN	61.3%	65.7%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	33.7%	33.3%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	4.3%	4.3%	4.4%	4.8%

- McKinley County ranks as the state's sixth most populous county and has a population density of 11 6 persons per square mile, accounting for 4 4% of the state's population

- The county population is projected to exceed 89,000 persons by the year 2000, with a relatively small percentage of the population being age 65 and older and a high percentage 19 and younger

- The county's birth rate of 27 2 births per 1000 population is 44% higher than the state rate and accounts for 6 3% of New Mexico resident births

- The percentage of births to single mothers is increasing and is nearly twice the state level while the percentage of births to teens rose above the state level and the percentage of low birth weight births is equal to the percentage statewide


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

MCKINLEY	1983	1984	1985
LIVE BIRTHS	1669	1670	1720
WHITE			
NON-HISPANIC	183	154	159
HISPANIC	198	181	182
BLACK	3	13	8
INDIAN	1280	1317	1370
OTHER	5	5	1
COUNTY % OF TOTAL STATE BIRTHS	6.1%	6.1%	6.3%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	MCKINLEY NUMBER	MCKINLEY %	NEW MEXICO %
ALL AGES	1720	100.0%	100.0%
<15	6	0.3%	0.2%
15-17	115	6.7%	5.4%
18-19	176	10.2%	9.9%
(15-19)	291	16.9%	15.3%
20-24	569	33.1%	33.6%
25-29	432	25.1%	28.8%
30-34	278	16.2%	15.7%
35-39	121	7.0%	5.5%
=> 40	23	1.3%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
		10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
MCKINLEY:										
1985	111.3	1.9	96.4	62.4	149.0	176.7	153.3	106.2	57.3	0.3
1984	110.0	0.6	72.3	44.1	119.8	191.9	144.1	119.9	56.4	8.9
1983	111.0	2.8	83.4	53.6	133.5	182.9	150.3	108.9	63.4	8.3
1982	113.6	1.2	84.1	50.0	141.7	174.4	167.4	121.4	54.9	8.9
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

MCKINLEY COUNTY

MCKINLEY COUNTY
DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	233	122	355	5.6
1984	227	130	357	5.8
1983	218	140	358	5.9
1982	194	139	333	5.6
COUNTY % OF TOTAL (1985) STATE DEATHS	4.3	2.9	3.7	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

MCKINLEY CO.	
MALE	1101.9
FEMALE	808.3
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 5.6 deaths per 1000 population accounts for 3.7% of all New Mexico resident deaths. Adjusted rates show notably higher than state levels of mortality for both females and males.
- Accidents, cancer and heart disease are the leading causes of death, accounting for 47% of all deaths. Observed deaths are lower than expected for cancer and heart disease, but nearly three and a half times higher than expected for accidents. Observed deaths are higher than expected for diabetes, suicide, influenza/pneumonia, cirrhosis and alcoholism.
- Postneonatal mortality is higher than statewide.
- Over 90% of resident births occur within the county. Overall, the county experiences 4.8 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

MCKINLEY CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1417	1100	129	100.0	100.0
HEART DISEASE	166	361	46	11.7	23.7
MALIGNANT NEOPLASMS	185	222	83	13.1	20.2
ALL ACCIDENT	320	92	348	22.6	10.0
MOTOR VEHICLE	191	51	375	13.5	6.1
ALL OTHER	129	41	315	9.1	3.9
STROKE	45	77	58	3.2	6.1
C.O.P.D.	16	27	59	1.1	4.7
DIABETES MELLITUS	26	17	153	1.8	2.7
SUICIDE	52	22	236	3.7	2.8
INFLUENZA/PNEUMONIA	54	25	216	3.8	2.7
CIRRHOSIS OF LIVER	33	19	174	2.3	1.9
HOMICIDE	40	22	182	2.8	1.9
ALCOHOLISM	63	3	2100	4.4	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	38	41	93	2.7	1.6
ATHEROSCLEROSIS	3	12	25	.2	1.1
CONGENITAL ANOMALIES	24	21	114	1.7	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS MCKINLEY	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	5.5	2.9
1 - 4	2.4	.8
5- 14	.8	.8
15-24	9.4	3.7
25-34	9.2	4.4
35-44	8.6	4.0
45-54	8.4	6.4
55-64	11.7	13.4
65-74	15.0	22.1
75-84	15.9	24.5
85+	13.1	16.9

INFANT MORTALITY

MCKINLEY COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	22	13.2	10	6.0	12	7.2
1981	14	8.4	4	2.4	10	6.0
1982	25	14.5	12	7.0	13	7.5
1983	23	13.8	15	9.0	8	4.8
1984	17	10.2	7	4.2	10	6.0
1985	19	11.0	8	4.7	11	6.4
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

MCKINLEY COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Crownpoint-All	113	24	4.7
Gallup-Urban	483	132	3.7
Gallup-Rural	25	21	1.2
Rehoboth-All	18	7	2.6
Zuni-All	181	29	6.2
Other & Unspecified	900	142	6.3
COUNTY TOTAL	1720	355	4.8
% OF COUNTY RESIDENT BIRTHS:			
Born in County	90.76%	XXXXX	XXXXX
Born in Other NM County	2.91%	XXXXX	XXXXX
Born Out of State	6.34%	XXXXX	XXXXX

****The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.*

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

MORA COUNTY

MORA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	4673	4205	4600	5400
% 19 AND YOUNGER	47.0%	37.4%	32.6%	29.9%
% 65 AND OLDER	11.9%	14.8%	15.2%	16.2%
% WHITE NON-HISPANIC	5.1%	12.9%	XXXXX	XXXXX
% WHITE HISPANIC	94.6%	86.4%	XXXXX	XXXXX
% BLACK	0.0%	0.0%	XXXXX	XXXXX
% INDIAN	0.0%	0.2%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	57.3%	36.9%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.5%	0.3%	0.3%	0.3%

- Mora County ranks as the state's 29th most populous county and has a population density of 2.4 persons per square mile, accounting for 0.3% of the state's population.
- The county population is projected to be at about 5,400 persons by the year 2000, with an increasing percentage of persons 65 and older. A large percentage of the county's population is Hispanic.
- The county's birth rate has been below state levels and is presently 17% below the state rate with a rate of 15.7 births per 1000 population accounting for 0.3% of New Mexico resident births.
- The percentages of low birth weight births, births to teen mothers and births to single women increased notably for 1985. The percentage of low birth weight births is more than twice the state level.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

MORA	YEAR		
	1983	1984	1985
LIVE BIRTHS	77	76	72
WHITE			
NON-HISPANIC	5	7	6
HISPANIC	71	69	65
BLACK	0	0	0
INDIAN	1	0	1
OTHER	0	0	0
COUNTY % OF TOTAL STATE BIRTHS	.3%	.3%	.3%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	MORA NUMBER	MORA %	NEW MEXICO %
ALL AGES	72	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	3	4.2%	5.4%
18-19	12	16.7%	9.9%
(15-19)	15	20.8%	15.3%
20-24	26	36.1%	33.6%
25-29	20	27.8%	28.8%
30-34	8	11.1%	15.7%
35-39	3	4.2%	5.5%
=) 40	0	0.0%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
MORA:										
1985	82.1	--	83.8	30.3	150.0	130.0	168.1	51.0	21.6	--
1984	84.0	--	34.5	16.9	72.3	205.9	129.6	86.1	38.5	4.7
1983	90.4	--	65.0	35.7	128.2	226.6	124.2	63.4	30.6	4.9
1982	98.3	--	83.3	61.0	131.6	254.0	100.0	79.1	31.3	10.1
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







MORA COUNTY

MORA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	18	13	31	6.7
1984	33	7	40	8.4
1983	25	19	44	9.8
1982	21	20	41	9.3
COUNTY % OF TOTAL (1985) STATE DEATHS	0.3	0.3	0.3	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
MORA CO.	
MALE	781.2
FEMALE	590.2
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate has dropped to 6.7 deaths per 1000 population, accounting for 0.3% of all New Mexico resident deaths. Adjusted rates show a lower than state level of mortality for both males and females.
- Cancer, heart disease and accidents are the leading causes of deaths, accounting for 55% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 75% higher than expected for accidents.
- Due to relatively small number of births, infant mortality rates fluctuate, but were well above state levels for 1985.
- Over 97% of resident births occur in another county. Overall, the county experiences 2.3 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

MORA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	150	187	80	100.0	100.0
HEART DISEASE	30	74	41	20.0	23.7
MALIGNANT NEOPLASMS	38	39	97	25.3	20.2
ALL ACCIDENT	14	8	175	9.3	10.0
MOTOR VEHICLE	6	4	150	4.0	6.1
ALL OTHER	8	4	200	5.3	3.9
STROKE	7	17	41	4.7	6.1
C.O.P.D.	6	6	100	4.0	4.7
DIABETES MELLITUS	6	3	200	4.0	2.7
SUICIDE	6	2	300	4.0	2.8
INFLUENZA/PNEUMONIA	7	5	140	4.7	2.7
CIRRHOSIS OF LIVER	7	2	350	4.7	1.9
HOMICIDE	7	2	350	4.7	1.9
ALCOHOLISM	1	.31	323	.7	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	1	2	50	.7	1.6
ATHEROSCLEROSIS	--	3	--	--	1.1
CONGENITAL ANOMALIES	--	1	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS MORA	NEW MEXICO
ALL	100.0	100.0
< 1	4.3	2.9
1 - 4	.9	.8
5- 14	1.7	.8
15-24	.9	3.7
25-34	3.5	4.4
35-44	4.3	4.0
45-54	7.0	6.4
55-64	8.7	13.4
65-74	21.7	22.1
75-84	20.9	24.5
85+	26.1	16.9

INFANT MORTALITY

MORA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	1	11.1	0	0.0	1	11.1
1981	0	0.0	0	0.0	0	0.0
1982	0	0.0	0	0.0	0	0.0
1983	0	0.0	0	0.0	0	0.0
1984	1	13.2	0	0.0	1	13.2
1985	4	55.6	3	41.7	1	13.9
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

MORA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Wagon Mound-All	8	7	1.1
Mora-All	25	8	3.1
Cleveland-All	8	2	4.0
Other & Unspecified	31	14	2.2
COUNTY TOTAL	72	31	2.3
% OF COUNTY RESIDENT BIRTHS:			
Born in County	2.78%	XXXXX	XXXXX
Born in Other NM County	97.22%	XXXXX	XXXXX
Born Out of State	----	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from these obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

OTERO COUNTY

OTERO COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	41097	44665	49400	62500
% 19 AND YOUNGER	44.9%	37.3%	35.1%	34.9%
% 65 AND OLDER	4.1%	6.6%	7.3%	9.4%
% WHITE NON-HISPANIC	67.0%	67.0%	XXXXX	XXXXX
% WHITE HISPANIC	23.7%	21.3%	XXXXX	XXXXX
% BLACK	4.5%	4.8%	XXXXX	XXXXX
% INDIAN	3.9%	4.8%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	12.3%	11.6%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	4.0%	3.4%	3.4%	3.4%

- Otero County ranks as the state's 9th most populous county and has a population density of 7.4 persons per 1000 population accounting for 3.4% of the state's population.
- The county population is projected to reach 62,500 people by the year 2000 with a 42% increase in the percentage of persons 65 and older.
- The county birth rate remains 20% higher than the state rate. At a rate of 22.6 births per 1000 population the county accounts for 4.1% of all New Mexico resident births.
- Decreased percentages of low birth weight births, births to single mothers and births to teen mothers leave county levels below statewide levels.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

OTERO	1983	1984	1985
LIVE BIRTHS	1163	1087	1114
WHITE			
NON-HISPANIC	634	598	612
HISPANIC	279	279	275
BLACK	98	83	96
INDIAN	116	93	88
OTHER	36	34	43
COUNTY % OF TOTAL STATE BIRTHS	4.2%	4.0%	4.1%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	OTERO NUMBER	OTERO %	NEW MEXICO %
ALL AGES	1114	100.0%	100.0%
<15	2	0.2%	0.2%
15-17	38	3.4%	5.4%
18-19	118	10.6%	9.9%
(15-19)	156	14.0%	15.3%
20-24	475	42.6%	33.6%
25-29	310	27.8%	28.8%
30-34	130	11.7%	15.7%
35-39	36	3.2%	5.5%
=> 40	5	0.4%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
OTERO:										
1985	96.5	1.1	81.9	33.2	155.5	206.1	140.7	64.9	21.1	--
1984	91.3	1.5	70.2	31.8	130.6	179.6	137.6	67.2	24.3	1.2
1983	100.1	--	85.1	46.7	145.1	209.6	130.2	74.8	22.8	1.2
1982	93.0	1.5	93.4	32.9	162.4	180.3	129.3	74.8	17.7	2.0
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







OTERO COUNTY

OTERO COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	156	111	267	5.4
1984	146	127	273	5.5
1983	147	113	260	5.4
1982	176	115	291	6.3
COUNTY % OF TOTAL (1985) STATE DEATHS	2.8	2.7	2.8	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

OTERO CO.	
MALE	803.5
FEMALE	752.8
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 5.4 deaths per 1000 population accounts for 2.8% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for females but lower for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 54% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 31% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), suicide, cirrhosis, alcoholism and atherosclerosis.
- Infant mortality is presently 10% higher than the statewide rate due to a higher postneonatal component.
- Almost 88% of resident births occur within the county. Overall, the county experience 4.2 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

OTERO CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1064	1048	102	100.0	100.0
HEART DISEASE	261	364	72	24.5	23.7
MALIGNANT NEOPLASMS	213	221	96	20.0	20.2
ALL ACCIDENT	105	80	131	9.9	10.0
MOTOR VEHICLE	67	44	152	6.3	6.1
ALL OTHER	38	36	106	3.6	3.9
STROKE	67	77	87	6.3	6.1
C.O.P.D.	51	27	189	4.8	4.7
DIABETES MELLITUS	18	17	106	1.7	2.7
SUICIDE	30	20	150	2.8	2.8
INFLUENZA/PNEUMONIA	28	24	117	2.6	2.7
CIRRHOSIS OF LIVER	31	19	163	2.9	1.9
HOMICIDE	22	19	116	2.1	1.9
ALCOHOLISM	22	3	733	2.1	1.7
CERTAIN CONDITIONS					
IN PERINATAL PERIOD	25	24	104	2.3	1.6
ATHEROSCLEROSIS	17	12	142	1.6	1.1
CONGENITAL ANOMALIES	7	13	54	.7	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS OTERO	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	4.2	2.9
1 - 4	.6	.8
5- 14	1.1	.8
15-24	4.2	3.7
25-34	4.2	4.4
35-44	4.7	4.0
45-54	7.0	6.4
55-64	15.0	13.4
65-74	22.2	22.1
75-84	23.2	24.5
85+	13.4	16.9

INFANT MORTALITY

OTERO COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	12	12.9	8	8.6	4	4.3
1981	9	9.1	7	7.1	2	2.0
1982	14	13.1	6	5.6	8	7.5
1983	8	6.9	3	2.6	5	4.3
1984	13	12.0	7	6.4	6	5.5
1985	13	11.7	5	4.5	8	7.2
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

OTERO COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Alamogordo-Urban	634	176	3.6
Alamogordo-Rural	65	15	4.3
Mescalero-All	71	15	4.7
Tularosa-Urban	51	18	2.8
Tularosa-Rural	19	7	2.7
Cloudcroft-All	12	7	1.7
Holloman AFB (Military)	181	4	45.3
Other & Unspecified	81	25	3.2
COUNTY TOTAL	1114	267	4.2
% OF COUNTY RESIDENT BIRTHS:			
Born in County	87.97%	XXXXX	XXXXX
Born in Other NM County	9.25%	XXXXX	XXXXX
Born Out of State	2.78%	XXXXX	XXXXX

****The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.*

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

QUAY COUNTY

QUAY COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	10903	10577	11700	11800
% 19 AND YOUNGER	38.8%	33.6%	29.3%	28.0%
% 65 AND OLDER	12.5%	15.7%	16.4%	20.3%
% WHITE NON-HISPANIC	61.9%	62.2%	XXXXX	XXXXX
% WHITE HISPANIC	36.0%	35.3%	XXXXX	XXXXX
% BLACK	1.1%	1.3%	XXXXX	XXXXX
% INDIAN	0.2%	0.4%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	19.3%	15.3%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.1%	0.8%	0.8%	0.6%

- Quay County ranks as the state's 24th most populous county and has a population density of 4.1 persons per square mile, accounting for 0.8% of the state's total population.
- The county population is projected to reach 11,800 persons by the year 2000, accounting for a decreasing percentage of the state population.
- The county birth rate of 15 births per 1000 population is 21% below the state level and accounts for 0.6% of New Mexico resident births.
- The percentage of low birth weight births is below the state level, while the percentage of births to teen women is twice the state percentage, as the fertility rate for county women age 18-19 is over twice that for women 18-19 statewide.


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

QUAY	1983	1984	1985
	YEAR		
LIVE BIRTHS	165	183	176
WHITE			
NON-HISPANIC	81	78	77
HISPANIC	83	100	96
BLACK	1	2	1
INDIAN	0	0	2
OTHER	0	3	0
COUNTY % OF TOTAL STATE BIRTHS	.6%	.7%	.6%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	QUAY NUMBER	QUAY %	NEW MEXICO %
ALL AGES	176	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	17	9.7%	5.4%
18-19	38	21.6%	9.9%
(15-19)	55	31.3%	15.3%
20-24	54	30.7%	33.6%
25-29	45	25.6%	28.8%
30-34	16	9.1%	15.7%
35-39	5	2.8%	5.5%
=> 40	1	0.6%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
QUAY:										
1985	91.8	--	181.5	91.9	322.0	178.2	148.5	39.7	12.4	--
1984	87.0	2.2	97.6	51.4	198.6	199.4	138.5	47.4	9.8	1.9
1983	77.9	2.2	79.3	38.5	169.0	171.6	107.4	52.6	32.6	3.7
1982	97.8	2.2	113.1	78.9	188.4	179.9	138.8	96.9	26.8	--
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

QUAY COUNTY

QUAY COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	64	51	115	9.8
1984	64	54	118	11.1
1983	82	48	130	12.1
1982	74	37	111	10.5
COUNTY % OF TOTAL (1985) STATE DEATHS	1.2	1.2	1.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
QUAY CO.	
MALE	880.9
FEMALE	781.6
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 9.8 deaths per 1000 population accounts for 1.2% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both males and females.
- Heart disease, cancer and stroke are the leading causes of death, accounting for 65% of all deaths. Observed deaths are lower than expected for stroke, equal to expected for cancer and higher than expected (8%) for heart disease. Observed deaths are higher than expected for accidents, chronic obstructive pulmonary disease (COPD), diabetes, influenza/pneumonia, cirrhosis and alcoholism.
- No infant deaths were reported for the county for 1985.
- Over 11% of resident births occur in another county. Overall, the county experiences 1.5 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

QUAY CO. — FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	511	461	111	100.0	100.0
HEART DISEASE	197	182	108	38.6	23.7
MALIGNANT NEOPLASMS	95	95	100	18.6	20.2
ALL ACCIDENT	34	20	170	6.7	10.0
MOTOR VEHICLE	15	9	167	2.9	6.1
ALL OTHER	19	10	190	3.7	3.9
STROKE	39	41	95	7.6	6.1
C.O.P.D.	22	14	157	4.3	4.7
DIABETES MELLITUS	11	8	138	2.2	2.7
SUICIDE	3	5	60	.6	2.8
INFLUENZA/PNEUMONIA	15	12	125	2.9	2.7
CIRRHOSIS OF LIVER	13	10	130	2.5	1.9
HOMICIDE	2	4	50	.4	1.9
ALCOHOLISM	6	1	600	1.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	3	5	60	.6	1.6
ATHEROSCLEROSIS	2	7	29	.4	1.1
CONGENITAL ANOMALIES	2	3	67	.4	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS QUAY	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	1.4	2.9
1 - 4	1.1	.8
5- 14	.8	.8
15-24	3.0	3.7
25-34	1.9	4.4
35-44	1.9	4.0
45-54	4.9	6.4
55-64	9.9	13.4
65-74	27.2	22.1
75-84	32.7	24.5
85+	15.1	16.9

INFANT MORTALITY

QUAY COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	22.3	3	16.8	1	5.6
1981	2	12.7	1	6.4	1	6.4
1982	0	0.0	0	0.0	0	0.0
1983	2	12.1	1	6.1	1	6.1
1984	2	10.9	0	0.0	2	10.9
1985	0	0.0	0	0.0	0	0.0
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

QUAY COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Tucumcari-Urban	124	87	1.4
Tucumcari-Rural	21	13	1.6
San Jon-All	8	1	8.0
Logan-All	14	8	1.8
Other & Unspecified	9	6	1.5
COUNTY TOTAL	176	115	1.5
% OF COUNTY RESIDENT BIRTHS:			
Born in County	82.39%	XXXXX	XXXXX
Born in Other NM County	11.36%	XXXXX	XXXXX
Born Out of State	6.25%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.*

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

RIO ARRIBA COUNTY

RIO ARRIBA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	25170	29282	32400	42300
% 19 AND YOUNGER	48.6%	40.3%	38.4%	37.1%
% 65 AND OLDER	7.4%	8.2%	7.9%	8.7%
% WHITE NON-HISPANIC	5.9%	14.5%	XXXXX	XXXXX
% WHITE HISPANIC	82.2%	73.3%	XXXXX	XXXXX
% BLACK	0.2%	0.1%	XXXXX	XXXXX
% INDIAN	10.9%	11.4%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	34.3%	25.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	2.5%	2.2%	2.2%	2.3%

- Rio Arriba County ranks as the state's 13th most populous county and has a population density of 5.5 persons per square mile, accounting for 2.2% of the state's total population.
- The county population is projected to exceed 42,000 persons by the year 2000, with over a third of the population 19 years of age and younger.
- The county birth rate has increased to over 24% higher than the state rate. The county rate of 23.4 births per 1000 population accounts for 2.8% of New Mexico resident births.
- Although decreasing, the percentage of low birth weight births is 36% higher than statewide. The percentage of births to teens is 31% higher than state levels, while the percentage of births to single mothers is 38% higher.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

RIO ARRIBA	1983	1984	1985
LIVE BIRTHS	676	733	758
WHITE			
NON-HISPANIC	62	67	58
HISPANIC	484	527	538
BLACK	2	3	6
INDIAN	127	136	156
OTHER	1	0	0
COUNTY % OF TOTAL STATE BIRTHS	2.5%	2.7%	2.8%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	RIO ARRIBA NUMBER	RIO ARRIBA %	NEW MEXICO %
ALL AGES	758	100.0%	100.0%
<15	1	0.1%	0.2%
15-17	53	7.0%	5.4%
18-19	100	13.2%	9.9%
(15-19)	153	20.2%	15.3%
20-24	258	34.0%	33.6%
25-29	184	24.3%	28.8%
30-34	99	13.1%	15.7%
35-39	57	7.5%	5.5%
=> 40	6	0.8%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP	15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
RIO ARRIBA:										
1985	99.4	0.7	110.6	65.4	174.8	163.3	133.0	83.5	48.1	0.6
1984	97.6	2.5	86.7	46.7	163.7	155.1	164.6	87.6	36.1	2.6
1983	91.9	1.9	79.9	39.0	158.6	157.5	138.8	85.9	32.5	5.3
1982	92.3	--	77.9	39.7	151.6	155.5	151.6	77.8	32.8	7.5
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







RIO ARRIBA COUNTY

RIO ARRIBA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	162	106	268	8.3
1984	153	81	234	7.3
1983	136	93	229	7.3
1982	153	79	232	7.5
COUNTY % OF TOTAL (1985) STATE DEATHS	3.0	2.5	2.8	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
RIO ARRIBA CO.	
MALE	1081.5
FEMALE	884.8
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 8.3 deaths per 1000 population accounts for 2.8% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for half of all deaths. Observed deaths are lower than expected for heart disease and cancer, but nearly 2 1/2 times the expected level for accidents. Observed deaths are notably higher than expected for diabetes, suicide, cirrhosis, homicide and alcoholism.
- The postneonatal component of infant mortality is higher than statewide.
- Over 60% of resident births occur in another county. Overall, the county experiences 2.8 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

RIO ARRIBA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	883	823	107	100.0	100.0
HEART DISEASE	174	302	58	19.7	23.7
MALIGNANT NEOPLASMS	145	171	85	16.4	20.2
ALL ACCIDENT	125	51	245	14.2	10.0
MOTOR VEHICLE	75	27	278	8.5	6.1
ALL OTHER	50	24	208	5.7	3.9
STROKE	49	66	74	5.5	6.1
C.O.P.D.	32	23	139	3.6	4.7
DIABETES MELLITUS	32	13	246	3.6	2.7
SUICIDE	37	13	285	4.2	2.8
INFLUENZA/PNEUMONIA	25	21	119	2.8	2.7
CIRRHOSIS OF LIVER	30	13	231	3.4	1.9
HOMICIDE	25	11	227	2.8	1.9
ALCOHOLISM	28	2	1400	3.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	15	17	88	1.7	1.6
ATHEROSCLEROSIS	6	11	55	.7	1.1
CONGENITAL ANOMALIES	8	9	89	.9	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS RIO ARRIBA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	4.0	2.9
1 - 4	.8	.8
5- 14	.7	.8
15-24	7.8	3.7
25-34	6.6	4.4
35-44	5.2	4.0
45-54	5.9	6.4
55-64	11.4	13.4
65-74	18.5	22.1
75-84	24.4	24.5
85+	14.9	16.9

INFANT MORTALITY

RIO ARRIBA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	2	3.0	1	1.5	1	1.5
1981	8	12.1	4	6.0	4	6.0
1982	9	13.9	7	10.8	2	3.1
1983	12	17.8	7	10.4	5	7.4
1984	9	12.3	6	8.2	3	4.1
1985	8	10.6	4	5.3	4	5.3
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

RIO ARRIBA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Espanola-Urban (Fairview incl.)	135	44	3.1
Espanola-Rural (Fairview incl.)	116	18	6.4
Chama-All	32	15	2.1
Tierra Amarilla-All	12	7	1.7
Abiquiu-All	10	8	1.3
El Rito-All	22	6	3.7
Other & Unspecified	431	170	2.5
COUNTY TOTAL	758	268	2.8
% OF COUNTY RESIDENT BIRTHS:			
Born in County	38.79%	XXXXX	XXXXX
Born in Other NM County	60.29%	XXXXX	XXXXX
Born Out of State	0.92%	XXXXX	XXXXX

***The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

ROOSEVELT COUNTY

ROOSEVELT COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	16479	15695	16500	18200
% 19 AND YOUNGER	38.8%	34.0%	29.9%	31.3%
% 65 AND OLDER	9.9%	12.9%	13.4%	12.1%
% WHITE NON-HISPANIC	82.4%	75.3%	XXXXX	XXXXX
% WHITE HISPANIC	15.3%	21.4%	XXXXX	XXXXX
% BLACK	0.7%	1.3%	XXXXX	XXXXX
% INDIAN	0.6%	0.7%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	21.5%	21.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.6%	1.2%	1.1%	1.0%

- Roosevelt County ranks as the state's 20th most populous county and has a population density of 6.7 persons per square mile, accounting for 1.1% of the state's total population.
- The county population is projected to reach 18,200 persons by the year 2000 accounting for a decreased percent of New Mexico's population.
- An increase in the county birth rate has left the rate 8% lower than the state rate. The county rate of 17.3 births per 1000 population accounts for 1% of the state's total resident births.
- The percentage of births to teen women has increased above state levels, while percentages of low birth weight births and births to single women remain below state levels.

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

ROOSEVELT	1983	1984	1985
LIVE BIRTHS	294	268	286
WHITE NON-HISPANIC	183	155	171
HISPANIC	103	106	103
BLACK	3	4	4
INDIAN	2	3	4
OTHER	3	0	4
COUNTY % OF TOTAL STATE BIRTHS	1.1%	1.0%	1.0%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	ROOSEVELT NUMBER	ROOSEVELT %	NEW MEXICO %
ALL AGES	286	100.0%	100.0%
<15	2	0.7%	0.2%
15-17	16	5.6%	5.4%
18-19	37	12.9%	9.9%
(15-19)	53	18.5%	15.3%
20-24	94	32.9%	33.6%
25-29	74	25.9%	28.8%
30-34	44	15.4%	15.7%
35-39	13	4.5%	5.5%
=> 40	6	2.1%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
ROOSEVELT:										
1985	70.5	4.0	87.7	44.8	149.8	93.3	73.6	87.5	25.8	--
1984	65.9	--	38.8	31.2	44.2	99.0	113.9	48.7	17.6	5.3
1983	72.7	--	49.3	51.6	47.8	95.5	135.0	75.7	35.6	5.3
1982	84.4	--	73.0	48.6	90.3	125.0	119.9	75.9	31.5	2.7
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



ROOSEVELT COUNTY

ROOSEVELT COUNTY
DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	65	50	115	7.0
1984	78	73	151	9.0
1983	80	43	123	7.4
1982	67	52	119	7.3
COUNTY % OF TOTAL (1985) STATE DEATHS	1.2	1.2	1.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

ROOSEVELT CO.	
MALE	895.8
FEMALE	684.8
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 7 deaths per 1000 population, accounting for 1.2% of all New Mexico resident deaths. Adjusted rates are lower than state levels of mortality for females but higher for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 59% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 74% higher than expected for accidents. Observed deaths are higher than expected for several causes including diabetes and influenza/pneumonia.
- Both components of infant mortality exceed the state rates.
- Over 13% of resident births occur in another county. Overall, the county experiences 2.5 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

ROOSEVELT CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	559	584	96	100.0	100.0
HEART DISEASE	162	227	71	29.0	23.7
MALIGNANT NEOPLASMS	112	120	93	20.0	20.2
ALL ACCIDENT	54	31	174	9.7	10.0
MOTOR VEHICLE	33	16	206	5.9	6.1
ALL OTHER	21	15	140	3.8	3.9
STROKE	45	52	87	8.1	6.1
C.O.P.D.	11	17	65	2.0	4.7
DIABETES MELLITUS	14	10	140	2.5	2.7
SUICIDE	8	7	114	1.4	2.8
INFLUENZA/PNEUMONIA	20	16	125	3.6	2.7
CIRRHOSIS OF LIVER	6	8	75	1.1	1.9
HOMICIDE	3	6	50	.5	1.9
ALCOHOLISM	--	1	--	--	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	5	6	83	.9	1.6
ATHEROSCLEROSIS	11	9	122	2.0	1.1
CONGENITAL ANOMALIES	9	4	225	1.6	.9

PERCENTAGE DISTRIBUTION
OF DEATHS BY AGE
1983-1985

AGE	% OF TOTAL DEATHS ROOSEVELT	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.1	2.9
1 - 4	.8	.8
5- 14	1.0	.8
15-24	2.8	3.7
25-34	2.6	4.4
35-44	1.8	4.0
45-54	4.1	6.4
55-64	13.3	13.4
65-74	21.3	22.1
75-84	27.4	24.5
85+	22.8	16.9

INFANT MORTALITY

ROOSEVELT COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	13.8	3	10.3	1	3.4
1981	4	13.7	3	10.3	1	3.4
1982	5	15.2	2	6.1	3	9.1
1983	2	6.8	2	6.8	0	0.0
1984	2	7.5	2	7.5	0	0.0
1985	4	14.0	2	7.0	2	7.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

ROOSEVELT COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Portales-Urban	197	92	2.1
Portales-Rural	66	13	5.1
Other & Unspecified	23	10	2.3
COUNTY TOTAL	286	115	2.5
% OF COUNTY RESIDENT BIRTHS:			
Born in County	84.27%	XXXXX	XXXXX
Born in Other NM County	13.99%	XXXXX	XXXXX
Born Out of State	1.75%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

SANDOVAL COUNTY

SANDOVAL COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	17492	34799	44100	68900
% 19 AND YOUNGER	48.3%	37.5%	35.3%	34.8%
% 65 AND OLDER	7.2%	9.8%	11.8%	12.6%
% WHITE NON-HISPANIC	36.8%	43.9%	XXXXX	XXXXX
% WHITE HISPANIC	23.6%	27.3%	XXXXX	XXXXX
% BLACK	0.1%	0.9%	XXXXX	XXXXX
% INDIAN	38.9%	27.3%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	37.5%	14.4%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.7%	2.7%	3.0%	3.7%

- Sandoval County ranks as the state's 10th most populous county and has a population density of 11.9 persons per square mile.
- The county population is projected to approach 69,000 persons by the year 2000, a 98% increase over 1980.
- The county birth rate remains at a level over 17% higher than the state rate. The county rate of 22.2 births per 1000 population accounts for 3.6% of New Mexico resident births.
- The percentage of births to single women is 7% higher than statewide, but percentages of low birth weight births and births to teen women are below state levels.

BIRTH RATES PER 1000 POPULATION



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SANDOVAL	1983	1984	1985
LIVE BIRTHS	796	899	981
WHITE NON-HISPANIC	255	305	360
HISPANIC	209	235	263
BLACK	3	11	17
INDIAN	324	338	331
OTHER	5	10	10
COUNTY % OF TOTAL STATE BIRTHS	2.9%	3.3%	3.6%

PERCENT LOW BIRTH WEIGHT



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	SANDOVAL NUMBER	SANDOVAL %	NEW MEXICO %
ALL AGES	981	100.0%	100.0%
<15	2	0.2%	0.2%
15-17	36	3.7%	5.4%
18-19	69	7.0%	9.9%
(15-19)	105	10.7%	15.3%
20-24	289	29.5%	33.6%
25-29	330	33.6%	28.8%
30-34	194	19.8%	15.7%
35-39	49	5.0%	5.5%
=> 40	12	1.2%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
SANDOVAL:										
1985	98.1	1.1	55.4	32.2	88.8	170.4	206.8	108.0	28.9	--
1984	112.0	--	67.7	35.8	124.8	204.3	183.0	110.7	54.8	5.9
1983	98.4	1.7	63.3	32.9	117.6	179.8	172.7	90.6	34.6	5.3
1982	93.0	--	67.0	40.7	114.0	175.2	145.7	84.1	40.5	4.7
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44

BIRTHS TO TEEN MOTHERS



BIRTHS TO SINGLE MOTHERS



LEVEL OF PRENATAL CARE 1985



SANDOVAL COUNTY

SANDOVAL COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	186	126	312	7.1
1984	178	119	297	8.4
1983	147	81	228	6.4
1982	140	93	233	6.4
COUNTY % OF TOTAL (1985) STATE DEATHS	3.4	3.0	3.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

SANDOVAL CO.	
MALE	741.2
FEMALE	600.7
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 7.1 deaths per 1000 population, accounting for 3.2% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for both females and males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 53% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 97% higher than expected for accidents. Observed deaths are higher than expected for diabetes, suicide, homicide and alcoholism.
- Both components of infant mortality are below state levels.
- Over 90% of all resident births occur in another county. Overall, the county experiences 3.1 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SANDOVAL CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	873	1039	84	100.0	100.0
HEART DISEASE	186	383	49	21.3	23.7
MALIGNANT NEOPLASMS	156	227	69	17.9	20.2
ALL ACCIDENT	120	61	197	13.7	10.0
MOTOR VEHICLE	81	31	261	9.3	6.1
ALL OTHER	39	30	130	4.5	3.9
STROKE	47	81	58	5.4	6.1
C.O.P.D.	24	30	80	2.7	4.7
DIABETES MELLITUS	23	17	135	2.6	2.7
SUICIDE	21	15	140	2.4	2.8
INFLUENZA/PNEUMONIA	17	25	68	1.9	2.7
CIRRHOSIS OF LIVER	17	17	100	1.9	1.9
HOMICIDE	25	14	179	2.9	1.9
ALCOHOLISM	23	2	1150	2.6	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	10	18	56	1.1	1.6
ATHEROSCLEROSIS	9	13	69	1.0	1.1
CONGENITAL ANOMALIES	12	10	120	1.4	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SANDOVAL	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.1	2.9
1 - 4	1.0	.8
5- 14	1.1	.8
15-24	3.2	3.7
25-34	5.1	4.4
35-44	5.0	4.0
45-54	7.2	6.4
55-64	12.2	13.4
65-74	22.6	22.1
75-84	23.4	24.5
85+	16.2	16.9

INFANT MORTALITY

SANDOVAL COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	5.8	2	2.9	2	2.9
1981	10	13.8	6	8.3	4	5.5
1982	8	10.8	7	9.4	1	1.3
1983	4	5.0	1	1.3	3	3.8
1984	12	13.3	5	5.6	7	7.8
1985	10	10.2	6	6.1	4	4.1
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SANDOVAL COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Bernalillo-Urban	44	20	2.2
Bernalillo-Rural	68	12	5.7
Cuba-All	53	27	2.0
Pena Blanca-All	8	3	2.7
Algodones-All	10	3	3.3
Jemez Springs-All	5	8	0.6
Other & Unspecified	793	239	3.3
COUNTY TOTAL	981	312	3.1
% OF COUNTY RESIDENT BIRTHS:			
Born in County	9.38%	XXXXX	XXXXX
Born in Other NM County	90.21%	XXXXX	XXXXX
Born Out of State	0.41%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

SAN JUAN COUNTY

SAN JUAN COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	52517	81433	91100	128000
% 19 AND YOUNGER	49.2%	41.7%	40.7%	40.2%
% 65 AND OLDER	4.9%	5.8%	6.5%	7.6%
% WHITE NON-HISPANIC	50.8%	54.6%	XXXXX	XXXXX
% WHITE HISPANIC	13.1%	11.4%	XXXXX	XXXXX
% BLACK	0.6%	0.4%	XXXXX	XXXXX
% INDIAN	35.1%	33.0%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	21.7%	16.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	5.2%	6.3%	6.3%	6.9%

- San Juan County ranks as the state's third most populous county and has a population density of 16.6 persons per square mile, accounting for 6.3% of the state's population.
- The county population is projected to reach 128,000 persons by the year 2000, a 57% increase over 1980, accounting for an increasing percentage of the state's population and maintaining a relatively high percentage of persons 19 and younger.
- The county birth rate increased to 24.5 births per 1000 population, still 30% higher than the state rate and accounting for 8.1% of New Mexico resident births.
- The percentage of births to single women is above the state level. Levels of low birth weight births and births to teens are below state levels.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SAN JUAN	1983	1984	1985
LIVE BIRTHS	2315	2240	2228
WHITE			
NON-HISPANIC	958	880	855
HISPANIC	388	340	353
BLACK	16	16	18
INDIAN	947	992	993
OTHER	6	12	9
COUNTY % OF TOTAL STATE BIRTHS	8.4%	8.2%	8.1%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	SAN JUAN NUMBER	SAN JUAN %	NEW MEXICO %
ALL AGES	2228	100.0%	100.0%
<15	6	0.3%	0.2%
15-17	101	4.5%	5.4%
18-19	219	9.8%	9.9%
(15-19)	320	14.4%	15.3%
20-24	760	34.1%	33.6%
25-29	636	28.5%	28.8%
30-34	364	16.3%	15.7%
35-39	123	5.5%	5.5%
=> 40	19	0.9%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
SAN JUAN										
1985	105.1	1.5	82.7	44.0	139.2	182.3	160.2	99.1	40.0	0.2
1984	97.6	0.9	67.1	34.0	119.9	177.4	145.2	83.3	40.4	5.2
1983	104.0	0.2	75.7	41.1	131.1	182.2	149.7	97.9	43.9	5.6
1982	118.6	0.7	85.1	49.9	141.5	207.5	166.3	85.7	37.4	5.8
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







SAN JUAN COUNTY

SAN JUAN COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	251	176	427	4.7
1984	278	187	465	4.9
1983	278	179	457	4.9
1982	320	174	494	5.5
COUNTY % OF TOTAL (1985) STATE DEATHS	4.6	4.2	4.4	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
SAN JUAN CO.	
MALE	866.4
FEMALE	613.1
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 4.7 deaths per 1000 population accounts for 4.4% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for females, and mortality for males higher than the state level.
- Accidents, heart disease and cancer are the leading causes of death, accounting for 53% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but over 2 1/2 times the expected level for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, influenza/pneumonia, homicide and alcoholism.
- Infant mortality is below state levels.
- Over 10% of resident births occur out of state. Overall, the county experiences 5.2 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SAN JUAN CO. — FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1779	1767	101	100.0	100.0
HEART DISEASE	286	595	48	16.1	23.7
MALIGNANT NEOPLASMS	282	365	77	15.9	20.2
ALL ACCIDENT	372	137	272	20.9	10.0
MOTOR VEHICLE	247	75	329	13.9	6.1
ALL OTHER	125	62	202	7.0	3.9
STROKE	83	126	66	4.7	6.1
C.O.P.D.	58	44	132	3.3	4.7
DIABETES MELLITUS	39	28	139	2.2	2.7
SUICIDE	61	34	179	3.4	2.8
INFLUENZA/PNEUMONIA	50	40	125	2.8	2.7
CIRRHOSIS OF LIVER	22	32	69	1.2	1.9
HOMICIDE	47	33	142	2.6	1.9
ALCOHOLISM	43	5	860	2.4	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	42	58	72	2.4	1.6
ATHEROSCLEROSIS	8	20	40	.4	1.1
CONGENITAL ANOMALIES	21	30	70	1.2	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SAN JUAN	NEW MEXICO
ALL	100.0	100.0
< 1	3.8	2.9
1 - 4	3.0	.8
5- 14	1.3	.8
15-24	6.6	3.7
25-34	8.3	4.4
35-44	6.1	4.0
45-54	7.6	6.4
55-64	13.3	13.4
65-74	19.0	22.1
75-84	19.5	24.5
85+	11.5	16.9

INFANT MORTALITY

SAN JUAN COUNTY	TOTAL INFANT Number	Rate	NEONATAL Number	Rate	POSTNEONATAL Number	Rate
1980	26	11.2	12	5.2	14	6.0
1981	25	10.3	11	4.5	14	5.8
1982	26	10.5	15	6.0	11	4.4
1983	17	7.3	6	2.6	11	4.8
1984	17	7.6	12	5.4	5	2.2
1985	19	8.5	11	4.9	8	3.6
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SAN JUAN COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Aztec-Urban	122	40	3.1
Aztec-Rural	82	16	5.1
Bloomfield-Urban	134	28	4.8
Bloomfield-Rural	85	17	5.0
Farmington-Urban	836	164	5.1
Farmington-Rural	76	25	3.0
Shiprock-All	296	50	5.9
Other & Unspecified	597	87	6.9
COUNTY TOTAL	2228	427	5.2
% OF COUNTY RESIDENT BIRTHS:			
Born in County	83.44%	XXXXX	XXXXX
Born in Other NM County	5.83%	XXXXX	XXXXX
Born Out of State	10.73%	XXXXX	XXXXX

****The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.*

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

SAN MIGUEL COUNTY

SAN MIGUEL COUNTY POPULATION

	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	21951	22751	24800	31300
% 19 AND YOUNGER	42.9%	36.7%	35.3%	35.5%
% 65 AND OLDER	9.9%	11.6%	11.4%	9.9%
% WHITE NON-HISPANIC	16.6%	17.3%	XXXXX	XXXXX
% WHITE HISPANIC	81.7%	81.3%	XXXXX	XXXXX
% BLACK	0.4%	0.4%	XXXXX	XXXXX
% INDIAN	0.4%	0.4%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	39.2%	26.9%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	2.2%	1.7%	1.7%	1.7%

- San Miguel County ranks as the state's 16th most populous county and has a population density of 5.2 persons per square mile, accounting for 1.7% of the total state population.
- The county population is projected to reach 31,300 persons by the year 2000, accounting for an unchanged percentage of the state's population. A large percentage of the county population is Hispanic.
- The county birth rate runs fairly close to the state level and for 1985 the rate of 18.8 births per 1000 population accounted for 1.7% of New Mexico resident births.
- Though showing decreases, the percentages of low birth weight births and births to single women remain above state levels, while the percent of births to teens dropped below state levels.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SAN MIGUEL	YEAR 1983	1984	1985
LIVE BIRTHS	482	495	467
WHITE			
NON-HISPANIC	42	48	46
HISPANIC	435	436	416
BLACK	3	6	1
INDIAN	1	4	3
OTHER	1	1	1
COUNTY % OF TOTAL STATE BIRTHS	1.8%	1.8%	1.7%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	SAN MIGUEL NUMBER	SAN MIGUEL %	NEW MEXICO %
ALL AGES	467	100.0%	100.0%
<15	2	0.4%	0.2%
15-17	23	4.9%	5.4%
18-19	37	7.9%	9.9%
(15-19)	60	12.8%	15.3%
20-24	162	34.7%	33.6%
25-29	143	30.6%	28.8%
30-34	70	15.0%	15.7%
35-39	26	5.6%	5.5%
=> 40	4	0.9%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
SAN MIGUEL:										
1985	75.9	2.1	51.4	33.3	77.7	111.0	133.6	80.0	29.7	--
1984	85.9	1.0	56.8	35.5	86.0	159.8	132.6	72.3	36.5	6.3
1983	85.0	2.0	67.1	39.9	104.5	152.9	121.1	80.4	31.2	1.8
1982	77.7	1.0	57.2	39.1	82.2	122.9	130.9	67.5	43.6	2.7
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







SAN MIGUEL COUNTY

SAN MIGUEL COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	111	76	187	7.5
1984	103	78	181	7.4
1983	105	68	173	7.2
1992	125	81	206	8.8
COUNTY % OF TOTAL (1985) STATE DEATHS	2.0	1.8	1.9	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

SAN MIGUEL CO.	
MALE	863.3
FEMALE	644.6
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 7 5 deaths per 1000 population accounts for 1 9% of all New Mexico resident deaths Adjusted rates show lower than state levels of mortality for females but higher for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for half of all deaths Observed deaths are lower than expected for heart disease and cancer, but 42% higher than expected for accidents Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), suicide, diabetes, influenza/pneumonia, cirrhosis, homicide and alcoholism
- The infant mortality rate is below the state rate.
- Nearly 13% of resident births occur in another county Overall, the county experiences 2 5 births for every death

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SAN MIGUEL CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	766	805	95	100.0	100.0
HEART DISEASE	182	310	59	23.8	23.7
MALIGNANT NEOPLASMS	137	164	84	17.9	20.2
ALL ACCIDENT	61	43	142	8.0	10.0
MOTOR VEHICLE	36	22	164	4.7	6.1
ALL OTHER	25	21	119	3.3	3.9
STROKE	42	71	59	5.5	6.1
C.O.P.D.	49	23	213	6.4	4.7
DIABETES MELLITUS	20	14	143	2.6	2.7
SUICIDE	19	11	173	2.5	2.8
INFLUENZA/PNEUMONIA	36	22	164	4.7	2.7
CIRRHOSIS OF LIVER	19	11	173	2.5	1.9
HOMICIDE	20	9	222	2.6	1.9
ALCOHOLISM	28	2	1400	3.7	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	6	11	55	.8	1.6
ATHEROSCLEROSIS	8	13	62	1.0	1.1
CONGENITAL ANOMALIES	2	6	33	.3	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SAN MIGUEL	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	1.7	2.9
1 - 4	.4	.8
5- 14	.6	.8
15-24	2.4	3.7
25-34	4.8	4.4
35-44	4.4	4.0
45-54	5.5	6.4
55-64	9.4	13.4
65-74	19.6	22.1
75-84	26.2	24.5
85+	25.0	16.9

INFANT MORTALITY

SAN MIGUEL COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	8.0	1	2.0	3	6.0
1981	4	8.5	3	6.4	1	2.1
1982	4	8.9	3	6.7	1	2.2
1983	3	6.2	2	4.1	1	2.1
1984	2	4.0	1	2.0	1	2.0
1985	4	8.6	3	6.4	1	2.1
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SAN MIGUEL COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Las Vegas-Urban	311	127	2.4
Las Vegas-Rural	52	16	3.3
Pecos-All	34	11	3.1
Other & Unspecified	70	33	2.1
COUNTY TOTAL	467	187	2.5
% OF COUNTY RESIDENT BIRTHS:			
Born in County	86.94%	XXXXX	XXXXX
Born in Other NM County	12.85%	XXXXX	XXXXX
Born Out of State	0.21%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

SANTA FE COUNTY

SANTA FE COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	53756	75360	84600	102600
% 19 AND YOUNGER	42.7%	33.6%	31.0%	29.0%
% 65 AND OLDER	7.8%	9.0%	10.0%	11.9%
% WHITE NON-HISPANIC	32.1%	40.8%	XXXXX	XXXXX
% WHITE HISPANIC	64.9%	55.0%	XXXXX	XXXXX
% BLACK	0.5%	0.5%	XXXXX	XXXXX
% INDIAN	2.0%	2.8%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	18.2%	11.2%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	5.3%	5.8%	5.8%	5.5%

- Santa Fe County ranks as the state's fourth most populous county and has a population density of 44 5 persons per square mile, accounting for 5 8% of the state's total population
- The county population is projected to exceed 102,000 persons by the year 2000, accounting for a decreasing percentage of the total state population and experiencing a drop in the percentage of population 19 and younger
- The county birth rate of 17 1 births per 1000 population remains 10% below the state rate, accounting for 5 3% of all New Mexico resident births.
- The percentage of births to teen mothers increased but remained below state levels, while the percentage of births to single women increased above state levels and the percent low birth weight exceeds state levels by 9%



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SANTA FE	1983	1984	1985
LIVE BIRTHS	1366	1420	1449
WHITE NON-HISPANIC	398	442	439
HISPANIC	900	895	914
BLACK	5	15	12
INDIAN	61	63	69
OTHER	2	5	15
COUNTY % OF TOTAL STATE BIRTHS	5.0%	5.2%	5.3%



BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	SANTA FE NUMBER	SANTA FE %	NEW MEXICO %
ALL AGES	1449	100.0%	100.0%
<15	2	0.1%	0.2%
15-17	76	5.2%	5.4%
18-19	113	7.8%	9.9%
(15-19)	189	13.0%	15.3%
20-24	411	28.4%	33.6%
25-29	395	27.3%	28.8%
30-34	296	20.4%	15.7%
35-39	133	9.2%	5.5%
=> 40	23	1.6%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
SANTA FE:										
1985	69.5	0.6	59.3	41.0	84.6	114.5	116.6	74.3	34.2	--
1984	70.2	0.3	42.3	20.0	78.6	125.5	98.5	80.0	38.6	5.0
1983	68.5	--	50.4	32.3	79.7	116.4	92.5	81.2	36.7	3.1
1982	62.5	0.3	48.9	28.2	82.3	106.4	84.2	71.1	35.4	1.7
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







SANTA FE COUNTY

SANTA FE COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	257	242	499	5.9
1984	276	224	500	6.1
1983	253	251	504	6.3
1982	252	238	490	6.3
COUNTY % OF TOTAL (1985) STATE DEATHS	4.7	5.8	5.2	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

SANTA FE CO.	
MALE	694.7
FEMALE	644.2
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 5.9 deaths per 1000 population, accounting for 5.2% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for males and females.

- Cancer, heart disease and accidents are the leading causes of death, accounting for half of all deaths. Observed deaths are lower than expected for cancer and heart disease, but 34% higher than expected for accidents. Observed deaths are higher than expected for chronic obstructive pulmonary disease (COPD), diabetes, suicide, cirrhosis, homicide and alcoholism.

- Infant mortality is lower than statewide.

- 11% of resident births occur in another county. Overall, the county experiences 2.9 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SANTA FE CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1898	2175	87	100.0	100.0
HEART DISEASE	377	804	47	19.9	23.7
MALIGNANT NEOPLASMS	403	457	88	21.2	20.2
ALL ACCIDENT	179	134	134	9.4	10.0
MOTOR VEHICLE	113	70	161	6.0	6.1
ALL OTHER	66	64	103	3.5	3.9
STROKE	118	179	66	6.2	6.1
C.O.P.D.	84	58	145	4.4	4.7
DIABETES MELLITUS	52	37	141	2.7	2.7
SUICIDE	59	35	169	3.1	2.8
INFLUENZA/PNEUMONIA	54	54	100	2.8	2.7
CIRRHOSIS OF LIVER	58	36	161	3.1	1.9
HOMICIDE	42	32	131	2.2	1.9
ALCOHOLISM	41	5	820	2.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	19	33	58	1.0	1.6
ATHEROSCLEROSIS	18	30	60	.9	1.1
CONGENITAL ANOMALIES	8	19	42	.4	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SANTA FE	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	1.6	2.9
1 - 4	.5	.8
5- 14	.6	.8
15-24	5.0	3.7
25-34	4.2	4.4
35-44	4.6	4.0
45-54	7.2	6.4
55-64	12.5	13.4
65-74	20.9	22.1
75-84	24.8	24.5
85+	18.1	16.9

INFANT MORTALITY

SANTA FE COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	14	11.0	9	7.1	5	3.9
1981	5	3.9	3	2.3	2	1.5
1982	4	3.3	3	2.5	1	0.8
1983	5	3.7	1	0.7	4	2.9
1984	13	9.2	9	6.3	4	2.8
1985	6	4.1	4	2.8	2	1.4
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SANTA FE COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Santa Fe-Urban	906	387	2.3
Santa Fe-Rural	315	54	5.8
Tesuque-All	21	5	4.2
Pojoaque-All	24	4	6.0
Other & Unspecified	183	49	3.7
COUNTY TOTAL	1449	499	2.9
% OF COUNTY RESIDENT BIRTHS:			
Born in County	88.54%	XXXXX	XXXXX
Born in Other NM County	11.11%	XXXXX	XXXXX
Born Out of State	0.35%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

SIERRA COUNTY

SIERRA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	7189	8454	9500	10500
% 19 AND YOUNGER	30.4%	24.2%	19.4%	16.9%
% 65 AND OLDER	24.7%	29.5%	35.1%	43.4%
% WHITE NON-HISPANIC	63.3%	73.9%	XXXXX	XXXXX
% WHITE HISPANIC	35.8%	25.2%	XXXXX	XXXXX
% BLACK	0.1%	0.1%	XXXXX	XXXXX
% INDIAN	0.2%	0.6%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	23.9%	18.9%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.7%	0.6%	0.7%	0.6%

- Sierra County ranks as the state's 25th most populous county and has a population density of 2.3 persons per square mile, accounting for 0.7% of the state's total population
- The county population is projected to reach 10,500 persons by the year 2000, with over 0% of the population expected to be 65 and older
- The county birth rate (10.3 births per 1000 population) is 46% lower than the state rate because of the large component of older population. The fertility rate is only 9% lower than that for the state
- The percentage of births to teens is slightly above state levels; the percent of births to single women dropped below the state percent, and the percent of low birth weight births decreased markedly below state levels


BIRTH RATES PER 1000 POPULATION

NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SIERRA	YEAR 1983	1984	1985
LIVE BIRTHS	128	95	98
WHITE			
NON-HISPANIC	70	51	49
HISPANIC	58	43	47
BLACK	0	0	1
INDIAN	0	1	1
OTHER	0	0	0
COUNTY % OF TOTAL STATE BIRTHS	.5%	.3%	.4%


PERCENT LOW BIRTH WEIGHT

BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	SIERRA NUMBER	SIERRA %	NEW MEXICO %
ALL AGES	98	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	2	2.0%	5.4%
18-19	14	14.3%	9.9%
(15-19)	16	16.3%	15.3%
20-24	34	34.7%	33.6%
25-29	29	29.6%	28.8%
30-34	16	16.3%	15.7%
35-39	3	3.1%	5.5%
=> 40	0	0.0%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
SIERRA:										
1985	73.0	--	50.5	11.2	101.4	149.1	156.8	97.6	14.6	--
1984	73.8	--	75.5	52.6	125.0	184.3	102.7	29.3	29.3	7.4
1983	99.8	--	115.5	63.2	229.9	194.4	190.2	63.7	19.6	4.9
1982	74.9	--	86.6	57.9	149.4	148.1	125.0	49.0	14.7	9.9
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44


BIRTHS TO TEEN MOTHERS


BIRTHS TO SINGLE MOTHERS


LEVEL OF PRENATAL CARE 1985

SIERRA COUNTY

SIERRA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	90	69	159	16.7
1984	94	70	164	18.3
1983	92	55	147	16.5
1982	87	56	143	16.1
COUNTY % OF TOTAL (1985) STATE DEATHS	1.6	1.7	1.6	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
SIERRA CO.	
MALE	830.4
FEMALE	768.5
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 16.7 deaths per 1000 population, accounting for 1.6% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for both females and males.
- Heart disease, cancer and chronic obstructive pulmonary disease (COPD) are the leading causes of death, accounting for 63% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but almost three times higher than expected for COPD. Observed deaths are higher than expected for accidents, suicide and cirrhosis.
- The infant mortality rate is well above the state rate.
- Almost 89% of all resident births occur in another county. Overall, the county experiences less than one birth (.6) for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SIERRA CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	598	605	99	100.0	100.0
HEART DISEASE	213	252	85	35.6	23.7
MALIGNANT NEOPLASMS	102	130	78	17.1	20.2
ALL ACCIDENT	38	19	200	6.4	10.0
MOTOR VEHICLE	19	8	238	3.2	6.1
ALL OTHER	19	11	173	3.2	3.9
STROKE	34	57	60	5.7	6.1
C.O.P.D.	59	20	295	9.9	4.7
DIABETES MELLITUS	12	11	109	2.0	2.7
SUICIDE	15	5	300	2.5	2.8
INFLUENZA/PNEUMONIA	15	17	88	2.5	2.7
CIRRHOSIS OF LIVER	16	7	229	2.7	1.9
HOMICIDE	2	3	67	.3	1.9
ALCOHOLISM	1	1	100	.2	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	2	2	100	.3	1.6
ATHEROSCLEROSIS	4	10	40	.7	1.1
CONGENITAL ANOMALIES	--	2	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SIERRA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.1	2.9
1 - 4	--	.8
5- 14	.2	.8
15-24	.8	3.7
25-34	.8	4.4
35-44	1.5	4.0
45-54	4.9	6.4
55-64	11.5	13.4
65-74	29.3	22.1
75-84	28.7	24.5
85+	20.2	16.9

INFANT MORTALITY

SIERRA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	1	11.0	1	11.0	0	0.0
1981	3	28.8	2	19.2	1	9.6
1982	1	10.4	0	0.0	1	10.4
1983	3	23.4	3	23.4	0	0.0
1984	5	52.6	4	42.1	1	10.5
1985	2	20.4	1	10.2	1	10.2
NEW MEXICO 1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SIERRA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Truth or Consequences-Urban	19	112	0.2
Truth or Consequences-Rural	50	6	8.3
Other & Unspecified	29	41	0.7
COUNTY TOTAL	98	159	0.6
% OF COUNTY RESIDENT BIRTHS:			
Born in County	8.16%	XXXXX	XXXXX
Born in Other NM County	88.78%	XXXXX	XXXXX
Born Out of State	3.06%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

SOCORRO COUNTY

SOCORRO COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	9763	12566	13600	18100
% 19 AND YOUNGER	45.1%	38.6%	36.0%	33.1%
% 65 AND OLDER	7.7%	8.8%	9.4%	11.0%
% WHITE NON-HISPANIC	31.0%	41.9%	XXXXX	XXXXX
% WHITE HISPANIC	60.0%	46.4%	XXXXX	XXXXX
% BLACK	0.7%	0.4%	XXXXX	XXXXX
% INDIAN	7.2%	9.9%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	29.5%	22.2%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.0%	1.0%	0.9%	1.0%

- Socorro County ranks as the state's 23rd most populous county and has a population density of 2.1 persons per square mile, accounting for 0.9% of the state's total population.
- The county population is projected to reach 18,100 persons by the year 2000; 1% of the total state population.
- The county birth rate of 18.2 births per 1000 population is 4% below the state rate and accounts for 0.9% of New Mexico resident live births.
- The percentage of births to teen women rose above state levels. The percentage of births to single women remained below state levels, while the percentage of low birth weight births decreased to former levels below the state average.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

SOCORRO	YEAR 1983	1984	1985
LIVE BIRTHS	277	240	248
WHITE			
NON-HISPANIC	76	64	66
HISPANIC	146	133	132
BLACK	2	0	1
INDIAN	51	40	46
OTHER	2	3	3
COUNTY % OF TOTAL STATE BIRTHS	1.0%	.9%	.9%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	SOCORRO NUMBER	SOCORRO %	NEW MEXICO %
ALL AGES	248	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	16	6.5%	5.4%
18-19	30	12.1%	9.9%
(15-19)	46	18.5%	15.3%
20-24	79	31.9%	33.6%
25-29	65	26.2%	28.8%
30-34	45	18.1%	15.7%
35-39	12	4.8%	5.5%
=> 40	1	0.4%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
SOCORRO										
1985	84.5	--	78.4	45.2	128.8	134.6	110.7	92.0	30.7	--
1984	74.7	1.5	43.1	36.9	51.4	123.6	122.4	82.8	25.4	3.1
1983	87.1	1.5	66.0	29.7	113.6	129.1	136.0	100.9	35.9	1.6
1982	72.9	--	56.2	56.9	55.2	107.4	123.7	68.7	30.8	3.1
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







SOCORRO COUNTY

SOCORRO COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	36	27	63	4.6
1984	55	38	93	6.4
1983	46	37	83	5.8
1982	52	45	97	7.0
COUNTY % OF TOTAL (1985) STATE DEATHS	0.7	0.6	0.7	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

SOCORRO CO.	
MALE	754.9
FEMALE	685.6
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 4.6 deaths per 1000 population accounting for 0.7% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for females, and for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 57% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but 65% higher than expected for accidents. Observed deaths are higher than expected for diabetes, suicide and alcoholism.
- There were no infant deaths reported for the county for 1985.
- 30% of resident births occur in another county. Overall, the county experiences 3.9 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

SOCORRO CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	363	353	103	100.0	100.0
HEART DISEASE	101	129	78	27.8	23.7
MALIGNANT NEOPLASMS	67	75	89	18.5	20.2
ALL ACCIDENT	38	23	165	10.5	10.0
MOTOR VEHICLE	28	12	233	7.7	6.1
ALL OTHER	10	11	91	2.8	3.9
STROKE	16	28	57	4.4	6.1
C.O.P.D.	11	10	110	3.0	4.7
DIABETES MELLITUS	14	6	233	3.9	2.7
SUICIDE	9	6	150	2.5	2.8
INFLUENZA/PNEUMONIA	5	9	56	1.4	2.7
CIRRHOSIS OF LIVER	6	6	100	1.7	1.9
HOMICIDE	6	5	120	1.7	1.9
ALCOHOLISM	4	.83	482	1.1	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	8	6	133	2.2	1.6
ATHEROSCLEROSIS	2	5	40	.6	1.1
CONGENITAL ANOMALIES	2	3	67	.6	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS SOCORRO	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.5	2.9
1 - 4	.4	.8
5- 14	1.7	.8
15-24	3.3	3.7
25-34	3.3	4.4
35-44	4.1	4.0
45-54	2.9	6.4
55-64	11.6	13.4
65-74	22.3	22.1
75-84	26.4	24.5
85+	21.5	16.9

INFANT MORTALITY

SOCORRO COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	4	16.2	3	12.1	1	4.0
1981	2	7.3	1	3.6	1	3.6
1982	4	17.3	1	4.3	3	13.0
1983	1	3.6	0	0.0	1	3.6
1984	5	20.8	2	8.3	3	12.5
1985	0	0.0	0	0.0	0	0.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

SOCORRO COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Magdalena-Urban	11	5	2.2
Magdalena-Rural	41	3	13.7
Socorro-Urban	144	36	4.0
Socorro-Rural	26	5	5.2
Other & Unspecified	26	14	1.9
COUNTY TOTAL	248	63	3.9
% OF COUNTY RESIDENT BIRTHS:			
Born in County	70.16%	XXXXX	XXXXX
Born in Other NM County	29.84%	XXXXX	XXXXX
Born Out of State	----	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

TAOS COUNTY

TAOS COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	17516	19456	21900	28000
% 19 AND YOUNGER	45.8%	37.6%	35.8%	33.2%
% 65 AND OLDER	9.3%	9.7%	9.7%	9.6%
% WHITE NON-HISPANIC	5.1%	23.9%	XXXXX	XXXXX
% WHITE HISPANIC	86.3%	68.8%	XXXXX	XXXXX
% BLACK	0.2%	0.2%	XXXXX	XXXXX
% INDIAN	6.8%	6.4%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	35.8%	24.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	1.7%	1.5%	1.5%	1.5%

- Taos County ranks as the states 17th most populous county and has a population density of 9.7 persons per square mile, accounting for 1.5% of the state population.
- The county population is projected to reach 28,000 persons by the year 2000 with a third of the population expected to be 19 and younger.
- A decreasing trend in the birth rate leaves the county birth rate (20.9 births per 1000 population) 11% above the state rate accounting for 1.7% of New Mexico resident births.
- A slight decrease in the percentage of births to single women dropped county figures to state levels, while fluctuating percentages leaves the county percentage of births to teens below state levels and the percentage of low birth weight births above the state level.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

TAOS	YEAR 1983	1984	1985
LIVE BIRTHS	450	443	458
WHITE			
NON-HISPANIC	121	101	88
HISPANIC	295	305	320
BLACK	2	1	2
INDIAN	29	34	45
OTHER	3	2	3
COUNTY % OF TOTAL STATE BIRTHS	1.6%	1.6%	1.7%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	TAOS NUMBER	TAOS %	NEW MEXICO %
ALL AGES	458	100.0%	100.0%
<15	1	0.2%	0.2%
15-17	22	4.8%	5.4%
18-19	39	8.5%	9.9%
(15-19)	61	13.3%	15.3%
20-24	135	29.5%	33.6%
25-29	122	26.6%	28.8%
30-34	102	22.3%	15.7%
35-39	29	6.3%	5.5%
=> 40	8	1.7%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
TAOS:										
1985	90.5	1.1	70.0	43.1	108.0	174.2	157.4	95.7	29.9	0.9
1984	91.3	--	59.2	31.7	113.4	196.1	127.3	82.9	50.9	5.8
1983	92.3	3.0	72.5	34.8	147.1	160.1	134.4	101.6	46.8	4.9
1982	98.6	--	67.1	35.7	129.2	203.3	143.2	95.2	46.4	5.0
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







TAOS COUNTY

TAOS COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	94	57	151	6.9
1984	83	77	160	7.7
1983	95	52	147	7.1
1982	89	51	140	6.9
COUNTY % OF TOTAL (1985) STATE DEATHS	1.7	1.4	1.6	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

TAOS CO.	
MALE	857.9
FEMALE	714.1
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate decreased to 6.9 deaths per 1000 population, accounting for 1.6% of all New Mexico resident deaths. Adjusted rates show a higher than state level of mortality for males, as well as for females.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 54% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but twice the expected level for accidents. Observed deaths are higher than expected for diabetes, suicide, influenza/pneumonia, cirrhosis, homicide and alcoholism.
- The neonatal component of infant mortality is higher than the state level.
- Over 26% of resident births occur in another county. Overall, the county experiences 3 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

TAOS CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	541	599	90	100.0	100.0
HEART DISEASE	118	224	53	21.8	23.7
MALIGNANT NEOPLASMS	103	125	82	19.0	20.2
ALL ACCIDENT	73	35	209	13.5	10.0
MOTOR VEHICLE	46	18	256	8.5	6.1
ALL OTHER	27	17	159	5.0	3.9
STROKE	38	50	76	7.0	6.1
C.O.P.D.	10	17	59	1.8	4.7
DIABETES MELLITUS	15	10	150	2.8	2.7
SUICIDE	16	9	178	3.0	2.8
INFLUENZA/PNEUMONIA	22	15	147	4.1	2.7
CIRRHOSIS OF LIVER	11	9	122	2.0	1.9
HOMICIDE	14	8	175	2.6	1.9
ALCOHOLISM	20	1	2000	3.7	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	7	10	70	1.3	1.6
ATHEROSCLEROSIS	8	9	89	1.5	1.1
CONGENITAL ANOMALIES	1	5	20	.2	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS TAOS	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.0	2.9
1 - 4	.9	.8
5- 14	1.5	.8
15-24	4.3	3.7
25-34	7.6	4.4
35-44	6.1	4.0
45-54	5.4	6.4
55-64	8.7	13.4
65-74	17.3	22.1
75-84	23.2	24.5
85+	22.1	16.9

INFANT MORTALITY

TAOS COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	3	7.2	2	4.8	1	2.4
1981	1	2.4	1	2.4	0	0.0
1982	2	4.3	1	2.2	1	2.2
1983	5	11.1	3	6.7	2	4.4
1984	4	9.0	3	6.8	1	2.3
1985	5	10.9	4	8.7	1	2.2
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

TAOS COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Questa-All	63	9	7.0
Rancho de Taos-All	59	17	3.5
Taos-Urban	120	52	2.3
Taos-Rural	61	14	4.4
Penasco-All	16	4	4.0
Other & Unspecified	139	55	2.5
COUNTY TOTAL	458	151	3.0
% OF COUNTY RESIDENT BIRTHS:			
Born in County	69.21%	XXXXX	XXXXX
Born in Other NM County	26.20%	XXXXX	XXXXX
Born Out of State	4.59%	XXXXX	XXXXX

***The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

TORRANCE COUNTY

TORRANCE COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	5290	7491	8600	11400
% 19 AND YOUNGER	41.1%	35.9%	35.1%	33.6%
% 65 AND OLDER	11.9%	12.0%	13.6%	14.4%
% WHITE NON-HISPANIC	46.8%	57.3%	XXXXX	XXXXX
% WHITE HISPANIC	52.6%	41.0%	XXXXX	XXXXX
% BLACK	---	0.2%	XXXXX	XXXXX
% INDIAN	0.1%	0.5%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	32.6%	19.7%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.5%	0.6%	0.6%	0.6%

- .Torrance County ranks as the state's 26th most populous county and has a population density of 2.6 persons per square mile, accounting for 0.6% of the total state population.
- .The county population is projected to reach over 11,000 persons by the year 2000.
- .The county birth rate remains below state levels with a birth rate of 15.8 births per 1000 population in 1985; a rate 16% below the state rate and accounting for one-half of one percent of New Mexico resident births.
- .The county percentage of births to single women has decreased to almost half the state percentage. The percentage of births to teen women is slightly below state levels, while the percentage low birth weight births in the county fluctuated notably over the last three years.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

TORRANCE	1983	1984	1985
	YEAR		
LIVE BIRTHS	146	136	136
WHITE			
NON-HISPANIC	62	60	58
HISPANIC	78	71	73
BLACK	2	1	1
INDIAN	4	3	4
OTHER	0	1	0
COUNTY % OF TOTAL STATE BIRTHS	.5%	.5%	.5%



BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	TORRANCE NUMBER	TORRANCE %	NEW MEXICO %
ALL AGES	136	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	8	5.9%	5.4%
18-19	13	9.6%	9.9%
(15-19)	21	15.4%	15.3%
20-24	45	33.1%	33.6%
25-29	44	32.4%	28.8%
30-34	17	12.5%	15.7%
35-39	7	5.1%	5.5%
=> 40	2	1.5%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
TORRANCE:										
1985	77.1	--	56.3	35.9	86.7	171.7	164.2	51.1	23.4	--
1984	81.4	--	59.9	19.0	130.1	182.8	107.1	81.2	31.1	--
1983	88.6	--	76.0	48.1	124.0	174.5	125.0	78.9	49.5	--
1982	78.5	--	65.6	29.7	127.1	126.9	141.4	57.4	55.6	--
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







TORRANCE COUNTY

TORRANCE COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	44	24	68	7.9
1984	45	20	65	7.9
1983	38	23	61	7.5
1982	36	21	57	7.2
COUNTY % OF TOTAL (1985) STATE DEATHS	0.8	0.6	0.7	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
TORRANCE CO.	
MALE	878.8
FEMALE	584.6
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 7.9 deaths per 1000 population, accounted for 0.7% of all New Mexico resident deaths. Adjusted rates show lower than state levels of mortality for females but higher for males.
- Heart disease, cancer and accidents are the leading causes of death, accounting for 52% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but over twice the expected level for accidents. Observed deaths are higher than expected for stroke, chronic obstructive pulmonary disease (COPD) and suicide.
- Due to the relatively small number of births, infant mortality fluctuates notably; and was twice the state rate in 1985.
- Over 98% of resident births occur in another county. Overall, the county experiences 2 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

TORRANCE CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	215	269	80	100.0	100.0
HEART DISEASE	41	103	40	19.1	23.7
MALIGNANT NEOPLASMS	39	57	68	18.1	20.2
ALL ACCIDENT	32	14	229	14.9	10.0
MOTOR VEHICLE	23	7	329	10.7	6.1
ALL OTHER	9	7	129	4.2	3.9
STROKE	41	23	178	19.1	6.1
C.O.P.D.	13	8	163	6.0	4.7
DIABETES MELLITUS	6	5	120	2.8	2.7
SUICIDE	9	4	225	4.2	2.8
INFLUENZA/PNEUMONIA	6	7	86	2.8	2.7
CIRRHOSIS OF LIVER	1	4	25	.5	1.9
HOMICIDE	2	3	67	.9	1.9
ALCOHOLISM	2	1	200	.9	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	3	4	75	1.4	1.6
ATHEROSCLEROSIS	3	4	75	1.4	1.1
CONGENITAL ANOMALIES	2	2	100	.9	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS TORRANCE	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	2.6	2.9
1 - 4	.5	.8
5- 14	2.1	.8
15-24	2.6	3.7
25-34	5.2	4.4
35-44	4.1	4.0
45-54	5.7	6.4
55-64	14.4	13.4
65-74	28.9	22.1
75-84	21.1	24.5
85+	12.9	16.9

INFANT MORTALITY

TORRANCE COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	1	7.7	1	7.7	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	2	15.9	1	7.9	1	7.9
1983	2	13.7	1	6.8	1	6.8
1984	0	0.0	0	0.0	0	0.0
1985	3	22.1	3	22.1	0	0.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

TORRANCE COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Estancia-All	23	14	1.6
Moriarty-All	60	18	3.3
Mountainair-Urban	7	17	0.4
Mountainair-Rural	16	2	8.0
Encino-All	4	1	4.0
Other & Unspecified	26	16	1.6
COUNTY TOTAL	136	68	2.0
% OF COUNTY RESIDENT BIRTHS:			
Born in County	1.47%	XXXXX	XXXXX
Born in Other NM County	98.53%	XXXXX	XXXXX
Born Out of State	----	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

UNION COUNTY

UNION COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	4925	4725	5100	4900
% 19 AND YOUNGER	37.6%	33.0%	32.1%	30.4%
% 65 AND OLDER	15.4%	17.2%	15.8%	16.8%
% WHITE NON-HISPANIC	70.5%	68.3%	XXXXX	XXXXX
% WHITE HISPANIC	28.3%	31.2%	XXXXX	XXXXX
% BLACK	0.1%	---	XXXXX	XXXXX
% INDIAN	0.2%	0.3%	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	26.2%	19.1%	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.5%	0.4%	0.4%	0.3%

- Union County ranks as the state's 28th most populous county and has a population density of 1.3 persons per square mile, accounting for 0.4% of the total state population.
- The county population is projected to decrease to 4,900 persons by the year 2000 with decreasing percentages in the 19 and younger component of population.
- The county birth rate remains below the state rate. As of 1985, the county rate of 14.9% births per 1000 population was 21% below the state rate, accounting for 0.3% of New Mexico resident births.
- A notable decrease in the percent of births to single women has dropped county levels below state levels. The percentage of births to teens is presently 36% above the state level and the percentage of low birth weight births shot up to 91% above the state percent.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

UNION	1983	1984	1985
LIVE BIRTHS	80	84	76
WHITE			
NON-HISPANIC	41	51	32
HISPANIC	39	33	44
BLACK	0	0	0
INDIAN	0	0	0
OTHER	3	0	0
COUNTY % OF TOTAL STATE BIRTHS	.3%	.3%	.3%



BIRTH DISTRIBUTION BY AGE OF MOTHER

AGE OF MOTHER	UNION NUMBER	UNION %	NEW MEXICO %
ALL AGES	76	100.0%	100.0%
<15	0	0.0%	0.2%
15-17	6	7.9%	5.4%
18-19	10	13.2%	9.9%
(15-19)	16	21.1%	15.3%
20-24	20	26.3%	33.6%
25-29	25	32.9%	28.8%
30-34	9	11.8%	15.7%
35-39	2	2.6%	5.5%
=> 40	4	5.3%	0.9%

NEW MEXICO RESIDENTS FERTILITY RATES BIRTH RATES BY AGE OF MOTHER BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-19 15-17	15-19 18-19	20-24	25-29	30-34	35-39	40-49
UNION:										
1985	70.2	--	85.1	54.1	129.9	74.6	140.4	62.9	13.0	3.9
1984	80.1	--	66.1	37.8	138.9	152.4	162.7	72.7	18.9	--
1983	77.2	--	47.2	27.3	98.6	160.5	176.8	55.2	19.1	3.8
1982	86.7	4.6	76.3	27.8	202.9	164.6	142.9	106.3	19.5	--
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







UNION COUNTY

UNION COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	32	27	59	11.6
1984	24	27	51	9.5
1983	34	20	54	10.2
1982	23	19	42	8.1
COUNTY % OF TOTAL (1985) STATE DEATHS	0.6	0.6	0.6	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE	
UNION CO.	
MALE	915.8
FEMALE	743.4
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate rose to 11 6 deaths per 1000 population accounting for 0 6% of all New Mexico resident deaths Adjusted rates show higher than state levels of mortality for both females and males

- Heart disease, cancer and chronic obstructive pulmonary disease (COPD) are the leading causes of death, accounting for 54% of all deaths Observed deaths are lower than expected for heart disease and cancer, but 2 1/2 times the expected level for COPD

- Infant mortality levels are above statewide levels, but are subject to fluctuation due to relatively few births

- 64% of resident births occur within the county. Overall, the county experiences 1 3 births for every death

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

UNION CO. FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	240	231	104	100.0	100.0
HEART DISEASE	66	93	71	27.5	23.7
MALIGNANT NEOPLASMS	45	47	96	18.8	20.2
ALL ACCIDENT	13	9	144	5.4	10.0
MOTOR VEHICLE	5	4	125	2.1	6.1
ALL OTHER	8	5	160	3.3	3.9
STROKE	13	22	59	5.4	6.1
C.O.P.D.	18	7	257	7.5	4.7
DIABETES MELLITUS	4	4	100	1.7	2.7
SUICIDE	4	2	200	1.7	2.8
INFLUENZA/PNEUMONIA	16	6	267	6.7	2.7
CIRRHOSIS OF LIVER	2	3	67	.8	1.9
HOMICIDE	--	2	--	--	1.9
ALCOHOLISM	3	.34	882	1.3	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	2	2	100	.8	1.6
ATHEROSCLEROSIS	1	4	25	.4	1.1
CONGENITAL ANOMALIES	--	1	--	--	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS UNION	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	1.2	2.9
1 - 4	--	.8
5- 14	.6	.8
15-24	4.3	3.7
25-34	.6	4.4
35-44	1.2	4.0
45-54	1.8	6.4
55-64	14.6	13.4
65-74	19.5	22.1
75-84	28.0	24.5
85+	28.0	16.9

INFANT MORTALITY

UNION COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	0	0.0	0	0.0	0	0.0
1981	0	0.0	0	0.0	0	0.0
1982	1	11.2	1	11.2	0	0.0
1983	0	0.0	0	0.0	0	0.0
1984	0	0.0	0	0.0	0	0.0
1985	2	26.3	2	26.3	0	0.0
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

UNION COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Clayton-Urban	57	41	1.4
Clayton-Rural	4	2	2.0
Des Moines-All	0	3	0.0
Other & Unspecified	15	13	1.2
COUNTY TOTAL	76	59	1.3
% OF COUNTY RESIDENT BIRTHS:			
Born in County	64.47%	XXXXX	XXXXX
Born in Other NM County	13.16%	XXXXX	XXXXX
Born Out of State	22.37%	XXXXX	XXXXX

***The SMR is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.

INFANT = <1 year, NEONATAL =<28 days, POSTNEONATAL = 28 days to <1 year.

VALENCIA COUNTY

VALENCIA COUNTY POPULATION

YEAR	1970	1980	ESTIMATED 1985	PROJECTED 2000
NUMBER	#####	30768	35900	48600
% 19 AND YOUNGER	###	38.7%	35.3%	34.2%
% 65 AND OLDER	###	10.4%	10.6%	11.5%
% WHITE NON-HISPANIC	###	###	XXXXX	XXXXX
% WHITE HISPANIC	###	###	XXXXX	XXXXX
% BLACK	###	###	XXXXX	XXXXX
% INDIAN	###	###	XXXXX	XXXXX
% FAMILIES BELOW POVERTY	###	###	XXXXX	XXXXX
COUNTY % OF STATE POPULATION	0.0%	2.4%	2.5%	2.6%

- Valencia County ranks as the states 12th most populous county and has a population density of 24.4 persons per square mile, accounting for 2.5% of the total state population.
- The county population is projected to reach 48,600 persons by the year 2000.
- The county birth rate increased to 17.5 births per 1000 population - a rate 7% below the state level and accounting for 2.3% of New Mexico resident births.
- The percentage of births to single women remains below state levels while the percentage of low birth weight births exceeds state levels; and births to teens have dropped below state levels.

***DATA AVAILABLE ONLY IN A FORM INCLUDING BOTH VALENCIA AND CIBOLA.



NUMBER OF LIVE BIRTHS BY RACE/ETHNIC

VALENCIA	YEAR 1983	1984	1985
LIVE BIRTHS	595	630	629
WHITE			
NON-HISPANIC	199	217	213
HISPANIC	347	379	379
BLACK	3	7	2
INDIAN	43	26	31
OTHER	3	1	4
COUNTY % OF TOTAL STATE BIRTHS	2.2%	2.3%	2.3%



BIRTH DISTRIBUTION
BY AGE OF MOTHER

AGE OF MOTHER	VALENCIA NUMBER	VALENCIA %	NEW MEXICO %
ALL AGES	629	100.0%	100.0%
<15	1	0.2%	0.2%
15-17	35	5.6%	5.4%
18-19	46	7.3%	9.9%
(15-19)	81	12.9%	15.3%
20-24	211	33.5%	33.6%
25-29	194	30.8%	28.8%
30-34	94	14.9%	15.7%
35-39	41	6.5%	5.5%
=> 40	7	1.1%	0.9%

NEW MEXICO RESIDENTS
FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER
BY COUNTY 1982, 1983, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP 10-14	15-19 TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
VALENCIA:										
1985	77.9	0.6	52.4	37.6	74.6	136.3	156.7	65.1	33.1	0.5
1984	77.0	--	66.2	37.0	119.7	154.1	132.0	59.7	20.6	4.1
1983	79.4	1.2	70.1	38.7	127.6	154.6	130.7	62.4	30.7	2.6
1982	90.2	2.0	71.8	43.2	124.3	189.4	144.3	83.6	22.6	3.1
NEW MEXICO:										
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	0.1

(a) Births Per 1,000 Women Ages 15-44







VALENCIA COUNTY

VALENCIA COUNTY DEATHS

YEAR	MALE	FEMALE	TOTAL NUMBER	TOTAL CRUDE DEATH RATE
1985	162	104	266	7.4
1984	145	116	261	7.0
1983	150	81	231	6.8
1982	158	105	263	7.9
COUNTY % OF TOTAL (1985) STATE DEATHS	3.0	2.5	2.8	

Rate Per 1000 Population

AGE ADJUSTED DEATHS RATES

1983 - 85 AVERAGE

VALENCIA CO.	
MALE	866.2
FEMALE	684.4
NEW MEXICO	
MALE	828.5
FEMALE	693.7

RATE PER 100,000 POPULATION

- The county crude death rate of 7.4 deaths per 1000 population accounts for 2.8% of all New Mexico resident deaths. Adjusted rates show higher than state levels of mortality for males, but lower than state levels for females.
- Mortality by cause is based on the combined Cibola/Valencia county data and shows heart disease, cancer and accidents to be the leading causes of death, accounting for 52% of all deaths. Observed deaths are lower than expected for heart disease and cancer, but higher than expected for accidents, chronic obstructive pulmonary disease (COPD), diabetes, suicide, influenza/pneumonia and alcoholism.
- Neonatal mortality exceeds state levels.
- 78% of resident births occur in another county. Overall, the county experiences 2.4 births for every death.

STANDARDIZED MORTALITY RATIOS - AGE/SEX ADJUSTED

VALENCIA CO. (CIBOLA INCLUDED) FOUR YEAR AGGREGATE 1979 - 1982

CAUSE	OBSERVED DEATHS	EXPECTED DEATHS	SMR	% OF TOTAL DEATHS	NM % OF TOTAL DEATHS
ALL CAUSES	1506	1545	97	100.0	100.0
HEART DISEASE	316	547	58	21.0	23.7
MALIGNANT NEOPLASMS	276	324	85	18.3	20.2
ALL ACCIDENT	194	105	185	12.9	10.0
MOTOR VEHICLE	126	56	225	8.4	6.1
ALL OTHER	68	49	139	4.5	3.9
STROKE	81	117	69	5.4	6.1
C.O.P.D.	54	41	132	3.6	4.7
DIABETES MELLITUS	48	25	192	3.2	2.7
SUICIDE	52	26	200	3.5	2.8
INFLUENZA/PNEUMONIA	45	36	125	3.0	2.7
CIRRHOSIS OF LIVER	26	27	96	1.7	1.9
HOMICIDE	24	25	96	1.6	1.9
ALCOHOLISM	22	4	550	1.5	1.7
CERTAIN CONDITIONS IN PERINATAL PERIOD	33	39	85	2.2	1.6
ATHEROSCLEROSIS	19	19	100	1.3	1.1
CONGENITAL ANOMALIES	24	20	120	1.6	.9

PERCENTAGE DISTRIBUTION OF DEATHS BY AGE 1983-1985

AGE	% OF TOTAL DEATHS VALENCIA	% OF TOTAL DEATHS NEW MEXICO
ALL	100.0	100.0
< 1	3.0	2.9
1 - 4	.9	.8
5- 14	1.5	.8
15-24	5.4	3.7
25-34	3.7	4.4
35-44	3.8	4.0
45-54	6.9	6.4
55-64	15.7	13.4
65-74	23.4	22.1
75-84	24.3	24.5
85+	11.5	16.9

INFANT MORTALITY

VALENCIA COUNTY	TOTAL INFANT Number	TOTAL INFANT Rate	NEONATAL Number	NEONATAL Rate	POSTNEONATAL Number	POSTNEONATAL Rate
1980	15	10.4	9	6.2	6	4.1
1981	12	9.1	6	4.6	6	4.6
1982	8	12.9	4	6.4	4	6.4
1983	9	15.1	3	5.0	6	10.1
1984	5	7.9	2	3.2	3	4.8
1985	9	14.3	8	12.7	1	1.6
NEW MEXICO						
1985	291	10.6	170	6.2	121	4.4

Rate Per 1000 Live Births

VALENCIA COUNTY

PLACE OF RESIDENCE/1985	BIRTHS	DEATHS	BIRTHS PER 1 DEATH
Belen-Urban	143	70	2.0
Belen-Rural	74	61	1.2
Los Lunas-Urban	154	59	2.6
Los Lunas-Rural	167	37	4.5
Bosque Farms-All	41	20	2.1
Other & Unspecified	50	19	2.6
COUNTY TOTAL	629	266	2.4
% OF COUNTY RESIDENT BIRTHS:			
Born in County	21.94%	XXXXX	XXXXX
Born in Other NM County	77.74%	XXXXX	XXXXX
Born Out of State	0.32%	XXXXX	XXXXX

****The SMR's is the ratio of the observed deaths to the expected deaths times 100. The number of expected deaths is derived using the national mortality as a standard. SMR's were computed before rounding and may differ slightly from those obtained using rounded figures.*

INFANT = <1 year, NEONATAL = <28 days, POSTNEONATAL = 28 days to <1 year.

APPENDIX

TABLE A-1 New Mexico Population by City 1980 Census and 1984 Estimates . . . 138

TABLE A-2 New Mexico Births Number By County of Residence 1970-1985 . . . 139

TABLE A-3 New Mexico Resident Live Births Percentage by Age of Mother, 1970, 1980, 1983, 1984, 1985 . . . 140

TABLE A-4 New Mexico Resident Fertility Rates and Age Specific Birth Rates by County 1970, 1980, 1984, 1985 . . . 141

TABLE A-5 New Mexico Resident Live Births by Age of Mother Distribution by Live Birth Order 1985 . . . 142

TABLE A-6 New Mexico Births by County of Occurrence By Facility and by Attendant 1985 . . . 144

TABLE A-7 New Mexico Births & Deaths By Place of Residence 1985 . . . 145

TABLE A-8 New Mexico Resident Mortality Rates By County, Sex, and Age 1983-1985 . . . 147

TABLE A-9 New Mexico Resident Deaths Number and Crude Death Rate by Cause 1985 . . . 149

TABLE A-10 New Mexico Residents, Selected Causes of Death by County 1985 . . . 150

TABLE A-11 New Mexico Residents, Selected Causes of Death by County Among Males 1985 . . . 151

TABLE A-12 New Mexico Residents, Selected Causes of Death by County Among Females 1985 . . . 152

TABLE A-13 Summary of Trends in New Mexico Health Statistics 1950-1985 . . . 153

TABLE A-1
NEW MEXICO
CITY POPULATIONS
1980 CENSUS AND 1984 ESTIMATES

CITY	COUNTY	1980 POPULATION	1984 POPULATION	% CHANGE 1980-1984
Alamogordo	Otero	24,024	27,485	14.4
Albuquerque	Bernalillo	331,767	350,575	5.7
Artesia	Eddy	10,385	11,938	15.0
Aztec	San Juan	5,512	6,936	25.8
Bayard	Grant	3,036	2,994	- 1.4
Belen	Valencia	5,617	6,794	21.0
Bernalillo	Sandoval	3,012	2,736	- 9.2
Bloomfield	San Juan	4,881	6,462	32.4
Bosque Farms	Valencia	3,353	4,029	20.2
Capitan	Lincoln	762	879	15.4
Carlsbad	Eddy	25,496	28,433	11.5
Carrizozo	Lincoln	1.222	1,233	0.9
Causey	Roosevelt	81	75	- 7.4
Central	Grant	1,968	2,126	8.0
Chama	Rio Arriba	1,090	1,093	0.3
Cimarron	Colfax	888	766	-13.7
Clayton	Union	2,968	3,517	18.5
Cloudcroft	Otero	521	642	23.2
Clovis	Curry	31,194	33,424	7.1
Columbus	Luna	414	432	4.3
Corona	Lincoln	236	286	21.2
Corrales	Bernalillo/ Sandoval	2,791	2,616	- 6.3
Cuba	Sandoval	609	519	-14.8
Deming	Luna	9,964	10,609	6.5
Des Moines	Union	178	163	- 8.4
Dexter	Chaves	882	1,112	26.1
Dora	Roosevelt	168	189	12.5
Eagle Nest	Colfax	202	233	15.3
Elida	Roosevelt	202	213	5.4
Encino	Torrance	155	159	2.6
Espanola	Santa Fe/ Rio Arriba	6,803	7,450	9.5
Estancia	Torrance	830	842	1.4
Eunice	Lea	2,970	3,234	8.9
Farmington	San Juan	31,222	37,332	19.6
Floyd	Roosevelt	146	156	6.8
Folsom	Union	73	67	- 8.2
Fort Sumner	De Baca	1,421	1,341	- 5.6
Gallup	McKinley	18,161	20,959	15.4
Grady	Curry	122	138	13.1
Grants	Cibola	11,439	9,808	-14.3
Grenville	Union	39	38	- 2.6
Hagerman	Chaves	936	939	0.3
Hatch	Dona Ana	1,028	1,084	5.4
Hobbs	Lea	29,153	35,029	20.2
Hope	Eddy	111	145	30.6
House	Quay	117	108	- 7.7
Hurley	Grant	1,616	1,670	3.3
Jal	Lea	2,675	2,943	10.0
Jemez Springs	Sandoval	316	306	- 3.2
La Mesilla	Dona Ana	2,029	2,624	29.3
Las Cruces	Dona Ana	45,086	50,275	11.5
Las Vegas (City & Town)	San Miguel	14,322	15,364	7.3
Logan	Quay	735	773	5.2
Lordsburg	Hidalgo	3,195	3,202	0.2
Los Lunas	Valencia	3,525	4,832	37.1
Los Ranchos de Albuquerque	Bernalillo	2,702	3,014	11.5
Loving	Eddy	1,355	1,524	12.5
Lovington	Lea	9,727	11,704	20.3
Magdalena	Socorro	1,022	1,117	9.3
Maxwell	Colfax	316	311	- 1.6
Melrose	Curry	649	669	3.1
Milan	Cibola	3,747	3,015	-19.5
Moriarty	Torrance	1,276	1,451	13.7
Mosquero	Harding	197	174	-11.7
Mountainair	Torrance	1,170	1,102	- 5.8
Pecos	San Miguel	885	1,046	18.2
Portales	Roosevelt	9,940	10,456	5.2
Questa	Taos	1,202	1,313	9.2
Raton	Colfax	8,225	7,923	- 3.7
Red River	Taos	332	370	11.4
Reserve	Catron	439	472	7.5
Roswell	Chaves	39,676	45,702	15.2
Roy	Harding	381	316	-17.1
Ruidoso	Lincoln	4,260	5,257	23.4
Ruidoso Downs	Lincoln	949	1,312	38.3
San Jon	Quay	341	262	-23.2
Santa Fe	Santa Fe	48,953	52,274	6.8
Santa Rosa	Guadalupe	2,469	2,453	- 0.6
San Ysidro	Sandoval	199	187	- 6.0
Silver City	Grant	9,887	11,014	11.4
Socorro	Socorro	7,173	8,100	12.9
Springer	Colfax	1,657	1,568	- 5.4
Taos	Taos	3,369	3,444	2.2
Tatum	Lea	896	974	8.7
Texico	Curry	958	955	- 0.3
Tijeras	Bernalillo	311	398	28.0
Truth or Consequences	Sierra	5,219	5,590	7.1
Tucumcari	Quay	6,765	6,957	2.8
Tularosa	Otero	2,536	2,704	6.6
Vaughn	Guadalupe	737	714	- 3.1
Virden	Hidalgo	246	262	6.5
Wagon Mound	Mora	416	428	2.9
Willard	Torrance	166	176	6.0
Williamsburg	Sierra	433	406	- 6.2

SOURCE: Bureau of the Census, Bureau of Business and Economic Research, University of New Mexico.

TABLE A-2
NEW MEXICO BIRTHS BY COUNTY OF RESIDENCE
1970-1985
NUMBER OF BIRTHS BY COUNTY

		COUNTY	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985
		NEW MEXICO	22004	22205	20813	20852	21339	21078	22180	23100	23907	24821	26589	26565	27630	27508	27350	27449
		CIBOLA*	*	*	*	*	*	*	*	*	*	*	*	*	661	574	533	525
	1	MCKINLEY	1340	1409	1420	1395	1403	1481	1551	1598	1607	1625	1661	1659	1722	1669	1670	1720
		SAN JUAN	1343	1328	1281	1409	1375	1512	1647	1788	1895	2089	2316	2422	2484	2315	2240	2228
		COLFAX	213	233	220	223	235	193	201	219	206	221	239	218	248	250	212	256
		HARDING	16	12	13	10	12	8	17	10	16	15	13	17	12	16	11	16
		LOS ALAMOS	217	191	165	179	165	166	184	171	209	193	200	205	197	206	200	204
		MORA	81	86	89	82	65	69	57	64	64	78	90	79	83	77	76	72
	2	RIO ARRIBA	661	703	647	677	661	611	606	626	664	635	658	662	649	676	733	758
H		SAN MIGUEL	466	542	426	435	437	393	453	432	456	460	503	472	449	482	495	467
E		SANTA FE	1102	1129	1040	1080	1059	991	1125	1165	1209	1233	1272	1293	1218	1366	1420	1449
A		TAOS	404	429	369	370	349	353	342	370	362	398	417	414	465	450	443	458
L		UNION	84	84	83	75	75	50	72	77	74	76	75	91	89	80	84	76
T																		
H		BERNALILLO	6448	6187	5895	5929	5995	5754	6073	6461	6750	7179	7699	7357	7674	7650	7785	7937
		SANDOVAL	448	476	467	487	573	507	532	563	555	544	686	725	741	796	899	981
D	3	TORRANCE	87	69	95	92	94	106	93	103	105	109	130	127	126	146	136	136
I		VALENCIA*	920	909	852	803	854	889	1083	1226	1325	1339	1449	1313	622	595	630	629
S																		
T		CHAVES	804	869	765	661	792	676	742	880	847	922	1007	990	1138	1173	1060	1012
R		CURRY	1018	1046	944	962	1003	966	970	969	995	970	966	905	928	904	980	893
I		DE BACA	33	35	33	36	24	32	26	31	33	42	41	35	40	42	38	36
C		EDDY	775	723	655	641	694	779	823	834	891	861	982	965	1161	1019	930	885
T	4	GUADALUPE	96	103	94	70	79	69	78	75	90	72	95	85	89	92	78	78
S		LEA	980	1025	985	964	1030	1097	1076	924	1113	1145	1312	1510	1623	1596	1368	1367
		LINCOLN	124	133	150	114	167	146	158	184	151	167	169	216	239	254	246	246
		QUAY	189	198	208	177	186	171	156	156	178	180	179	157	201	165	183	176
		ROOSEVELT	290	347	299	270	274	271	258	248	276	273	290	291	330	294	268	286
		CATRON	33	31	37	39	24	27	47	55	39	45	49	41	40	37	30	34
		DONA ANA	1666	1638	1480	1564	1557	1559	1580	1595	1584	1697	1839	1947	1992	2112	2283	2140
		GRANT	512	506	478	447	490	495	478	514	512	546	548	598	546	464	470	515
	5	HIDALGO	111	113	116	108	133	142	149	144	148	130	139	112	142	144	136	107
		LUNA	238	282	264	285	276	305	283	292	230	297	296	295	325	296	291	302
		OTERO	977	1052	950	983	974	941	1006	974	999	944	931	985	1069	1163	1087	1114
		SIERRA	70	85	86	87	85	92	100	108	72	99	91	104	96	128	95	98
		SOCORRO	258	232	207	198	199	227	214	244	252	237	247	275	231	277	240	248

*Prior to 1982, Cibola County data were included in Valencia County.

TABLE A-3
NEW MEXICO RESIDENT LIVE BIRTHS
PERCENTAGE BY AGE OF MOTHER
1970, 1980, 1983, 1984, 1985

			PERCENTAGE DISTRIBUTION BY AGE OF MOTHER						
COUNTY		TOTAL	< 15	15-17	18-19	20-24	25-29	30-34	35+
NEW MEXICO	1970	22,004	.2	5.6	12.7	38.6	23.9	11.6	7.4
	1980	26,589	.2	6.5	11.4	35.9	27.6	13.8	4.6
	1983	27,508	.3	6.1	11.0	34.3	27.8	14.9	5.6
	1984	27,350	.2	5.3	10.1	34.4	28.7	15.5	5.7
	1985	27,449	.2	5.4	9.9	33.6	28.8	15.7	6.4
DISTRICT I									
CIBOLA	1970		Cibola County included in Valencia County						
	1980		prior to 1982.						
	1983	574	.7	6.3	14.5	32.8	28.4	11.8	5.6
	1984	533	.6	9.4	11.6	34.0	27.0	12.6	4.9
	1985	525	.6	6.9	11.4	32.4	26.7	15.2	6.9
MCKINLEY	1970	1,340	.1	4.3	12.5	35.6	21.9	15.4	10.3
	1980	1,661	.5	8.0	12.2	36.1	24.7	12.5	6.0
	1983	1,669	.5	7.2	10.6	34.0	25.1	14.5	8.1
	1984	1,670	.1	6.0	9.6	36.1	24.4	16.2	7.6
	1985	1,720	.3	6.7	10.2	33.1	25.1	16.2	8.4
SAN JUAN	1970	1,343	.1	4.9	10.9	36.2	26.2	12.9	8.7
	1980	2,316	.3	6.6	12.2	38.9	26.3	11.1	4.6
	1983	2,315	.0	5.3	10.6	36.2	26.9	14.7	6.3
	1984	2,240	.2	4.7	10.3	37.5	27.8	13.3	6.2
	1985	2,228	.3	4.5	9.8	34.1	28.5	16.3	6.4
DISTRICT II									
COLFAX	1970	213	--	5.2	15.5	35.7	25.8	11.3	6.6
	1980	239	--	6.3	12.6	37.7	27.6	10.0	5.9
	1983	250	.4	5.2	10.8	35.6	27.2	14.0	6.8
	1984	212	.5	5.2	10.8	33.0	31.1	14.2	5.2
	1985	256	.4	8.2	10.2	33.6	28.9	15.2	3.5
HARDING	1970	16	--	--	12.5	43.8	25.0	6.3	12.5
	1980	13	--	--	--	53.8	38.5	7.7	--
	1983	16	--	--	18.8	37.5	12.5	25.0	6.3
	1984	11	--	--	--	18.2	54.5	18.2	9.1
	1985	16	--	6.3	6.3	31.3	43.8	12.5	--
LOS ALAMOS	1970	217	--	2.8	4.6	28.6	37.3	18.0	8.8
	1980	200	--	3.0	4.5	20.0	34.0	28.0	10.5
	1983	206	--	2.4	3.4	20.9	35.0	28.2	10.2
	1984	200	--	.5	2.0	16.5	38.5	31.5	11.0
	1985	204	--	.5	2.9	22.5	36.8	24.5	12.7
MORA	1970	81	--	7.4	14.8	44.4	12.3	12.3	8.6
	1980	90	1.1	12.2	13.3	26.7	27.8	14.4	4.4
	1983	77	--	7.8	13.0	37.7	24.7	11.7	5.2
	1984	76	--	3.9	7.9	36.8	27.6	17.1	6.6
	1985	72	--	4.2	16.7	36.1	27.8	11.1	4.2
RIO ARRIBA	1970	661	.2	4.2	11.3	36.8	23.0	13.9	10.6
	1980	658	.6	7.8	12.8	35.3	25.2	12.5	5.9
	1983	676	.4	6.7	14.1	33.0	25.6	14.6	5.6
	1984	733	.5	7.5	13.6	30.6	28.5	14.1	5.2
	1985	758	.1	7.0	13.2	34.0	24.3	13.1	8.3
SAN MIGUEL	1970	466	.2	6.9	13.1	42.1	20.6	9.4	7.7
	1980	503	.2	8.3	9.9	37.0	25.4	14.7	4.4
	1983	482	.4	6.6	12.7	36.9	24.1	14.5	4.9
	1984	495	.2	5.9	10.3	38.2	26.1	12.9	6.5
	1985	467	.4	4.9	7.9	34.7	30.6	15.0	6.4
SANTA FE	1970	1,102	.1	4.6	12.0	36.0	24.5	13.3	9.2
	1980	1,272	.1	5.5	9.0	27.8	32.4	17.9	7.3
	1983	1,366	--	5.5	8.3	28.4	26.4	22.8	8.6
	1984	1,420	.1	3.3	8.0	29.9	27.4	22.0	9.4
	1985	1,449	.1	5.2	7.8	28.4	27.3	20.4	10.8
TAOS	1970	404	--	5.9	10.9	37.1	22.3	10.9	12.9
	1980	417	--	5.3	12.0	29.7	29.0	18.9	5.0
	1983	450	.7	5.1	10.9	26.9	27.3	20.9	8.2
	1984	443	--	4.7	8.6	33.6	26.4	17.4	9.3
	1985	458	.2	4.8	8.5	29.5	26.6	22.3	8.1
UNION	1970	84	--	3.6	9.5	44.0	26.2	10.7	6.0
	1980	75	--	8.0	13.3	33.3	24.0	16.0	5.3
	1983	80	--	6.3	8.8	32.5	36.3	11.3	5.0
	1984	84	--	8.3	11.9	29.8	32.1	14.3	3.6
	1985	76	--	7.9	13.2	26.3	32.9	11.8	7.9
DISTRICT III									
BERNALILLO	1970	6,448	.2	5.5	12.2	38.2	27.0	10.8	6.2
	1980	7,699	.2	5.6	10.1	33.4	30.9	16.0	3.8
	1983	7,650	.3	5.0	9.4	32.6	30.2	17.1	5.4
	1984	7,785	.3	4.4	9.5	31.1	30.7	18.3	5.8
	1985	7,937	.2	4.4	9.0	31.8	30.1	18.0	6.5
SANDOVAL	1970	448	.4	7.6	10.0	32.6	24.3	13.6	11.4
	1980	686	.3	7.6	10.2	29.4	29.7	16.9	5.8
	1983	796	.4	4.5	9.0	31.5	31.8	16.8	5.9
	1984	899	--	4.3	8.5	31.6	29.7	18.1	7.8
	1985	981	.2	3.7	7.0	29.5	33.6	19.8	6.2
TORRANCE	1970	87	--	12.6	10.3	36.8	14.9	11.5	13.8
	1980	130	.8	3.8	15.4	32.3	28.5	12.3	6.9
	1983	146	--	6.8	10.3	32.9	26.0	16.4	7.5
	1984	136	--	2.9	11.8	37.5	24.3	18.4	5.1
	1985	136	--	5.9	9.6	33.1	32.4	12.5	6.6
VALENCIA	1970	920	.1	5.0	13.0	38.2	23.5	11.2	9.0
	1980	1,449	.2	7.9	11.0	38.1	25.3	13.0	4.5
	1983	595	.3	6.9	12.4	31.1	28.7	13.9	6.6
	1984	630	--	6.8	12.1	32.1	30.0	13.8	5.2
	1985	629	.2	5.6	7.3	33.5	30.8	14.9	7.6
DISTRICT IV									
CHAVES	1970	804	.1	8.6	15.3	36.9	22.6	10.9	5.5
	1980	1,007	.8	7.5	11.8	36.7	24.8	12.7	5.6
	1983	1,173	.3	6.1	12.9	37.9	28.1	10.1	4.6
	1984	1,060	.4	7.7	10.9	37.8	25.8	13.2	4.2
	1985	1.012	.6	7.6	10.1	38.5	26.6	12.4	4.2
CURRY	1970	1,018	.1	4.6	14.7	46.4	19.4	9.9	4.8
	1980	966	.5	6.0	15.1	43.6	21.3	10.1	3.3
	1983	904	.3	7.4	13.6	43.0	23.8	8.6	3.2
	1984	980	.2	6.4	11.8	44.2	25.7	9.5	2.1
	1985	893	.3	5.7	12.3	42.7	26.1	10.2	2.7
DE BACA	1970	33	--	12.1	9.1	30.3	21.2	18.2	9.1
	1980	41	--	4.9	14.6	36.6	24.4	12.2	7.3
	1983	42	--	7.1	19.0	31.0	26.2	11.9	4.8
	1984	38	--	10.5	10.5	34.2	26.3	15.8	2.6
	1985	36	--	8.3	22.2	30.6	30.6	2.8	5.6
EDDY	1970	775	.1	9.4	16.3	39.5	18.8	9.0	6.8
	1980	982	.1	8.7	12.6	38.4	26.8	9.9	3.6
	1983	1,019	.4	8.2	12.6	35.5	27.3	11.9	4.1
	1984	930	.3	7.5	12.5	37.0	26.9	12.3	3.5
	1985	885	.1	6.8	12.5	35.1	30.3	10.2	5.0
GUADALUPE	1970	96	--	9.4	13.5	34.4	14.6	13.5	14.6
	1980	95	--	6.3	23.2	31.6	20.0	11.6	7.4
	1983	92	--	12.0	8.7	43.5	20.7	10.9	4.3
	1984	78	--	12.8	12.8	33.3	24.4	14.1	2.6
	1985	78	--	9.0	15.4	35.9	20.5	11.5	7.7
LEA	1970	980	.3	9.0	15.9	42.9	19.3	8.0	4.7
	1980	1,312	.1	8.1	13.6	43.0	22.5	9.2	3.6
	1983	1,596	.5	8.0	14.1	39.5	25.3	9.3	3.3
	1984	1,368	.3	6.8	10.6	41.2	26.9	11.3	2.9
	1985	1,367	.4	7.0	10.5	36.1	31.5	11.1	3.3
LINCOLN	1970	124	--	4.8	15.3	41.1	22.6	11.3	4.8
	1980	169	--	7.7	9.5	32.5	32.5	14.2	3.6
	1983	254	.4	3.9	9.4	35.0	29.9	14.2	7.1
	1984	246	--	2.4	14.6	30.5	31.3	17.5	3.7
	1985	246	--	3.7	7.7	39.0	29.3	14.6	5.7
QUAY	1970	189	.5	9.0	15.9	39.2	19.6	10.6	5.3
	1980	179	1.1	8.4	17.9	34.1	26.8	9.5	2.2
	1983	165	.6	7.3	14.5	35.2	23.6	11.5	7.3
	1984	183	.5	8.7	15.3	36.6	27.3	9.3	2.2
	1985	176	--	9.7	21.6	30.7	25.6	9.1	3.4
ROOSEVELT	1970	290	.7	5.2	18.6	48.6	18.6	3.4	4.8
	1980	290	1.0	6.9	17.2	39.7	22.4	10.3	2.4
	1983	294	--	7.8	10.2	36.4	26.9	12.6	6.1
	1984	268	--	5.2	10.4	41.8	29.5	9.0	4.1
	1985	286	.7	5.6	12.9	32.9	25.9	15.4	6.6
DISTRICT V									
CATRON	1970	33	--	6.1	12.1	36.4	30.3	9.1	6.1
	1980	49	--	8.2	14.3	32.7	26.5	14.3	4.1
	1983	37	--	--	18.9	24.3	21.6	29.7	5.4
	1984	30	--	--	16.7	30.0	13.3	20.0	20.0
	1985	34	--	--	8.8	32.4	29.4	14.7	14.7
DONA ANA	1970	1,666	.1	4.1	11.8	37.7	23.5	13.6	9.3
	1980	1,839	.2	4.1	10.5	36.4	29.4	14.0	5.4
	1983	2,112	.5	6.2	10.1	32.3	29.8	15.2	6.0
	1984	2,283	.3	4.7	9.1	32.2	32.3	15.6	5.7
	1985	2,140	.2	5.1	10.4	35.3	29.0	13.8	6.3
GRANT	1970	512	.2	4.9	12.1	41.2	24.0	11.1	6.4
	1980	548	.2	5.8	12.8	37.4	22.8	16.6	4.4
	1983	464	.4	6.5	13.6	30.4	26.1	15.7	7.3
	1984	470	.2	6.6	10.6	37.0	25.5	13.8	6.2
	1985	515	.2	7.8	12.6	32.0	27.4	13.4	6.6
HIDALGO	1970	111	--	6.3	8.1	40.5	27.0	6.3	11.7
	1980	139	--	6.5	9.4	44.6	24.5	12.9	2.2
	1983	144	--	12.5	10.4	40.3	23.6	10.4	2.8
	1984	136	.7	5.9	9.6	38.2	25.0	9.6	11.0
	1985	107	--	10.3	9.3	38.3	29.0	8.4	4.7
LUNA	1970	238	--	8.0	10.9	35.7	21.8	15.5	8.0
	1980	296	.3	7.8	13.9	41.6	16.9	14.2	5.4
	1983	296	--	11.1	15.5	32.8	23.0	11.8	5.7
	1984	291	.3	6.5	14.4	34.7	28.9	8.6	6.5
	1985	302	--	9.6	13.6	32.8	24.2	12.6	7.3
OTERO	1970	977	.2	4.6	12.0	44.1	22.6	11.8	4.7
	1980	931	--	5.6	11.1	41.2	27.5	10.8	3.8
	1983	1,163	--	5.9	11.8	43.1	24.4	11.6	3.2
	1984	1,087	.3	4.4	11.6	40.4	28.2	11.4	3.7
	1985	1,114	.2	3.4	10.6	42.6	27.8	11.7	3.7
SIERRA	1970	70	--	10.0	7.1	32.9	30.0	11.4	8.6
	1980	91	--	7.7	16.5	39.6	28.6	4.4	3.3
	1983	128	--	9.4	15.6	32.8	27.3	10.2	4.7
	1984	95	--	10.5	11.6	42.1	20.0	6.3	9.5
	1985	98	--	2.0	14.3	34.7	29.6	16.3	3.1
SOCORRO	1970	258	.4	4.3	16.3	31.4	23.6	13.6	10.5
	1980	247	.4	8.5	13.4	35.2	24.7	12.1	5.7
	1983	277	.4	4.3	12.6	32.5	27.8	17.0	5.4
	1984	240	.4	6.3	6.7	36.3	29.2	16.3	5.0
	1985	248	--	6.5	12.1	31.9	26.2	18.1	5.2

TABLE A-4
NEW MEXICO RESIDENT FERTILITY RATES
BIRTH RATES BY AGE OF MOTHER BY COUNTY 1970, 1980, 1984, 1985

COUNTY	FERTILITY RATE (a)	BIRTH RATES BY AGE GROUP (b)								
			15-19							
		10-14	TOTAL	15-17	18-19	20-24	25-29	30-34	35-39	40-49
NEW MEXICO:										
1970	100.8	.6	76.7	37.2	145.8	201.9	157.7	82.3	41.1	7.0
1980	85.4	1.2	72.8	43.9	115.9	151.8	126.9	72.5	25.0	3.5
1984	80.3	1.1	59.2	34.0	96.6	136.6	124.1	76.6	30.8	3.1
1985	80.0	1.1	72.6	42.9	116.2	140.0	124.0	71.5	28.2	.1
CIBOLA:										
1970										
1980	(Prior to 1982, Cibola County Data is included in Valencia County)									
1984	83.2	2.4	82.4	56.8	129.2	125.3	120.9	66.1	28.9	3.6
1985	99.0	2.5	99.1	61.9	155.0	263.2	162.6	67.6	33.0	--
1 MCKINLEY:										
1970	142.2	.3	95.2	36.2	214.4	269.3	187.6	147.2	79.0	19.9
1980	118.7	2.7	101.3	63.9	164.4	207.8	157.9	100.0	48.0	8.1
1984	110.0	.6	72.3	44.1	119.8	191.9	144.1	119.9	56.4	8.9
1985	111.3	1.9	96.4	62.4	149.0	176.7	153.3	106.2	57.3	.3
SAN JUAN:										
1970	118.0	.5	75.3	33.4	173.6	254.7	204.3	102.1	52.9	9.5
1980	118.2	1.5	101.9	57.8	172.4	223.3	166.5	84.3	33.7	6.6
1984	97.6	.9	67.1	34.0	119.9	177.4	145.2	83.3	40.4	5.2
1985	105.1	1.5	82.7	44.0	139.2	182.3	160.2	99.1	40.0	.2
COLFAX:										
1970	100.8	--	88.9	31.6	224.5	216.5	170.8	89.6	35.4	4.2
1980	85.8	--	76.5	39.6	143.5	171.8	126.7	52.3	33.3	1.7
1984	77.5	1.8	59.1	29.6	112.7	136.7	129.4	66.8	26.2	1.7
1985	83.4	1.7	79.4	60.2	107.0	145.3	150.1	79.1	14.2	--
HARDING:										
1970	69.6	--	29.9	--	166.7	200.0	148.1	47.6	27.0	12.2
1980	69.1	--	--	--	--	250.0	142.9	24.4	--	--
1984	65.1	--	--	--	--	80.0	193.5	54.1	66.7	--
1985	98.8	--	66.7	58.8	76.9	147.1	411.8	71.4	--	--
LOS ALAMOS:										
1970	65.1	--	23.2	10.5	83.3	192.5	157.6	68.2	23.4	3.9
1980	49.7	--	19.9	8.9	46.6	96.6	119.5	77.0	24.3	--
1984	45.9	--	6.1	1.6	19.1	73.7	125.0	80.1	22.5	.7
1985	50.0	--	8.8	2.3	16.7	66.1	188.4	83.8	33.0	--
MORA:										
1970	102.4	--	69.7	32.8	146.3	238.4	109.9	117.6	55.6	4.7
1980	111.7	4.8	100.0	70.1	164.4	200.0	174.8	97.7	43.5	--
1984	84.0	--	34.5	16.9	72.3	205.9	129.6	86.1	38.5	4.7
1985	82.1	--	83.8	30.3	150.0	130.0	168.1	51.0	21.6	--
2 RIO ARRIBA:										
1970	130.0	.6	72.0	31.1	170.8	260.5	207.1	128.1	67.4	18.4
1980	95.9	2.8	82.8	47.5	150.8	176.3	143.4	76.5	38.5	4.3
1984	97.6	2.5	86.7	46.7	163.7	155.1	164.6	87.6	36.1	2.6
1985	99.4	.7	110.6	65.4	174.8	163.3	133.0	83.5	48.1	.6
SAN MIGUEL:										
1970	103.2	.8	70.8	44.7	102.0	186.7	157.9	82.6	44.5	13.9
1980	94.2	1.0	70.3	55.4	90.7	172.0	141.6	90.0	29.9	2.9
1984	85.9	1.0	56.8	35.5	86.0	159.8	132.6	72.3	36.5	6.3
1985	75.9	2.1	51.4	33.3	77.7	111.0	133.6	80.0	29.7	--
SANTA FE:										
1970	98.3	.3	70.5	29.8	149.5	195.3	156.1	85.3	47.8	9.1
1980	67.9	.3	52.3	32.1	84.9	113.2	112.8	63.2	29.3	3.8
1984	70.2	.3	42.3	20.0	78.6	125.5	98.5	80.0	38.6	5.0
1985	69.5	.6	59.3	41.0	84.6	114.5	116.6	74.3	34.2	--
TAOS:										
1970	114.2	--	70.5	36.7	141.5	235.5	168.5	87.3	90.3	12.3
1980	91.8	--	77.2	35.5	159.7	174.4	140.7	90.9	24.9	5.2
1984	91.3	--	59.2	31.7	113.4	196.1	127.3	82.9	50.9	5.8
1985	90.5	1.1	70.0	43.1	108.0	174.2	157.4	95.7	29.9	.9
UNION:										
1970	96.3	--	50.9	18.4	150.9	296.0	178.9	73.8	39.1	--
1980	81.3	--	70.8	36.8	158.7	173.6	123.3	82.8	14.3	8.5
1984	80.1	--	66.1	37.0	138.9	152.4	162.7	72.7	18.9	--
1985	70.2	--	85.1	54.1	129.9	74.6	140.4	62.9	13.0	3.9
BERNALILLO:										
1970	89.4	.7	69.0	34.9	122.9	166.7	149.1	69.7	31.8	4.6
1980	71.6	.7	58.0	35.9	88.5	114.8	113.4	68.2	17.7	2.1
1984	67.5	1.3	48.3	26.6	78.3	100.8	106.3	73.4	27.9	1.8
1985	66.5	.8	62.3	34.7	102.6	113.1	98.1	63.9	24.7	--
SANDOVAL:										
1970	127.7	1.7	85.2	54.9	146.1	225.7	206.0	124.2	78.5	15.3
1980	86.7	1.2	73.1	48.6	117.1	148.0	142.5	80.2	27.9	6.0
1984	112.0	--	67.7	35.8	124.8	204.3	183.0	110.7	54.8	5.9
1985	98.1	1.1	55.4	32.2	88.8	170.4	206.8	108.0	28.9	--
3 TORRANCE:										
1970	91.2	--	73.8	57.6	112.5	200.0	100.8	82.0	58.8	11.5
1980	84.8	2.8	82.2	26.0	178.6	165.4	131.7	56.9	34.1	5.4
1984	81.4	--	59.9	19.0	130.1	182.8	107.1	81.2	31.1	--
1985	77.1	--	56.3	35.9	86.7	171.7	164.2	51.1	23.4	--
VALENCIA:										
1970	107.9	.4	80.1	33.5	171.9	230.9	158.1	78.4	58.4	7.4
1980	101.2	1.0	88.3	56.8	146.0	194.2	138.0	81.7	27.5	5.0
1984	77.0	--	66.2	37.0	119.7	154.1	132.0	59.7	20.6	4.1
1985	77.9	.6	52.4	37.6	74.6	136.3	156.7	65.1	33.1	.5
CHAVES:										
1970	92.2	.4	87.3	47.1	167.6	209.9	146.7	68.9	30.4	2.3
1980	94.4	3.9	80.8	51.8	125.9	184.3	133.1	76.7	31.5	5.4
1984	87.3	1.7	71.6	48.8	107.1	174.3	126.9	73.3	23.5	2.7
1985	85.8	3.0	81.2	60.1	110.6	169.3	141.4	59.4	20.0	--
CURRY:										
1970	123.4	.4	113.5	42.8	235.1	258.9	155.8	76.3	32.1	5.3
1980	100.1	2.8	104.1	48.5	191.1	205.7	115.0	66.3	22.9	2.0
1984	97.7	1.1	87.5	50.5	145.4	202.5	134.7	60.2	14.9	1.0
1985	96.8	1.7	98.5	51.3	171.9	248.7	142.5	55.7	13.1	--
DE BACA:										
1970	82.1	--	72.9	53.3	136.4	200.0	114.8	107.1	43.5	--
1980	93.0	--	77.7	30.3	162.2	223.9	131.6	75.8	27.4	7.9
1984	91.1	--	82.5	64.5	114.3	206.3	138.9	96.8	14.5	--
1985	80.7	--	123.6	56.6	222.2	118.3	180.3	14.1	16.4	--
EDDY:										
1970	94.0	.4	89.8	48.6	176.3	233.8	131.4	64.9	33.2	4.7
1980	96.9	.5	95.6	63.4	146.7	188.6	134.9	63.0	23.4	2.1
1984	81.3	1.4	75.5	46.3	121.8	152.8	113.8	65.7	19.4	1.9
1985	80.4	.5	82.2	49.0	129.8	157.0	142.4	47.8	20.8	--
4 GUADALUPE:										
1970	107.6	--	91.3	50.8	203.1	217.1	116.7	102.4	58.0	26.7
1980	95.8	--	101.4	35.5	205.6	167.6	119.5	78.6	53.6	4.1
1984	79.3	--	73.0	59.5	94.3	146.1	120.3	79.1	18.0	--
1985	82.1	--	96.4	61.4	144.6	133.3	108.1	60.0	38.2	--
LEA:										
1970	89.5	1.0	94.0	50.8	181.0	246.8	122.2	49.1	20.4	3.4
1980	105.8	.4	110.9	67.2	181.0	218.4	126.7	62.6	25.0	3.1
1984	97.9	1.5	82.5	52.4	131.0	193.5	140.4	71.2	17.5	3.0
1985	99.5	2.1	105.3	69.8	159.3	199.4	160.7	59.0	16.5	--
LINCOLN:										
1970	88.6	--	73.3	24.8	191.9	210.7	123.3	72.5	34.1	--
1980	72.6	--	67.9	44.1	121.2	154.5	113.9	55.0	13.9	1.8
1984	85.7	--	79.8	16.5	220.9	170.8	129.4	80.1	20.3	--
1985	81.0	--	59.2	32.8	95.5	203.0	152.2	63.5	15.9	1.2
QUAY:										
1970	95.8	1.7	92.3	52.8	160.4	218.3	123.3	70.7	40.0	--
1980	85.1	4.4	104.7	48.5	228.6	182.6	133.7	47.6	13.2	--
1984	87.0	2.2	97.6	51.4	193.6	199.4	138.5	47.4	9.8	1.9
1985	91.8	--	181.5	91.9	322.0	178.2	148.5	39.7	12.4	--
ROOSEVELT:										
1970	76.7	2.7	59.5	34.2	74.8	125.8	133.0	28.5	26.5	5.3
1980	75.3	5.3	69.0	47.5	84.2	108.5	117.5	64.9	16.1	1.4
1984	65.9	--	38.8	31.2	44.2	99.0	113.9	48.7	17.6	5.3
1985	70.5	4.0	87.7	44.8	149.8	93.3	73.6	87.5	25.8	--
CATRON:										
1970	85.7	--	67.4	29.4	166.7	200.0	144.9	60.0	40.0	--
1980	88.3	--	83.3	47.8	145.8	238.8	122.6	74.5	22.2	--
1984	53.0	--	37.0	--	102.0	132.4	37.0	62.5	65.2	--
1985	54.8	--	25.9	--	56.6	85.9	144.9	47.6	31.3	6.5
DONA ANA:										
1970	104.0	.2	67.5	31.0	115.1	171.2	163.7	105.8	57.2	10.8
1980	75.4	.9	47.5	26.7	68.5	110.9	130.8	75.9	30.1	4.0
1984	80.3	1.2	47.9	32.6	63.4	104.8	153.3	90.2	36.5	3.2
1985	74.3	.8	59.6	32.9	99.1	109.3	102.6	77.6	32.7	.2
GRANT:										
1970	111.4	.8	77.7	37.9	134.8	223.8	171.5	99.1	34.7	8.6
1980	94.8	.8	78.9	40.2	141.1	194.3	122.8	91.1	24.0	4.0
1984	77.0	.8	59.4	36.9	95.6	156.3	111.7	61.7	28.7	3.8
1985	81.0	.8	87.2	54.0	140.4	137.0	140.4	68.7	28.9	1.3
5 HIDALGO:										
1970	124.3	--	74.4	47.3	134.3	306.1	208.3	54.7	92.4	7.7
1980	104.5	--	71.7	43.1	132.7	247.0	148.5	81.4	18.4	--
1984	96.2	3.1	64.8	36.2	126.2	196.2	140.5	55.8	81.4	3.3
1985	76.3	--	72.9	62.9	88.5	156.5	147.6	44.1	12.6	--
LUNA:										
1970	106.9	--	78.7	48.7	142.9	234.2	181.8	117.1	42.0	6.2
1980	102.1	1.5	96.2	51.6	187.2	236.1	114.2	89.9	30.5	5.1
1984	94.5	1.4	86.2	40.0	180.3	182.0	180.3	50.3	26.2	9.5
1985	100.2	--	120.1	84.5	170.8	169.8	150.2	78.2	37.0	--
OTERO:										
1970	108.4	.8	88.1	36.3	194.7	251.5	152.0	75.9	24.3	5.6
1980	86.9	--	69.4	38.1	118.3	174.1	127.2	60.6	20.4	3.0
1984	91.3	1.5	70.2	31.8	130.6	179.6	137.6	67.2	24.3	1.2
1985	96.5	1.1	81.9	33.2	155.5	206.1	140.7	64.9	21.1	--
SIERRA:										
1970	67.3	--	49.6	40.7	71.4	153.3	114.8	54.8	33.8	2.8
1980	74.7	--	83.7	38.6	180.7	175.6	148.6	20.6	15.5	--
1984	73.8	--	75.5	52.6	125.0	184.3	102.7	29.3	29.3	7.4
1985	73.0	--	50.5	11.2	101.4	149.1	156.8	97.6	14.6	--
SOCORRO:										
1970	131.8	1.8	105.6	37.2	203.9	210.4	213.3	135.1	66.4	16.6
1980	89.0	1.8	87.0	59.7	122.7	141.4	123.5	76.2	23.5	10.7
1984	74.7	1.5	43.1	36.9	51.4	123.6	122.4	82.8	25.4	3.1
1985	84.5	--	78.4	45.2	128.8	134.6	110.7	92.0	30.7	--

New Mexico rates based on 1970 and 1980 Census figures, 1980 age composition as applied to 1984 total state and county population estimates furnished by the Bureau of the Census. 1985 based on 1985 BBER projections by age and sex adjusted to estimated totals by state and county.

(a) Births Per 1,000 Women Ages 15-44

(b) Also referred to as Age-Specific Fertility Rates.
Birth rates by age group per 1,000 women.

TABLE A-5
NEW MEXICO RESIDENT
LIVE BIRTHS BY AGE OF MOTHER
1985
Percentage Distribution By Live Birth Order

COUNTY	TOTAL BIRTHS	LIVE BIRTH ORDER 1	2	3	4	>5
DISTRICT I						
CIBOLA COUNTY	525	35.2	29.7	19.2	9.3	6.5
< 15	3	100.0	--	--	--	--
15-17	36	86.1	11.1	2.8	--	--
18-19	60	81.7	13.3	5.0	--	--
(15-19)	(96)	(83.3)	(12.5)	(4.2)	(--)	(--)
20-24	170	37.6	42.9	14.1	3.5	1.8
25-29	140	18.6	34.3	32.1	12.1	2.9
30-34	80	12.5	26.3	26.3	20.0	15.0
35-39	32	6.3	6.3	18.8	31.3	37.5
> 40	4	--	--	25.0	--	75.0
MCKINLEY COUNTY	1720	32.8	26.0	18.8	11.2	11.2
< 15	6	100.0	--	--	--	--
15-17	115	87.0	12.2	0.9	--	--
18-19	176	69.3	23.3	7.4	--	--
(15-19)	(291)	(76.3)	(18.9)	(4.8)	(--)	(--)
20-24	569	36.9	38.0	18.3	6.0	0.9
25-29	432	18.5	26.4	26.6	16.4	12.0
30-34	278	11.9	18.0	24.5	20.5	25.2
35-39	121	10.7	9.1	18.2	22.3	39.7
> 40	23	--	4.3	4.3	13.0	78.3
SAN JUAN COUNTY	2228	32.8	31.7	18.3	9.2	8.0
< 15	6	100.0	--	--	--	--
15-17	101	90.1	8.9	1.0	--	--
18-19	219	68.5	27.9	3.7	--	--
(15-19)	(320)	(75.3)	(21.9)	(2.8)	(--)	(--)
20-24	760	37.9	40.0	14.7	5.7	1.7
25-29	636	20.1	35.2	25.9	12.3	6.4
30-34	364	14.6	24.7	25.3	15.4	20.1
35-39	123	10.6	11.4	22.0	20.3	35.8
> 40	19	5.3	21.1	15.8	15.8	42.1
DISTRICT II						
COLFAX COUNTY	256	41.8	30.1	19.9	5.5	2.7
< 15	1	100.0	--	--	--	--
15-17	21	81.0	14.3	--	4.8	--
18-19	26	65.4	30.8	3.8	--	--
(15-19)	(47)	(72.3)	(23.4)	(2.1)	(2.1)	(--)
20-24	86	43.0	37.2	17.4	2.3	--
25-29	74	33.8	29.7	24.3	9.5	2.7
30-34	39	17.9	25.6	41.0	5.1	10.3
35-39	7	42.9	28.6	14.3	14.3	--
> 40	2	--	--	--	50.0	50.0
HARDING COUNTY	16	43.8	37.5	12.5	6.3	--
< 15	0	--	--	--	--	--
15-17	1	100.0	--	--	--	--
18-19	1	--	100.0	--	--	--
(15-19)	(2)	(50.0)	(50.0)	(--)	(--)	(--)
20-24	5	40.0	60.0	--	--	--
25-29	7	57.1	14.3	14.3	14.3	--
30-34	2	--	50.0	50.0	--	--
35-39	0	--	--	--	--	--
> 40	0	--	--	--	--	--
LOS ALAMOS COUNTY	204	40.7	39.7	14.2	3.4	2.0
< 15	0	--	--	--	--	--
15-17	1	100.0	--	--	--	--
18-19	6	83.3	16.7	--	--	--
(15-19)	(7)	(85.7)	(14.3)	(--)	(--)	(--)
20-24	46	52.2	39.1	6.5	2.2	--
25-29	75	44.0	40.0	16.0	--	--
30-34	50	32.0	40.0	14.0	10.0	4.0
35-39	23	17.4	43.5	30.4	4.3	4.3
> 40	3	--	66.7	--	--	33.3
DISTRICT II						
MORA COUNTY	72	40.3	31.9	13.9	6.9	6.9
< 15	0	--	--	--	--	--
15-17	3	66.7	33.3	--	--	--
18-19	12	75.0	8.3	16.7	--	--
(15-19)	(15)	(73.0)	(13.3)	(13.3)	(--)	(--)
20-24	26	42.3	46.2	7.7	3.8	--
25-29	20	25.0	40.0	15.0	10.0	10.0
30-34	8	12.5	12.5	12.5	25.0	37.5
35-39	3	33.3	--	66.7	--	--
> 40	0	--	--	--	--	--
RIO ARRIBA COUNTY	758	41.3	30.7	14.9	7.1	5.9
< 15	1	100.0	--	--	--	--
15-17	53	88.7	9.4	1.9	--	--
18-19	100	84.0	14.0	2.0	--	--
(15-19)	(153)	(85.6)	(12.4)	(2.0)	(--)	(--)
20-24	258	45.3	36.8	14.3	2.3	1.2
25-29	184	23.9	40.8	20.7	11.4	3.3
30-34	99	13.1	32.3	23.2	15.2	16.2
35-39	57	10.5	21.1	21.1	17.5	29.8
> 40	6	16.7	--	--	33.3	50.0

COUNTY	TOTAL BIRTHS	LIVE BIRTH ORDER 1	2	3	4	>5
DISTRICT II (continued)						
SAN MIGUEL COUNTY	467	36.0	33.4	20.3	6.2	4.1
< 15	2	100.0	--	--	--	--
15-17	23	100.0	--	--	--	--
18-19	37	75.7	21.6	--	2.7	--
(15-19)	(60)	(85.0)	(13.3)	(--)	(1.7)	(--)
20-24	162	45.1	38.3	13.0	3.1	0.6
25-29	143	16.8	36.4	35.7	9.1	2.1
30-34	70	20.0	41.4	27.1	4.3	7.1
35-39	26	15.4	15.4	15.4	26.9	26.9
> 40	4	--	25.0	--	--	75.0
SANTA FE COUNTY	1449	44.2	33.7	13.9	5.2	3.0
< 15	2	100.0	--	--	--	--
15-17	76	92.1	7.9	--	--	--
18-19	113	85.0	10.6	4.4	--	--
(15-19)	(189)	(87.8)	(9.5)	(2.6)	(--)	(--)
20-24	411	51.1	37.5	9.2	1.5	0.7
25-29	395	34.4	39.2	20.3	4.6	1.5
30-34	296	30.7	35.8	17.6	9.8	6.1
35-39	133	23.3	36.1	18.8	13.5	8.3
> 40	23	21.7	30.4	4.3	17.4	26.1
TAOS COUNTY	458	38.4	37.1	15.5	6.8	2.2
< 15	1	100.0	--	--	--	--
15-17	22	90.9	9.1	--	--	--
18-19	39	74.4	25.6	--	--	--
(15-19)	(61)	(80.3)	(19.7)	(--)	(--)	(--)
20-24	135	43.0	44.4	11.1	1.5	--
25-29	122	32.0	45.1	16.4	4.9	1.6
30-34	102	20.6	32.4	25.5	18.6	2.9
35-39	29	20.7	24.1	31.0	10.3	13.8
> 40	8	25.0	37.5	12.5	12.5	12.5
UNION COUNTY	76	32.9	28.9	23.7	10.5	3.9
< 15	0	--	--	--	--	--
15-17	6	100.0	--	--	--	--
18-19	10	70.0	30.0	--	--	--
(15-19)	(16)	(81.3)	(18.8)	(--)	(--)	(--)
20-24	20	40.0	40.0	20.0	--	--
25-29	25	12.0	28.0	36.0	24.0	--
30-34	9	--	44.4	22.2	11.1	22.2
35-39	2	50.0	--	50.0	--	--
> 40	4	--	--	50.0	25.0	25.0
DISTRICT III						
BERNALILLO COUNTY	7937	45.4	33.1	14.1	4.9	2.6
< 15	13	100.0	--	--	--	--
15-17	351	88.0	11.4	0.6	--	--
18-19	714	73.8	22.5	3.4	0.3	--
(15-19)	(1065)	(78.5)	(18.9)	(2.4)	(0.2)	(--)
20-24	2524	50.6	34.7	11.3	2.7	0.6
25-29	2393	38.1	36.6	17.3	6.1	2.0
30-34	1425	29.7	37.3	20.2	7.8	5.0
35-39	469	27.5	28.8	19.6	11.7	12.4
> 40	48	27.1	10.4	25.0	12.5	25.0
SANDOVAL COUNTY	981	38.9	30.5	18.1	7.1	5.3
< 15	2	100.0	--	--	--	--
15-17	36	86.1	11.1	2.8	--	--
18-19	69	79.7	17.4	2.9	--	--
(15-19)	(105)	(81.9)	(15.2)	(2.9)	(--)	(--)
20-24	289	48.4	34.3	14.5	2.8	--
25-29	330	30.6	34.5	22.7	7.6	4.5
30-34	194	23.2	29.9	23.2	13.4	10.3
35-39	49	10.2	24.5	22.4	18.4	24.5
> 40	12	25.0	--	16.7	16.7	41.7
TORRANCE COUNTY	136	38.2	33.1	19.1	5.1	4.4
< 15	0	--	--	--	--	--
15-17	8	100.0	--	--	--	--
18-19	13	69.2	30.8	--	--	--
(15-19)	(21)	(81.0)	(19.0)	(--)	(--)	(--)
20-24	45	46.7	37.8	11.1	4.4	--
25-29	44	22.7	38.6	34.1	2.3	2.3
30-34	17	17.6	23.5	35.3	5.9	17.6
35-39	7	14.3	42.9	--	28.6	14.3
> 40	2	--	--	--	50.0	50.0
VALENCIA COUNTY	629	38.5	31.8	18.0	7.8	4.0
< 15	1	100.0	--	--	--	--
15-17	35	97.1	2.9	--	--	--
18-19	46	71.7	21.7	6.5	--	--
(15-19)	(81)	(82.7)	(13.6)	(3.7)	(--)	(--)
20-24	211	45.5	36.5	13.3	3.3	1.4
25-29	194	26.8	40.7	23.2	7.2	2.1
30-34	94	22.3	24.5	28.7	18.1	6.4
35-39	41	12.2	22.0	22.0	24.4	19.5
> 40	7	--	14.3	14.3	14.3	57.1

TABLE A-5
NEW MEXICO RESIDENT
LIVE BIRTHS BY AGE OF MOTHER
1985
Percentage Distribution By Live Birth Order

COUNTY	TOTAL BIRTHS	LIVE BIRTH ORDER				
		1	2	3	4	5
DISTRICT IV						
CHAVES COUNTY	1012	35.9	35.4	17.2	6.4	5.1
< 15	6	100.0	--	--	--	--
15-17	77	83.1	16.9	--	--	--
18-19	102	62.7	29.4	6.9	1.0	--
(15-19)	(179)	(71.5)	(24.0)	(3.9)	(0.6)	(--)
20-24	390	34.9	40.8	17.2	4.6	2.6
25-29	269	23.4	40.1	23.0	6.3	7.1
30-34	125	20.8	33.6	20.8	13.6	11.2
35-39	36	11.1	16.7	22.2	27.8	22.2
> 40	7	--	--	57.1	28.6	14.3
CURRY COUNTY	893	50.6	29.6	12.2	3.8	3.8
< 15	3	100.0	--	--	--	--
15-17	51	96.1	3.9	--	--	--
18-19	110	70.9	25.5	3.6	--	--
(15-19)	(161)	(78.9)	(18.6)	(2.5)	(--)	(--)
20-24	381	51.7	35.4	10.2	2.1	0.5
25-29	233	37.3	30.9	21.9	4.3	5.6
30-34	91	33.0	28.6	12.1	13.2	13.2
35-39	20	30.0	5.0	20.0	20.0	25.0
> 40	4	50.0	--	--	--	50.0
DE BACA COUNTY	36	50.0	25.0	13.9	5.6	5.6
< 15	0	--	--	--	--	--
15-17	3	100.0	--	--	--	--
18-19	8	75.0	25.0	--	--	--
(15-19)	(11)	(81.8)	(18.2)	(--)	(--)	(--)
20-24	11	45.5	45.5	9.1	--	--
25-29	11	36.4	9.1	27.3	9.1	18.2
30-34	1	--	--	100.0	--	--
35-39	1	--	100.0	--	--	--
> 40	1	--	--	--	100.0	--
EDDY COUNTY	885	33.4	34.5	18.4	9.2	4.5
< 15	1	100.0	--	--	--	--
15-17	60	90.0	8.3	1.7	--	--
18-19	111	67.6	21.6	9.9	0.9	--
(15-19)	(171)	(75.4)	(17.0)	(7.0)	(0.6)	(--)
20-24	311	36.0	41.2	15.1	5.5	2.3
25-29	268	15.3	43.3	25.7	11.6	4.1
30-34	90	12.2	24.4	32.2	20.0	11.1
35-39	37	5.4	24.3	13.5	35.1	21.6
> 40	7	--	14.3	14.3	14.3	57.1
GUADALUPE COUNTY	78	37.2	28.2	17.9	9.0	7.7
< 15	0	--	--	--	--	--
15-17	7	85.7	14.3	--	--	--
18-19	12	50.0	41.7	8.3	--	--
(15-19)	(19)	(63.2)	(31.6)	(5.3)	(--)	(--)
20-24	28	46.4	25.0	21.4	7.1	--
25-29	16	6.3	31.3	37.5	18.8	6.3
30-34	9	33.3	33.3	--	22.2	11.1
35-39	5	--	--	20.0	--	80.0
> 40	1	--	100.0	--	--	--
LEA COUNTY	1367	38.3	32.6	18.1	6.1	4.8
< 15	6	100.0	--	--	--	--
15-17	96	86.5	12.5	1.0	--	--
18-19	144	69.4	24.3	6.3	--	--
(15-19)	(240)	(76.3)	(19.6)	(4.2)	(--)	(--)
20-24	494	40.3	38.3	15.8	4.3	1.4
25-29	430	25.3	36.0	23.3	9.1	6.3
30-34	152	15.1	31.6	28.3	12.5	12.5
35-39	36	8.3	19.4	36.1	8.3	27.8
> 40	9	11.1	--	33.3	22.2	33.3
LINCOLN COUNTY	246	44.3	31.7	15.9	5.7	2.4
< 15	0	--	--	--	--	--
15-17	9	77.8	22.2	--	--	--
18-19	19	94.7	5.3	--	--	--
(15-19)	(28)	(89.3)	(10.7)	(--)	(--)	(--)
20-24	96	46.9	37.5	12.5	3.1	--
25-29	72	36.1	36.1	19.4	8.3	--
30-34	36	33.3	27.8	22.2	11.1	5.6
35-39	9	11.1	33.3	33.3	--	22.2
> 40	5	--	--	40.0	20.0	40.0
QUAY COUNTY	176	43.8	29.0	15.9	5.1	6.3
< 15	0	--	--	--	--	--
15-17	17	76.5	17.6	5.9	--	--
18-19	38	60.5	26.3	10.5	--	2.6
(15-19)	(55)	(65.5)	(23.6)	(9.1)	(--)	(1.8)
20-24	54	50.0	29.6	14.8	3.7	1.9
25-29	45	31.1	37.8	15.6	8.9	6.7
30-34	16	--	18.8	37.5	12.5	31.3
35-39	5	--	20.0	40.0	20.0	20.0
> 40	1	--	100.0	--	--	--
ROOSEVELT COUNTY	286	42.0	28.7	15.7	8.7	4.9
< 15	2	100.0	--	--	--	--
15-17	16	100.0	--	--	--	--
18-19	37	64.9	27.0	5.4	2.7	--
(15-19)	(53)	(75.5)	(18.9)	(3.8)	(1.9)	(--)
ROOSEVELT COUNTY (continued)						
20-24	94	44.7	35.1	16.0	3.2	1.1
25-29	74	32.4	39.2	14.9	9.5	4.1
30-34	44	18.2	18.2	29.5	22.7	11.4
35-39	13	15.4	15.4	15.4	30.8	23.1
> 40	6	33.3	--	33.3	--	33.3
DISTRICT V						
CATRON COUNTY	34	41.2	29.4	20.6	5.9	2.9
< 15	0	--	--	--	--	--
15-17	0	--	--	--	--	--
18-19	3	66.7	33.3	--	--	--
(15-19)	(3)	(66.7)	(33.3)	(--)	(--)	(--)
20-24	11	54.5	--	36.4	9.1	--
25-29	10	40.0	50.0	10.0	--	--
30-34	5	--	60.0	40.0	--	--
35-39	3	33.3	33.3	--	33.3	--
> 40	2	50.0	--	--	--	50.0
DONA ANA COUNTY	2140	39.8	29.5	17.5	7.9	5.3
< 15	4	100.0	--	--	--	--
15-17	109	87.2	11.9	0.9	--	--
18-19	222	68.9	26.1	5.0	--	--
(15-19)	(331)	(74.9)	(21.5)	(3.6)	(--)	(--)
20-24	756	48.3	32.7	12.3	4.9	1.9
25-29	620	27.6	31.0	25.0	11.3	5.2
30-34	295	14.9	31.2	27.5	14.6	11.9
35-39	118	13.6	23.7	26.3	13.6	22.9
> 40	16	18.8	6.3	18.8	25.0	31.3
GRANT COUNTY	515	35.7	35.3	16.9	8.0	4.1
< 15	1	100.0	--	--	--	--
15-17	40	92.5	7.5	--	--	--
18-19	65	66.2	30.8	--	3.1	--
(15-19)	(105)	(76.2)	(21.9)	(--)	(1.9)	(--)
20-24	165	40.0	43.6	12.1	4.2	--
25-29	141	17.0	41.8	29.8	6.4	5.0
30-34	69	13.0	29.0	24.6	20.3	13.0
35-39	29	13.8	24.1	27.6	24.1	10.3
> 40	5	--	20.0	--	40.0	40.0
HIDALGO COUNTY	107	32.7	31.8	18.7	10.3	6.5
< 15	0	--	--	--	--	--
15-17	11	63.6	27.3	9.1	--	--
18-19	10	60.0	30.0	10.0	--	--
(15-19)	(21)	(61.9)	(28.6)	(9.5)	(--)	(--)
20-24	41	36.6	34.1	22.0	7.3	--
25-29	31	16.1	32.3	29.0	16.1	6.5
30-34	9	22.2	22.2	--	33.3	22.2
35-39	3	--	33.3	--	--	66.7
> 40	2	--	50.0	--	--	50.0
LUNA COUNTY	302	35.4	33.1	17.2	7.6	6.6
< 15	0	--	--	--	--	--
15-17	29	79.3	17.2	3.4	--	--
18-19	41	68.3	31.7	--	--	--
(15-19)	(70)	(72.9)	(25.7)	(1.4)	(--)	(--)
20-24	99	28.3	45.5	20.2	6.1	--
25-29	73	26.0	30.1	23.3	11.0	9.6
30-34	38	15.8	26.3	26.3	10.5	21.1
35-39	18	11.1	27.8	16.7	27.8	16.7
> 40	4	25.0	--	25.0	--	50.0
OTERO COUNTY	1114	41.1	36.1	15.0	4.7	3.1
< 15	2	100.0	--	--	--	--
15-17	38	86.8	13.2	--	--	--
18-19	118	65.3	34.7	--	--	--
(15-19)	(156)	(70.5)	(29.5)	(--)	(--)	(--)
20-24	475	47.6	37.7	12.2	1.9	0.6
25-29	310	30.3	39.7	19.7	7.1	3.2
30-34	130	13.8	33.8	32.3	8.5	11.5
35-39	36	19.4	27.8	13.9	25.0	13.9
> 40	5	20.0	--	20.0	20.0	40.0
SIERRA COUNTY	98	39.8	32.7	17.3	8.2	2.0
< 15	0	--	--	--	--	--
15-17	2	100.0	--	--	--	--
18-19	14	64.3	28.6	7.1	--	--
(15-19)	(16)	(68.8)	(25.0)	(6.3)	(--)	(--)
20-24	34	50.0	23.5	26.5	--	--
25-29	29	24.1	44.8	13.8	17.2	--
30-34	16	18.8	43.8	12.5	18.8	6.3
35-39	3	33.3	--	33.3	--	33.3
> 40	0	--	--	--	--	--
SOCORRO COUNTY	248	37.5	30.2	17.3	8.9	6.0
< 15	0	--	--	--	--	--
15-17	16	93.8	6.3	--	--	--
18-19	30	76.7	13.3	6.7	3.3	--
(15-19)	(46)	(82.6)	(10.9)	(4.3)	(2.2)	(--)
20-24	79	38.0	38.0	20.3	2.5	1.3
25-29	65	26.2	27.7	20.0	16.9	9.2
30-34	45	17.8	42.2	20.0	17.8	2.2
35-39	12	--	25.0	25.0	--	50.0
> 40	1	--	--	--	--	100.0

TABLE A-6
NEW MEXICO
BIRTHS BY COUNTY OF OCCURRENCE BY FACILITY AND ATTENDANT AT BIRTH
1985

COUNTY/PLACE OF BIRTH	TOTAL BIRTHS	ATTENDANT AT BIRTH					
		M.D.	D.O.	LAY MIDWIFE	NURSE MIDWIFE	R.N.	OTHER/ UNSPEC.
STATE TOTAL	27,300	23,935	1,126	286	1,732	15	206
DISTRICT I							
CIBOLA COUNTY:	265	264	0	0	0	0	1
Acoma-Canoncito-Laguna PHS Hospital	3	2					1
Cibola General Hospital	262	262					
MCKINLEY COUNTY:	2,045	2,021	0	2	14	1	7
Crownpoint Indian Hospital (PHS)	104	104					
Gallup Indian Hospital (PHS)	1,319	1,301			14	1	3
Rehoboth/Mckinley Christian Hospital	458	458					
Zuni Comprehensive Comm. Hlth Ctr (PHS)	154	153					1
Home	8	3		2			3
Clinic and Other Places	2	2					
SAN JUAN COUNTY:	2,186	1,652	0	4	521	0	9
Shiprock PHS Hospital	1,012	486			521		5
San Juan Regional Medical Center	1,161	1,160					1
Home	12	6		4			2
Clinic and Other Places	1						1
DISTRICT II							
COLFAX COUNTY:	235	232	0	0	0	0	3
Colfax General Hospital	11	11					
Northern Colfax County Hospital	224	221					3
HARDING COUNTY:	0						
LOS ALAMOS COUNTY:	415	336	0	1	78	0	0
Los Alamos Medical Center	413	336			77		
Home	2			1	1		
MORA COUNTY:	2	1	0	0	0	0	1
Home	2	1					1
RIO ARRIBA COUNTY:	385	362	6	5	2	0	10
Espanola Hospital	360	360					
Home	10			5	2		3
Clinic and Other Places	15	2	6				7
SAN MIGUEL COUNTY:	532	466	0	21	39	0	6
Northeastern Regional Hospital	505	466			39		
Home	7			2			5
Clinic and Other Places	20			19			1
SANTA FE COUNTY:	1,813	1,669	56	33	30	0	25
PHS Indian Hospital	403	343	55				5
St. Vincent Hospital	1,336	1,323	1				12
Home	64	3		29	24		8
Clinic and Other Places	10			4	6		
TAOS COUNTY:	342	282	0	54	1	0	5
Holy Cross Hospital	278	277					1
Home	23	3		17	1		2
Clinic and Other Places	41	2		37			2
UNION COUNTY:	65	64	0	0	0	0	1
Union General Hospital	64	64					
Home	1						1
DISTRICT III							
BERNALILLO COUNTY:	9,721	8,321	352	42	939	12	55
Lovelace-Bataan Medical Center	1,447	954			493		
University of New Mexico Hospital	2,772	2,732	1		33		6
St. Joseph West Mesa Hospital	573	551					22
Heights General Hospital	357	21	335		1		
Presbyterian Hospital Center	3,969	3,932	15			11	11
St. Joseph Hospital	1	1					
USAF Hospital-Kirtland AFB	466	123			342		1
Home	82	5		38	28		11
Enroute	2						2
Southwest Maternity Center (clinic)	40				40		
Clinics and Other Places	12	2	1	4	2	1	2
SANDOVAL COUNTY:	148	133	0	7	5	0	3
Cuba Hospital	131	131					
Home	14			6	5		3
Clinic and Other Places	3	2		1			
TORRANCE COUNTY:	2	0	0	2	0	0	0
Home	2			2			
DISTRICT III (continued)							
VALENCIA COUNTY:	152	126	21	4	0	0	1
Valencia Presbyterian Hospital	147	126	21				
Home	5			4			1
DISTRICT IV							
CHAVES COUNTY:	1,063	639	418	1	0	0	5
Eastern N.M. Medical Center	1,060	639	418				3
Home	2			1			1
Clinic and Other Places	1						1
CURRY COUNTY:	749	745	1	0	0	0	3
USAF Hospital-Cannon AFB	343	342	1				
Clovis High Plains Hospital	402	402					
Home	4	1					3
DE BACA COUNTY:	18	18	0	0	0	0	0
De Baca General Hospital	18	18					
EDDY COUNTY:	866	861	0	0	2	0	3
Artesia General Hospital	179	177			2		
Guadalupe Medical Center	681	681					
Home	5	2					3
Clinic and Other Places	1	1					
GUADALUPE COUNTY:	33	33	0	0	0	0	0
Guadalupe General Hospital	33	33					
LEA COUNTY:	1,272	1,252	5	0	0	0	15
Lea Regional Hospital	1,257	1,250	5				2
Home	9						9
Clinic and Other Places	6	2					4
LINCOLN COUNTY:	308	305	0	1	0	0	2
Ruidiso Hondo Valley Hospital	303	303					
Home	4	2		1			1
Clinic and Other Places	1						1
QUAY COUNTY:	150	148	0	0	0	0	2
Trigg Memorial Hospital	148	148					
Home	2						2
ROOSEVELT COUNTY:	478	256	218	0	0	0	4
Roosevelt General Hospital	474	256	218				
Home	2						2
Clinic and Other Places	2						2
DISTRICT V							
CATRON COUNTY:	2	0	0	0	0	0	2
Clinic and Other Places	2						2
DONA ANA COUNTY:	1,947	1,825	47	62	0	2	11
McAfee Hospital-WSMR	1	1					
Memorial General Hospital	1,873	1,821	47			2	3
Home	41	3		33			5
Clinic and Other Places	32			29			3
GRANT COUNTY:	565	543	1	16	1	0	4
Gila Regional Hospital	544	543	1				
Home	16			14	1		1
Clinic and Other Places	5			2			3
HIDALGO COUNTY:	0						
LUNA COUNTY:	331	328	0	0	0	0	3
Mimbres Memorial Hospital	325	325					
Home	3	1					2
Enroute	2	2					
Clinic and Other Places	1						1
OTERO COUNTY:	1,011	857	0	30	100	0	24
Gerald Champion Memorial Hospital	433	433					
Mescalero Indian Hospital (PHS)	2	2					
USAF Hospital-Holloman AFB	533	420			100		13
Home	31	2		22			7
Clinic and Other Places	12			8			4
SIERRA COUNTY:	8	7	0	1	0	0	0
Sierra Vista Hospital	7	7					
Home	1			1			
SOCORRO COUNTY:	191	189	1	0	0	0	1
Socorro General Hospital	190	189	1				
Clinic and Other Places	1						1

TABLE A-7
NEW MEXICO 1985 BIRTHS AND DEATHS BY PLACE OF RESIDENCE

	RESIDENT	
LOCATION	BIRTHS	DEATHS
DISTRICT I		
CIBOLA COUNTY:		
Grants-Urban	196	50
Grants-Rural	38	0
Milan-Urban	32	11
Milan-Rural	9	0
Other & Unspecified	250	70
COUNTY TOTAL	525	131
McKINLEY COUNTY:		
Crownpoint-All	113	24
Gallup-Urban	483	132
Gallup-Rural	25	21
Rehoboth-All	18	7
Zuni-All	181	29
Other & Unspecified	900	142
COUNTY TOTAL	1720	355
SAN JUAN COUNTY:		
Aztec-Urban	122	40
Aztec-Rural	82	16
Bloomfield-Urban	134	28
Bloomfield-Rural	85	17
Farmington-Urban	836	164
Farmington-Rural	76	25
Shiprock-All	296	50
Other & Unspecified	597	87
COUNTY TOTAL	2228	427
DISTRICT II		
COLFAX COUNTY:		
Cimarron-All	28	14
Raton-Urban	144	81
Raton-Rural	12	5
Springer-Urban	22	14
Springer-Rural	3	2
Eagle Nest-All	15	1
Other & Unspecified	32	8
COUNTY TOTAL	256	125
HARDING COUNTY:		
Roy-All	9	9
Mosquero-All	2	3
Other & Unspecified	5	3
COUNTY TOTAL	16	15
LOS ALAMOS COUNTY:		
Los Alamos-Urban	190	66
Los Alamos-Rural	12	2
White Rock-All	2	4
Other & Unspecified	0	0
COUNTY TOTAL	204	72
MORA COUNTY:		
Wagon Mound-All	8	7
Mora-All	25	8
Cleveland-All	8	2
Other & Unspecified	31	14
COUNTY TOTAL	72	31
RIO ARRIBA COUNTY:		
Espanola-Urban (Includes Fairview)	135	44
Espanola-Rural (Includes Fairview)	116	18
Chama-All	32	15
Tierra Amarilla-All	12	7
Abiquiu-All	10	8
El Rito-All	22	6
Other & Unspecified	431	170
COUNTY TOTAL	758	268
SAN MIGUEL COUNTY:		
Las Vegas-Urban	311	127
Las Vegas-Rural	52	16
Pecos-All	34	11
Other & Unspecified	70	33
COUNTY TOTAL	467	187
DISTRICT II (Continued)		
SANTA FE COUNTY:		
Santa Fe-Urban	906	387
Santa Fe-Rural	315	54
Tesuque-All	21	5
Pojoaque-All	24	4
Other & Unspecified	183	49
COUNTY TOTAL	1449	499
TAOS COUNTY:		
Questa-All	63	9
Ranchos de Taos-All	59	17
Taos-Urban	120	52
Taos-Rural	61	14
Penasco-All	16	4
Other & Unspecified	139	55
COUNTY TOTAL	458	151
UNION COUNTY:		
Clayton-Urban	57	41
Clayton-Rural	4	2
Des Moines-All	0	3
Other & Unspecified	15	13
COUNTY TOTAL	76	59
DISTRICT III		
BERNALILLO COUNTY:		
Albuquerque-Urban	7248	2852
Albuquerque-Rural	518	141
Sandia/Kirtland Airforce Base (Military	9	1
Other & Unspecified	162	44
COUNTY TOTAL	7937	3038
SANDOVAL COUNTY:		
Bernalillo-Urban	44	20
Bernalillo-Rural	68	12
Cuba-All	53	27
Pena Blanca-All	8	3
Algodones-All	10	3
Jemez Springs-All	5	8
Other & Unspecified	793	239
COUNTY TOTAL	981	312
TORRANCE COUNTY:		
Estancia-All	23	14
Moriarty-All	60	18
Mountainair-Urban	7	17
Mountainair-Rural	16	2
Encino-All	4	1
Other & Unspecified	26	16
COUNTY TOTAL	136	68
VALENCIA COUNTY:		
Belen-Urban	143	70
Belen-Rural	74	61
Los Lunas-Urban	154	59
Los Lunas-Rural	167	37
Bosque Farms	41	20
Other & Unspecified	50	19
COUNTY TOTAL	629	266
DISTRICT IV		
CHAVES COUNTY:		
Dexter-All	55	24
Hagerman-Urban	14	11
Hagerman-Rural	9	3
Roswell-Urban	842	451
Roswell-Rural	66	39
Other & Unspecified	26	6
COUNTY TOTAL	1012	534
CURRY COUNTY:		
Clovis-Urban	694	249
Clovis-Rural	44	8
Texico-All	28	5

TABLE A-7
NEW MEXICO 1985 BIRTHS AND DEATHS BY PLACE OF RESIDENCE

LOCATION	RESIDENT BIRTHS	RESIDENT DEATHS
DISTRICT IV (Continued)		
Melrose-All	4	10
Cannon Airforce Base (Military)	112	2
Other & Unspecified	11	6
COUNTY TOTAL	893	280
DE BACA COUNTY:		
Ft. Sumner-Urban	19	32
Ft. Sumner-Rural	14	9
Other & Unspecified	3	3
COUNTY TOTAL	36	44
EDDY COUNTY:		
Artesia-Urban	253	109
Artesia-Rural	52	22
Carlsbad-Urban	422	290
Carlsbad-Rural	98	23
Loving-Urban	27	11
Loving-Rural	5	2
Other & Unspecified	28	6
COUNTY TOTAL	885	463
GUADALUPE COUNTY:		
Santa Rosa-Urban	51	19
Santa Rosa-Rural	12	1
Vaughn-Urban	5	9
Vaughn-Rural	4	1
Other & Unspecified	6	6
COUNTY TOTAL	78	36
LEA COUNTY:		
Eunice-Urban	62	16
Eunice-Rural	4	2
Hobbs-Urban	794	245
Hobbs-Rural	142	19
Jal-Urban	47	15
Jal-Rural	7	2
Lovington-Urban	229	79
Lovington-Rural	35	9
Tatum-Urban	21	5
Tatum-Rural	4	2
Other & Unspecified	22	4
COUNTY TOTAL	1367	398
LINCOLN COUNTY:		
Capitan-All	23	9
Carrizozo-Urban	17	16
Carrizozo-Rural	2	2
Ruidoso-Urban	117	45
Ruidoso-Rural	12	1
Ruidoso Downs-All	39	13
Other & Unspecified	36	18
COUNTY TOTAL	246	104
QUAY COUNTY:		
Tucumcari-Urban	124	87
Tucumcari-Rural	21	13
San Jon-All	8	1
Logan-All	14	8
Other & Unspecified	9	6
COUNTY TOTAL	176	115
ROOSEVELT COUNTY:		
Portales-Urban	197	92
Portales-Rural	66	13
Other & Unspecified	23	10
COUNTY TOTAL	286	115

LOCATION	RESIDENT BIRTHS	RESIDENT DEATHS
DISTRICT V		
CATRON COUNTY:		
Reserve-All	14	11
Other & Unspecified	20	15
COUNTY TOTAL	34	26
DONA ANA COUNTY:		
Hatch-All	56	25
Las Cruces-Urban	1232	360
Las Cruces-Rural	49	46
Mesilla-Urban	2	8
Mesilla-Rural	54	1
University Park-All	0	0
White Sands-All	69	6
La Mesa-All	56	16
Other & Unspecified	622	123
COUNTY TOTAL	2140	585
GRANT COUNTY:		
Bayard-Urban	55	19
Bayard-Rural	2	1
Santa Rita-Urban	0	0
Santa Rita-Rural	0	0
Central-Urban	42	17
Central-Rural	2	2
Hurley-Urban	21	14
Hurley-Rural	4	1
Silver City-Urban	238	90
Silver City-Rural	52	17
Fort Bayard-All	6	1
Other & Unspecified	93	25
COUNTY TOTAL	515	187
HIDALGO COUNTY:		
Lordsburg-Urban	63	35
Lordsburg-Rural	8	0
Other & Unspecified	36	7
COUNTY TOTAL	107	42
LUNA COUNTY:		
Deming-Urban	190	157
Deming-Rural	91	43
Other Unspecified	21	15
COUNTY TOTAL	302	215
OTERO COUNTY:		
Alamogordo-Urban	634	176
Alamogordo-Rural	65	15
Mescalero-All	71	15
Tularosa-Urban	51	18
Tularosa-Rural	19	7
Cloudcroft-All	12	7
Holloman AFB (Military)	181	4
Other & Unspecified	81	25
COUNTY TOTAL	1114	267
SIERRA COUNTY:		
Truth or Consequences-Urban	19	112
Truth or Consequences-Rural	50	6
Other & Unspecified	29	41
COUNTY TOTAL	98	159
SOCORRO COUNTY:		
Magdalena-Urban	11	5
Magdalena-Rural	41	3
Socorro-Urban	144	36
Socorro-Rural	26	5
Other & Unspecified	26	14
COUNTY TOTAL	248	63

TABLE A-8
AGE-SPECIFIC DEATH RATES BY COUNTY AND SEX NEW MEXICO RESIDENTS, 1983-85 AVERAGE UNITED STATES 1984
(Rates per 100,000 Population)

		AGE GROUP										
HEALTH DISTRICTS	COUNTY	< 1	1 - 4	5 - 14	15 - 24	25 - 34	35 - 44	45 - 54	55 - 64	65 - 74	75 - 84	=>85
	MALE											
	UNITED STATES	1198.2	57.0	32.1	141.6	173.9	270.2	676.6	1705.2	3813.0	8445.9	18119.1
	NEW MEXICO	1191.0	73.8	41.1	216.9	252.8	265.7	616.0	1378.4	3095.8	7134.0	15117.3
	CIBOLA	749.1	147.1	34.4	619.2	278.7	262.1	476.2	929.1	2850.1	4801.1	6666.7
1	MCKINLEY	1382.5	168.7	35.7	463.7	500.2	624.5	828.2	1552.8	3458.4	8736.2	19090.9
	SAN JUAN	704.2	110.8	34.8	294.9	375.2	335.9	571.9	1274.0	3003.4	8455.3	12476.7
	COLFAX	769.2	59.5	56.4	193.1	304.3	338.2	675.7	1858.1	2907.4	6577.0	15068.5
	HARDING	4166.7	--	--	483.1	--	--	--	--	1025.6	10416.7	25925.9
	LOS ALAMOS	1634.0	--	51.6	93.1	33.5	98.6	287.2	753.8	2079.1	8333.3	20289.9
	MORA	3703.7	183.2	86.6	76.3	277.8	546.4	683.8	1185.2	2394.1	4834.6	15384.6
2	RIO ARRIBA	1666.7	44.5	56.2	495.6	459.0	404.9	804.8	1659.6	4148.6	8290.2	14814.8
	SAN MIGUEL	706.2	64.8	16.3	163.9	360.8	342.8	800.0	1371.0	2313.6	7827.6	21183.8
	SANTA FE	651.8	48.1	31.4	272.4	199.8	203.6	560.4	1186.6	2402.7	5825.2	13459.1
	TAOS	757.6	106.0	68.8	275.2	472.8	384.5	657.3	1247.3	2749.1	6998.8	14953.3
	UNION	833.3	--	81.3	442.8	105.8	273.2	136.1	2279.2	3549.4	7407.4	15686.3
	BERNALILLO	1431.7	58.7	36.0	159.7	210.9	224.4	604.7	1392.9	3213.9	7969.2	17088.4
3	SANDOVAL	1166.0	97.5	41.8	232.0	345.3	381.8	597.7	1206.4	2401.8	5563.0	12473.1
	TORRANCE	1687.8	125.3	123.0	204.5	490.2	420.4	647.2	1649.3	3554.3	5416.7	8771.9
	VALENCIA	1592.4	92.6	86.2	290.8	284.3	337.9	703.2	1639.0	2995.6	7336.2	9713.0
	CHAVES	1234.6	77.0	43.4	234.4	251.8	246.9	718.8	1714.7	2936.9	7056.0	16181.2
	CURRY	958.2	71.6	45.3	151.8	158.2	274.1	775.2	1468.5	3289.8	6178.4	11711.7
	DE BACA	1388.9	--	--	376.6	1139.6	476.2	289.9	1960.8	4325.7	8333.3	20833.3
	EDDY	1223.9	45.2	43.9	250.2	281.8	336.5	641.0	1418.8	3468.0	7588.4	24342.1
4	GUADALUPE	3252.0	199.6	85.5	144.9	476.2	436.7	917.4	578.0	1766.0	8558.6	12121.2
	LEA	1350.7	28.0	32.0	209.4	246.7	300.9	698.8	1746.9	3501.5	7250.6	15123.5
	LINCOLN	555.6	61.5	64.1	96.2	469.9	224.1	906.3	837.7	2824.0	11111.1	28282.8
	QUAY	757.6	421.9	66.1	330.8	110.1	41.4	715.1	1320.1	3256.1	10040.2	11111.1
	ROOSEVELT	1207.7	100.0	60.2	154.7	124.2	207.3	496.7	1655.6	3200.9	8843.5	20467.8
	CATRON	2222.2	--	--	--	660.1	196.1	447.4	1041.7	4201.7	5714.3	37500.0
	DONA ANA	741.4	45.1	28.8	114.8	169.4	148.7	591.5	1158.3	2807.3	4932.8	12214.6
	GRANT	1371.7	28.7	41.5	271.7	350.7	249.1	415.3	1207.7	3320.9	6582.3	9968.8
5	HIDALGO	2525.3	--	58.6	66.4	335.9	194.4	326.8	995.0	4590.8	6425.7	22222.2
	LUNA	427.4	63.5	26.4	293.7	190.5	228.6	771.2	1886.1	3617.1	6166.4	11838.0
	OTERO	1085.0	77.4	40.6	184.0	183.3	207.9	607.6	1369.3	3387.8	7407.4	11805.6
	SIERRA	4761.9	--	119.5	189.4	274.3	244.2	707.7	1515.2	2798.7	5186.7	14062.5
	SOCORRO	1388.9	71.4	85.2	170.3	170.4	303.0	227.1	1158.8	2721.1	6578.9	19697.0

TABLE A-8
AGE-SPECIFIC DEATH RATES BY COUNTY AND SEX
NEW MEXICO RESIDENTS, 1983-85 AVERAGE
UNITED STATES 1984

(Rates per 100,000 Population)

AGE GROUP

HEALTH DISTRICTS

COUNTY	< 1	1 - 4	5 - 14	15 - 24	25 - 34	35 - 44	45 - 54	55 - 64	65 - 74	75 - 84	=>85
FEMALE											
UNITED STATES	967.5	46.5	21.1	51.1	68.5	141.5	374.6	918.4	2096.4	5188.2	14053.9
NEW MEXICO	815.6	54.2	26.1	62.8	82.3	143.2	321.6	778.0	1738.6	4221.3	12962.7
CIBOLA	890.6	58.6	--	144.5	81.5	165.1	309.6	557.1	1548.0	5031.4	16666.7
1 MCKINLEY	927.0	85.7	10.0	90.8	153.3	232.8	382.1	920.3	2017.5	4090.4	16142.6
SAN JUAN	850.0	164.6	30.9	74.6	113.4	164.9	344.5	783.8	1695.5	3383.5	9128.2
COLFAX	961.5	--	--	84.5	33.8	75.1	112.6	483.1	1801.8	4199.5	12202.4
HARDING	--	--	--	--	--	--	--	966.2	1360.5	8974.4	20512.8
LOS ALAMOS	666.7	--	25.8	--	33.5	125.7	128.9	609.4	1007.6	3973.5	11805.6
MORA	877.2	--	103.5	--	120.8	150.2	577.2	307.2	1951.2	1424.5	14285.7
2 RIO ARRIBA	913.2	101.5	--	112.5	181.7	225.0	269.9	675.1	2067.5	5694.4	17994.1
SAN MIGUEL	544.2	--	34.3	25.4	102.8	171.3	280.4	651.6	1981.0	3598.1	11754.4
SANTA FE	464.3	46.5	16.2	93.5	94.9	134.8	342.9	724.8	1875.7	3870.8	10377.4
TAOS	1304.3	39.9	57.3	81.0	235.4	193.4	137.5	420.5	1573.3	4395.6	15000.0
UNION	833.3	--	--	--	--	--	283.7	1228.9	1260.5	5063.3	16666.7
BERNALILLO	796.2	34.3	19.5	51.6	62.8	137.0	311.8	782.4	1683.4	4223.7	13470.9
3 SANDOVAL	660.8	60.9	38.2	18.6	117.9	111.4	413.7	617.9	1473.2	3038.0	12500.0
TORRANCE	584.8	--	45.8	52.4	110.9	63.4	253.8	688.1	1563.8	2824.9	12500.0
VALENCIA	843.9	69.6	33.4	150.8	74.6	88.1	285.1	1034.3	2019.4	3982.3	9750.6
CHAVES	1142.5	60.4	56.8	88.7	99.9	197.5	380.1	940.9	1730.4	4619.8	10954.3
CURRY	740.7	59.0	9.9	52.6	61.2	72.5	479.8	779.0	1468.5	3610.1	12111.8
DE BACA	--	--	--	--	252.5	252.5	--	--	3268.0	5128.2	14285.7
EDDY	976.0	93.9	15.3	98.5	62.0	84.1	307.3	922.1	1569.9	4690.6	16570.9
4 GUADALUPE	1801.8	--	--	81.9	223.7	408.2	--	673.4	1960.8	5555.6	17543.9
LEA	1035.5	20.9	31.5	63.1	95.2	154.6	468.5	1033.9	2055.3	5399.1	12131.5
LINCOLN	1322.8	192.1	156.7	35.2	192.3	160.3	220.5	617.3	1936.6	4186.0	13043.5
QUAY	1087.0	--	41.3	110.0	188.9	330.0	206.8	662.3	2200.2	4793.0	12714.8
ROOSEVELT	719.4	59.0	60.2	82.8	88.4	73.6	331.4	1041.7	1036.5	3580.6	16071.4
CATRON	3921.6	--	465.1	273.2	--	181.2	--	490.2	1881.7	4615.4	12121.2
DONA ANA	613.3	55.9	16.6	29.4	44.0	114.0	312.6	634.2	1913.6	3946.1	13636.4
GRANT	400.0	--	27.7	55.4	49.8	87.4	359.9	842.9	1494.0	4342.3	13075.5
5 HIDALGO	--	--	--	60.6	--	76.1	335.6	669.3	2873.6	4427.1	12766.0
LUNA	709.2	--	131.9	57.2	34.3	152.4	304.8	748.4	1476.8	3885.9	11868.7
OTERO	975.0	--	35.0	63.4	55.4	193.2	202.5	923.9	1830.3	5178.0	13224.6
SIERRA	1626.0	--	--	--	191.0	371.2	908.5	944.2	1569.0	3891.1	12202.4
SOCORRO	268.8	--	26.2	28.4	62.0	97.3	170.4	624.6	1870.7	4992.7	13131.3

TABLE A-9
NEW MEXICO RESIDENT DEATHS
NUMBER AND CRUDE DEATH RATE BY CAUSE
1985

CAUSE OF DEATH	9TH REVISION ICDA	TOTAL	RATE	MALE	RATE	FEMALE	RATE
ALL CAUSES		9637	664.6	5477	767.9	4160	564.7
TUBERCULOSIS (ALL FORMS)	010-018	14	1.0	10	1.4	4	0.5
SYPHILIS AND ITS SEQUELAE	090-097	-	0.0	-	0.0	-	0.0
ALL OTHER INFECTIOUS AND PARASITIC DISEASES	001-009,020-088,098-139	73	5.0	38	5.3	35	4.8
MALIGNANT NEOPLASMS	140-208	1921	132.5	1029	144.3	892	121.1
Buccal Cavity and Pharynx	140-149	21	1.4	15	2.1	6	0.8
Digestive Organs and Peritoneum	150-159	493	34.0	250	35.0	243	33.0
Respiratory System	160-165	477	32.9	318	44.6	159	21.6
Breast	174-175	176	12.1	2	0.3	174	23.6
Genital Organs	179-187	207	14.3	134	18.8	73	9.9
Urinary Organs	188,189	69	4.8	46	6.4	23	3.1
Other and Unspecified Sites	170-173,[redacted]190-199	274	18.9	155	21.7	119	16.2
Lymphatic and Hematopietic Tissues	200-208	204	14.1	109	15.3	95	12.9
DIABETES	250	251	17.3	108	15.1	143	19.4
MAJOR CARDIOVASCULAR DISEASE	390-459	3507	241.9	1891	265.1	1616	219.4
Diseases of the Heart	390-398,402,404-429	2700	186.2	1542	216.2	1158	157.2
Active Rheumatic Fever	390-392	1	0.1	0	0.0	1	0.1
Chronic Rheumatic Fever	393-398	28	1.9	11	1.5	17	2.3
Hypertension with Heart Disease	402,404	79	5.4	30	4.2	49	6.7
Acute Myocardial Infarction	410	860	59.3	532	74.6	328	44.5
Other Ischemic Heart Diseases	411-414	692	47.7	357	50.0	335	45.5
Diseases of Pulmonary Circulation	415-417	58	4.0	33	4.6	25	3.4
Other Diseases of the Heart	420-429	982	67.7	579	81.2	403	54.7
Hypertension with or without Renal Disease	401,403	32	2.2	12	1.7	20	2.7
Cerebrovascular Disease	430-438	575	39.7	234	32.8	341	46.3
Atherosclerosis	440	88	6.1	35	4.9	53	7.2
Other Diseases of the Circulatory System	441-459	112	7.7	68	9.5	44	6.0
PNEUMONIA	480-486	266	18.3	134	18.8	132	17.9
INFLUENZA	487	3	0.2	1	0.1	2	0.3
CHRONIC OBSTRUCTIVE PULMONARY DISEASES (COPD) AND ALLIED CONDITIONS	490-496	483	33.3	301	42.2	182	24.7
Bronchitis, Chronic and Unspecified	490-491	12	0.8	4	0.6	8	1.1
Emphysema	492	119	8.2	81	11.4	38	5.2
Asthma	493	26	1.8	12	1.7	14	1.9
Other COPD and Allied Conditions	494-496	326	22.5	204	28.6	122	16.6
PEPTIC ULCER	531-533	37	2.6	19	2.7	18	2.4
APPENDICITIS	540-543	1	0.1	1	0.1	0	0.0
HERNIA AND INTESTINAL OBSTRUCTION	550-553,560	26	1.8	8	1.1	18	2.4
CHRONIC LIVER DISEASE AND CIRRHOSIS	571	198	13.7	135	18.9	63	8.6
NEPHRITIS, NEPHROTIC SYNDROME, AND NEPHROSIS	580-589	106	7.3	61	8.6	45	6.1
COMPLICATIONS OF PREGNANCY, CHILDBIRTH, AND PUERPERIUM	630-676	3	0.2	0	0.0	3	0.4
CONGENITAL ANOMALIES	740-759	112	7.7	62	8.7	50	6.8
CERTAIN CONDITIONS ORIGINATING IN THE PERINATAL PERIOD	760-779	122	8.4	74	10.4	48	6.5
SYMPTOMS, SIGNS, AND ILL-DEFINED CONDITIONS	780-799	205	14.1	117	16.4	88	11.9
ALL OTHER DISEASES	Residual	975	67.2	526	73.7	449	60.9
ACCIDENTS AND ADVERSE EFFECTS	E800-E949	861	59.4	615	86.2	246	33.4
Motor Vehicle Accidents	E810-E825	490	33.8	358	50.2	132	17.9
All other Accidents and Adverse Effects	E800-E807,E826-E949	371	25.6	257	36.0	114	15.5
SUICIDE	E950-E959	257	17.7	194	27.2	63	8.6
HOMICIDE AND LEGAL INTERVENTION	E960-E978	160	11.0	118	16.5	42	5.7
ALL OTHER EXTERNAL CAUSES	E980-E999	56	3.9	35	4.9	21	2.9

TABLE A-10
NEW MEXICO RESIDENTS
SELECTED CAUSES OF DEATH BY COUNTY
1985

	COUNTY	MALIGNANT NEOPLASMS (140-208)	STOMACH (151)	LARGE INTESTINE (153)	RECTUM (154)	LIVER & BILE DUCTS (155)	TRACHEA, BRONCHUS LUNG (162-163)	BLADDER (188)	DIABETES MELLITUS (250)	HYPERTENSIVE DISEASE (401-405)	ISCHEMIC HEART DISEASE (410-414)	CEREBRO-VASCULAR (430-438)	INFLUENZA & PNEUMONIA (480-487)	CHRONIC OBSTRUCTIVE PULMONARY DISEASE (490-496)	CIRRHOSIS OF LIVER (571)	ACCIDENTS (E800-E949)	MOTOR VEHICLE (E810-E825)	ACCIDENTS AT WORK	ACCIDENTS AT HOME	PUBLIC NON-MOTOR VEHICLE	SUICIDE (E950-E959)	HOMICIDE (E960-E978)
	NEW MEXICO	1,921	71	172	38	39	450	28	251	111	1,552	575	269	483	198	861	490	15	158	198	257	160
	Cibola	26	1	--	1	1	8	--	4	--	17	8	1	4	5	16	7	--	2	7	3	8
1	Mckinley	49	7	3	--	--	5	1	13	2	31	7	11	7	16	74	48	--	8	18	12	10
	San Juan	88	4	9	1	1	21	1	14	3	41	16	8	21	6	78	44	3	8	23	17	10
	Colfax	21	1	1	--	--	5	--	2	3	33	5	3	8	2	11	4	1	2	4	4	2
	Harding	4	--	2	--	--	--	--	--	--	2	--	--	--	--	1	--	--	--	1	--	--
	Los Alamos	14	--	1	--	--	6	--	3	--	8	6	1	4	1	5	1	--	3	1	3	--
	Mora	8	1	--	--	--	2	--	--	2	2	--	--	1	--	1	--	--	--	1	--	--
2	Rio Arriba	41	2	1	1	--	6	1	6	3	31	13	4	9	11	48	35	--	6	7	8	7
	San Miguel	26	2	1	2	2	2	1	6	3	30	10	5	12	7	13	8	--	3	2	7	5
H	Santa Fe	104	3	8	4	1	22	--	10	4	65	34	16	20	13	48	30	--	7	11	20	7
E	Taos	22	--	4	1	--	4	--	4	2	15	8	8	4	4	17	13	1	1	2	4	1
A	Union	14	--	2	--	--	5	--	--	--	17	5	1	3	1	1	--	--	--	1	2	1
L																						
T	Bernalillo	629	20	59	14	10	137	5	72	39	480	185	89	170	65	215	104	1	49	61	88	55
H 3	Sandoval	71	3	5	1	4	7	2	9	--	45	14	6	8	6	38	26	--	8	4	9	4
	Torrance	13	--	--	--	--	4	1	2	--	12	2	--	4	2	3	1	--	--	2	1	4
D	Valencia	62	2	2	1	--	21	1	15	4	34	11	6	12	4	27	19	--	4	4	8	3
I																						
S	Chaves	107	4	8	2	2	26	2	10	10	103	41	15	29	8	32	21	1	6	4	6	9
T	Curry	47	1	6	--	3	9	1	3	5	61	26	10	10	2	18	11	--	4	3	3	4
R	De Baca	11	--	3	--	--	4	--	--	1	8	1	--	3	3	5	1	--	2	2	2	--
I	Eddy	122	4	17	4	2	31	2	12	5	84	28	20	25	6	33	18	2	9	4	8	3
C 4	Guadalupe	3	--	--	--	--	2	--	--	1	6	1	4	1	1	--	--	--	--	--	--	2
T	Lea	75	3	6	--	5	22	3	10	4	88	30	11	28	6	28	14	2	4	8	9	5
S	Lincoln	19	--	2	--	--	7	1	2	--	20	7	1	8	3	9	4	--	2	3	1	--
	Quay	23	2	3	1	--	4	1	3	--	17	5	3	6	--	8	4	1	3	--	1	1
	Roosevelt	14	--	3	--	1	2	--	2	2	32	7	2	5	1	15	10	--	3	2	--	--
	Catron	6	--	--	1	--	2	--	--	--	4	5	--	--	--	1	--	--	--	1	1	--
	Dona Ana	128	5	10	2	5	35	1	16	7	90	33	15	23	13	49	28	1	9	11	15	14
	Grant	32	2	3	1	--	11	1	5	3	34	12	8	13	2	11	5	1	3	2	6	--
5	Hidalgo	10	--	1	--	--	4	--	3	--	7	5	1	2	--	3	2	--	1	--	--	1
	Luna	41	--	4	--	1	13	2	17	--	33	19	8	17	2	7	5	--	2	--	7	--
	Otero	52	2	5	1	1	12	--	8	2	51	17	5	13	4	30	16	1	6	7	7	2
	Sierra	24	1	1	--	--	8	--	--	5	47	9	5	8	4	9	6	--	2	1	4	1
	Socorro	15	1	2	--	--	3	1	--	1	4	5	2	5	--	7	5	--	1	1	1	--

TABLE A-11
NEW MEXICO RESIDENT
SELECTED CAUSES OF DEATH BY COUNTY AMONG MALES
1985

	COUNTY	MALIGNANT NEOPLASMS (140-208)	STOMACH (151)	LARGE INTESTINE (153)	RECTUM (154)	LIVER & BILE DUCTS (155)	TRACHEA, BRONCHUS LUNG (162-163)	PROSTATE (185)	BLADDER (188)	DIABETES MELLITUS (250)	HYPERTENSIVE DISEASE (401-405)	ISCHEMIC HEART DISEASE (410-414)	CEREBRO-VASCULAR (430-438)	INFLUENZA & PNEUMONIA (480-487)	CHRONIC OBSTRUCTIVE PULMONARY DISEASE (490-496)	CIRRHOSIS OF LIVER (571)	ACCIDENTS (E800-E949)	MOTOR VEHICLE (E810-E825)	ACCIDENTS AT WORK	ACCIDENTS AT HOME	PUBLIC NON-MOTOR VEHICLE	SUICIDE (E950-E959)	HOMICIDE (E960-E978)
	NEW MEXICO	1,029	44	81	20	17	298	133	20	108	42	889	234	135	301	135	615	358	15	97	145	194	118
	Cibola	15	1	--	1	--	6	3	--	4	--	9	2	1	3	3	9	5	--	1	3	3	4
	1 McKinley	30	4	1	--	--	5	6	--	5	2	19	2	8	3	10	59	37	--	6	16	11	9
	San Juan	46	4	4	1	--	17	7	1	7	1	19	7	5	16	5	58	32	3	4	19	9	7
	Colfax	12	1	--	--	--	3	1	--	1	1	21	1	1	5	2	7	3	1	1	2	3	2
	Harding	3	--	2	--	--	--	--	--	--	--	1	--	--	--	--	--	--	--	--	--	--	--
	Los Alamos	12	--	1	--	--	5	1	--	1	--	8	3	1	2	1	2	1	--	--	1	3	--
	Mora	2	--	--	--	--	1	--	--	--	--	1	--	--	1	--	1	--	--	--	1	--	--
	2 Rio Arriba	21	1	--	--	--	4	4	1	3	2	19	7	2	5	6	35	23	--	5	7	5	7
	San Miguel	11	1	--	1	1	2	3	1	4	2	18	7	4	6	5	9	6	--	2	1	7	2
H	Santa Fe	49	2	5	2	1	11	6	--	2	1	39	8	10	10	7	32	19	--	4	9	18	5
E	Taos	13	--	3	1	--	1	1	--	3	2	10	2	3	3	4	11	9	1	--	1	3	1
A	Union	9	--	1	--	--	3	1	--	--	--	10	2	1	2	1	--	--	--	--	--	2	1
L																							
T	Bernalillo	326	14	28	10	7	88	43	4	27	14	268	83	44	100	43	154	82	1	31	40	58	40
H	3 Sandoval	35	1	3	--	1	3	3	2	3	--	34	9	5	6	4	28	19	--	6	3	6	3
	Torrance	10	--	--	--	--	3	1	1	1	--	7	1	--	4	1	2	1	--	--	1	1	4
D	Valencia	40	2	1	1	--	13	3	1	7	3	22	7	2	6	1	21	14	--	3	4	7	2
I																							
S	Chaves	58	3	3	--	--	16	11	1	4	5	59	18	9	19	7	19	11	1	4	3	6	7
T	Curry	23	1	2	--	2	6	5	1	--	--	29	6	4	6	2	12	9	--	--	3	1	3
R	De Baca	6	--	1	--	--	2	--	--	--	--	4	1	--	3	2	4	1	--	2	1	1	--
I	Eddy	72	1	6	1	2	22	7	2	6	2	50	11	7	13	3	25	12	2	7	4	7	2
C	4 Guadalupe	2	--	--	--	--	2	--	--	--	1	4	1	1	--	1	--	--	--	--	--	--	2
T	Lea	42	1	3	--	1	19	1	3	5	2	43	10	4	14	5	19	10	2	2	5	7	4
S	Lincoln	10	--	2	--	--	5	--	1	2	--	13	5	1	4	3	6	4	--	--	2	1	--
	Quay	10	1	1	--	--	3	2	--	1	--	12	1	2	4	--	5	3	1	1	--	1	--
	Roosevelt	8	--	1	--	--	2	1	--	2	--	18	2	1	5	1	7	5	--	1	1	--	--
	Catron	4	--	--	--	--	1	--	--	--	--	4	3	--	--	--	1	--	--	--	1	--	--
	Dona Ana	61	3	5	1	1	21	11	--	6	1	48	11	5	18	11	36	22	1	5	8	14	11
	Grant	14	1	1	--	--	4	2	--	3	--	22	3	3	8	1	9	4	1	2	1	5	--
	5 Hidalgo	4	--	--	--	--	2	--	--	--	--	2	1	1	1	--	3	2	--	1	--	--	--
	Luna	28	--	3	--	1	12	4	1	7	--	17	6	4	15	2	6	4	--	2	--	5	--
	Otero	32	1	3	1	--	9	4	--	4	1	28	7	2	9	3	23	12	1	4	6	7	1
	Sierra	15	1	--	--	--	6	1	--	--	1	28	4	2	6	1	6	4	--	1	1	2	1
	Socorro	6	--	1	--	--	1	1	--	--	1	3	3	2	4	--	6	4	--	1	1	1	--

TABLE A-12
NEW MEXICO RESIDENTS
SELECTED CAUSES OF DEATH BY COUNTY AMONG FEMALES
1985

	COUNTY	MALIGNANT NEOPLASMS (140-208)	STOMACH (151)	LARGE INTESTINE (153)	RECTUM (154)	LIVER & BILE DUCTS (155)	TRACHEA, BRONCHUS LUNG (162-163)	BREAST (174)	CERVIX UTERI (180)	BLADDER (188)	DIABETES MELLITUS (250)	HYPERTENSIVE DISEASE (401-405)	ISCHEMIC HEART DISEASE (410-414)	CEREBRO-VASCULAR (430-438)	INFLUENZA PNEUMONIA (480-487)	CHRONIC OBSTRUCTIVE PULMONARY DISEASE (490-496)	CIRRHOSIS OF LIVER (571)	ACCIDENTS (E800-E949	MOTOR VEHICLE (E810-E825)	ACCIDENTS AT WORK	ACCIDENTS AT HOME	PUBLIC NON-MOTOR VEHICLE	SUICIDE (E950-E959)	HOMICIDE (E960-E978)
	NEW MEXICO	892	27	91	18	22	152	174	17	8	143	69	663	341	134	182	63	246	132	0	61	53	63	42
	Cibola	11	--	--	--	1	2	1	--	--	--	--	8	6	--	1	2	7	2	--	1	4	--	4
1	McKinley	19	3	2	--	--	--	3	1	1	8	--	12	5	3	4	6	15	11	--	2	2	1	1
	San Juan	42	--	5	--	1	4	9	--	--	7	2	22	9	3	5	1	20	12	--	4	4	8	3
	Colfax	9	--	1	--	--	2	4	--	--	1	2	12	4	2	3	--	4	1	--	1	2	1	--
	Harding	1	--	--	--	--	--	--	--	--	--	--	1	--	--	--	--	1	--	--	--	1	--	--
	Los Alamos	2	--	--	--	--	1	1	--	--	2	--	--	3	--	2	--	3	--	--	3	--	--	--
	Mora	6	1	--	--	--	1	--	1	--	--	2	1	--	--	--	--	--	--	--	--	--	--	--
2	Rio Arriba	20	1	1	1	--	2	3	--	--	3	1	12	6	2	4	5	13	12	--	1	--	3	--
	San Miguel	15	1	1	1	1	--	3	--	--	2	1	12	3	1	6	2	4	2	--	1	1	--	3
H	Santa Fe	55	1	3	2	--	11	9	1	--	8	3	26	26	6	10	6	16	11	--	3	2	2	2
E	Taos	9	--	1	--	--	3	3	--	--	1	--	5	6	5	1	--	6	4	--	1	1	1	--
A	Union	5	--	1	--	--	2	--	--	--	--	--	7	3	--	1	--	1	--	--	--	1	--	--
L																								
T	Bernalillo	303	6	31	4	3	49	69	8	1	45	25	212	102	45	70	22	61	22	--	18	21	30	15
H 3	Sandoval	36	2	2	1	3	4	8	1	--	6	--	11	5	1	2	2	10	7	--	2	1	3	1
	Torrance	3	--	--	--	--	1	--	--	--	1	--	5	1	--	--	1	1	--	--	--	1	--	--
D	Valencia	22	--	1	--	--	8	2	--	--	8	1	12	4	4	6	3	6	5	--	1	--	1	1
I																								
S	Chaves	49	1	5	2	2	10	10	1	1	6	5	44	23	6	10	1	13	10	--	2	1	--	2
T	Curry	24	--	4	--	1	3	6	1	--	3	5	32	20	6	4	--	6	2	--	4	--	2	1
R	De Baca	5	--	2	--	--	2	--	--	--	--	1	4	--	--	--	1	1	--	--	--	1	1	--
I	Eddy	50	3	11	3	--	9	7	--	--	6	3	34	17	13	12	3	8	6	--	2	--	1	1
C 4	Guadalupe	1	--	--	--	--	--	--	--	--	--	--	2	--	3	1	--	--	--	--	--	--	--	--
T	Lea	33	2	3	--	4	3	6	1	--	5	2	45	20	7	14	1	9	4	--	2	3	2	1
S	Lincoln	9	--	--	--	--	2	--	--	--	--	--	7	2	--	4	--	3	--	--	2	1	--	--
	Quay	13	1	2	1	--	1	2	--	1	2	--	5	4	1	2	--	3	1	--	2	--	--	1
	Roosevelt	6	--	2	--	1	--	2	--	--	--	2	14	5	1	--	--	8	5	--	2	1	--	--
	Catron	2	--	--	1	--	1	--	--	--	--	--	--	2	--	--	--	--	--	--	--	--	1	--
	Dona Ana	67	2	5	1	4	14	11	2	1	10	6	42	22	10	5	2	13	6	--	4	3	1	3
	Grant	18	1	2	1	--	7	3	--	1	2	3	12	9	5	5	1	2	1	--	--	1	1	--
5	Hidalgo	6	--	1	--	--	2	--	--	--	3	--	5	4	--	1	--	--	--	--	--	--	--	1
	Luna	13	--	1	--	--	1	3	--	1	10	--	16	13	4	2	--	1	1	--	--	--	2	--
	Otero	20	1	2	--	1	3	8	--	--	4	1	23	10	3	4	1	7	4	--	2	1	--	2
	Sierra	9	--	1	--	--	2	1	--	--	--	4	19	5	3	2	3	3	2	--	1	--	1	--
	Socorro	9	1	1	--	--	2	--	--	1	--	--	1	2	--	1	--	1	1	1	--	--	--	--

TABLE A-13
SUMMARY OF TRENDS IN NEW MEXICO HEALTH STATISTICS
NEW MEXICO RESIDENTS
1950-1985

		BIRTHS		DEATHS		MATERNAL DEATHS		INFANT DEATHS				FETAL DEATHS	
								ALL INFANTS		NEONATAL	POST		
YEAR	POPULATION ESTIMATE MID-YEAR	NUMBER	RATE	NUMBER	RATE	NUMBER	RATE	NUMBER	RATE	RATE	NEONATAL RATE	NUMBER	RATE
1985	1,450,000	27,449	18.9	9,637	6.6	3	10.9	291	10.6	6.2	4.4	157	5.7
1984	1,426,000	27,350	19.2	9,504	6.7	5	18.3	263	9.6	6.0	3.6	186	6.8
1983	1,399,000	27,508	19.7	9,138	6.5	3	10.9	274	10.0	5.5	4.5	192	7.0
1982	1,367,000	27,630	20.2	9,186	6.7	3	10.9	316	11.4	6.7	4.7	202	7.3
1981	1,334,000	26,565	19.9	8,668	6.5	2	7.5	256	9.6	5.7	3.9	180	6.8
1980	1,302,894*	26,589	20.4	9,032	6.9	3	11.3	293	11.0	7.0	4.0	204	7.7
1979	1,283,000	24,821	19.3	8,617	6.7	10	40.3	352	14.2	8.8	5.4	206	8.4
1978	1,254,000	23,907	19.1	8,331	6.6	2	8.4	330	13.8	8.7	5.1	199	8.3
1977	1,227,000	23,100	18.8	8,223	6.7	2	8.7	333	14.4	9.8	4.6	201	8.7
1976	1,196,000	22,180	18.5	8,204	6.9	3	13.5	347	15.6	10.0	5.6	196	8.8
1975	1,164,000	21,078	18.1	8,003	6.9	2	9.5	357	16.9	11.0	6.0	169	8.0
1974	1,130,000	21,339	18.9	8,029	7.1	10	46.8	391	18.3	12.2	6.1	184	8.6
1973	1,104,000	20,852	18.9	8,139	7.4	2	9.6	421	20.2	12.9	7.3	211	10.1
1972	1,078,000	20,813	19.3	7,877	7.3	5	24.0	403	19.4	14.4	5.0	248	11.9
1971	1,053,000	22,205	21.1	7,638	7.3	5	22.5	460	20.7	15.0	5.8	254	11.4
1970	1,017,055*	22,004	21.6	7,411	7.3	6	27.3	463	21.0	15.0	6.0	245	11.1
1969	1,011,000	21,543	21.3	7,180	7.1	5	23.2	508	23.6	16.1	7.5	266	12.3
1968	994,000	20,346	20.5	7,128	7.2	7	34.4	487	23.9	16.8	7.2	264	13.0
1967	1,000,000	21,243	21.2	6,897	6.9	4	18.8	527	24.8	16.1	8.7	249	11.7
1966	1,007,000	22,363	22.2	6,971	6.9	9	40.2	601	26.9	16.7	10.2	278	12.4
1965	1,012,000	24,352	24.1	6,801	6.7	10	41.1	657	27.0	17.8	9.2	326	13.4
1964	1,006,000	26,862	26.7	6,902	6.9	14	52.1	781	29.1	19.2	9.9	349	13.0
1963	989,000	27,820	28.1	6,837	6.9	10	35.9	848	30.5	20.0	10.5	371	13.3
1962	979,000	29,226	29.9	6,507	6.6	16	54.7	877	30.0	19.5	10.5	409	14.0
1961	965,000	30,009	31.1	6,344	6.6	19	63.3	880	29.3	20.0	9.3	370	12.3
1960	951,023*	30,747	32.3	6,503	6.8	13	42.3	1,022	33.2	20.2	13.0	414	13.5
1959	937,000	30,186	32.2	6,174	6.6	16	53.0	994	32.9	22.6	11.3	388	12.9
1958	912,000	28,698	31.5	6,088	6.7	9	31.4	1,073	37.4	22.3	15.0	389	13.6
1957	877,000	27,947	31.9	6,080	6.9	14	50.1	1,093	39.1	23.4	16.0	383	13.7
1956	830,000	26,650	32.1	5,446	6.6	10	37.5	956	35.9	22.0	13.9	405	15.2
1955	787,000	25,723	32.7	5,481	7.0	18	70.0	1,121	43.6	24.3	19.2	415	16.1
1954	767,000	25,381	33.1	5,293	6.9	28	110.3	1,087	42.8	23.8	18.9	404	15.9
1953	752,000	24,897	33.1	5,493	7.3	21	84.3	1,134	45.6	24.1	21.6	421	16.9
1952	719,000	24,059	33.5	5,369	7.5	27	112.2	1,163	48.3	26.9	21.3	417	17.3
1951	701,000	23,447	33.4	5,554	7.9	34	145.0	1,295	55.2	29.1	26.2	383	16.3
1950	681,187*	22,014	32.3	5,438	8.0	36	163.3	1,212	55.1	26.9	28.1	381	17.3

Note: Rates for Maternal Deaths are per 100,000 live births;
Rates for Infant and Fetal Deaths are per 1,000 live births;
Rates for Births and Deaths are per 1,000 population.

* BASED ON CENSUS ENUMERATION FOR APRIL 1, AND REVISED POPULATION ESTIMATES.